PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PC1)


(51) International Patent Classification 7: WO 00/40749 (11) International Publication Number: A2 C12Q 1/68 (43) International Publication Date: 13 July 2000 (13.07.00) (21) International Application Number: PCT/CA00/00005 (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, (22) International Filing Date: 5 January 2000 (05.01.00) ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, (30) Priority Data: SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, 60/115,125 6 January 1999 (06.01.99) US 09/477,148 4 January 2000 (04.01.00) US AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, (71)(72) Applicant and Inventor: LIEW, Choong-Chin [CA/CA]; MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, 81 Millersgrove Drive, Willowdale, Ontario M2R 3S1 (CA). GA, GN, GW, ML, MR, NE, SN, TD, TG). (74) Agent: DEETH WILLIAMS WALL; National Bank Building, **Published** Suite 400, 150 York Street, Toronto, Ontario M5H 3S5 (CA). Without international search report and to be republished upon receipt of that report.

(54) Title: METHOD FOR THE DETECTION OF GENE TRANSCRIPTS IN BLOOD AND USES THEREOF

(57) Abstract

The present invention is directed to detection and measurement of gene transcripts in blood. Specifically provided is a RT-PCR analysis performed on a drop of blood for detecting, diagnosing and monitoring diseases using tissue-specific primers. The present invention also describes methods by which delineation of the sequence and/or quantitation of the expression levels of disease-associated genes allows for an immediate and accurate diagnostic/prognostic test for disease or to assess the effect of a particular treatment regimen.





FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| 1 | | | | | _ | | | |
|------|--------------------------|----|---------------------|----|-----------------------|----|-------------------------------------|--|
| . AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia | |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia | |
| AT | Austria | FR | France | LU | Luxembourg | SN | Senegal | |
| AU | Australia | GA | Gabon | LV | Latvia | SZ | Swaziland | |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad | |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo | |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan | |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav | TM | Turkmenistan | |
| BF | Burkina Faso | GR | Greece | | Republic of Macedonia | TR | Turkey | |
| BG | Bulgaria | HU | Hungary | ML | Mali | TT | Trinidad and Tobago | |
| BJ | Benin | IE | Ireland | MN | Mongolia | UA | Ukraine | |
| BR | Brazil | IL | Israel | MR | Mauritania | UG | Uganda | |
| BY | Belarus | IS | Iceland | MW | Malawi | US | • | |
| CA | Canada | IT | Italy | MX | Mexico | UZ | United States of America Uzbekistan | |
| CF | Central African Republic | JP | Japan | NE | Niger | VN | | |
| CG | Congo | KE | Kenya | NL | Netherlands | YU | Vict Nam | |
| СН | Switzerland | KG | Kyrgyzstan | NO | Norway | | Yugoslavia | |
| CI | Côte d'Ivoire | KP | Democratic People's | NZ | New Zealand | zw | Zimbabwe | |
| CM | Cameroon | | Republic of Korea | PL | Poland | | | |
| CN | China | KR | Republic of Korea | PT | | | | |
| CU | Cuba | KZ | Kazakstan | RO | Portuga! Romania | | | |
| cz | Czech Republic | LC | Saint Lucia | RU | Russian Federation | | | |
| DE | Germany | Li | Liechtenstein | SD | Sudan | | | |
| DK | Denmark | LK | Sri Lanka | SE | | | | |
| EE | Estonia | LR | Liberia | SG | Sweden | | | |
| _ | | | Siocing. | 30 | Singapore | | | |
| | | | | | • | | | |
| | | | | | | | | |

METHOD FOR THE DETECTION OF GENE TRANSCRIPTS IN BLOOD AND USES THEREOF

5

10

15

BACKGROUND OF THE INVENTION

Cross-Reference to Related Application

This application claims the benefit of priority of provisional patent application U.S. Serial Number 60/115,125, filed January 6, 1999 and of a U.S. application entitled "Method for the Detection of Gene Transcripts in Blood and uses Thereof" filed on January 4, 2000 (application number not yet assigned).

Field of the Invention

The present invention relates generally to the molecular biology of human diseases. More specifically, the present invention relates to a process using the genetic information contained in human peripheral whole blood for the diagnosis, prognosis and monitoring of genetic and infectious disease in the human body.

Description of the Related Art

20

25

The blood is a vital part of the human circulatory system for the human body. Numerous cell types make up the blood tissue including monocytes, leukocytes, lymphocytes and erythrocytes. Although many blood cell types have been described, there are likely many as yet undiscovered cell types in the human blood. Some of these undiscovered cells may exist transiently, such as those derived from tissues and organs that are constantly interacting with the circulating blood in health and disease. Thus, the blood can provide an immediate picture of what is happening in the human body at any given time.

The turnover of cells in the hematopoietic system is enormous. It was reported that over one trillion cells, including 200 billion erythrocytes and 70 billion neutrophilic leukocytes, turn over each day in the human body (Ogawa 1993). As a consequence of continuous interactions between the blood and the body, genetic changes that occur within the cells or tissues of the body will trigger specific changes in gene expression within blood. It is the goal of the present invention that these genetic alterations be harnessed for diagnostic and prognostic purposes, which may lead to the development of therapeutics for ameliorating disease.

5

10

15

20

25

The complete profile of gene expression in the circulating blood remains totally unexplored. It is hypothesized that gene expression in the blood is reflective of body state and, as such, the resultant disruption of homeostasis under conditions of disease can be detected through analysis of transcripts differentially expressed in the blood alone. Thus, the identification of several key transcripts or genetic markers in blood will provide information about the genetic state of the cells, tissues, organs and systems of the human body in health and disease.

The prior art is deficient in non-invasive methods of screening for tissue-specific diseases. The present invention fulfills this long-standing need and desire in the art.

SUMMARY OF THE INVENTION

This present invention discloses a process of using the genetic information contained in human peripheral whole blood in the diagnosis, prognosis and monitoring of genetic and infectious disease in the human body. The process described herein requires a simple blood sample and is, therefore, non-invasive compared to conventional practices used to detect tissue specific disease, such as biopsies.

One object of the present invention is to provide a non-invasive method for the diagnosis, prognosis and monitoring of genetic and infectious disease in humans and animals.

In one embodiment of the present invention, there is provided a method for detecting expression of a gene in blood from a subject, comprising the steps of: a) quantifying RNA from a subject blood sample; and b) detecting expression of the gene in the quantified RNA, wherein the expression of the gene in quantified RNA indicates the expression of the gene in the subject blood.

5

10

15

20

25

In another embodiment of the present invention, there is provided a method for detecting expression of one or more genes in blood from a subject, comprising the steps of: a) obtaining a subject blood sample; b) extracting RNA from the blood sample; c) amplifying the RNA; d) generating expressed sequence tags (ESTs) from the amplified RNA product; and e) detecting expression of the genes in the ESTs, wherein the expression of the genes in the ESTs indicates the expression of the genes in the subject blood. Preferably, the genes are tissue-specific genes.

In still another embodiment of the present invention, there is provided a method for detecting expression of one or more genes in blood from a subject, comprising the steps of: a) obtaining a subject blood sample; b) extracting DNA fragments from the blood sample; c) amplifying the DNA fragments; and d) detecting expression of the genes in the amplified DNA product, wherein the expression of the genes in the subject blood.

In yet another embodiment of the present invention, there is provided a method for monitoring a course of a therapeutic treatment in an individual, comprising the steps of: a) obtaining a blood sample from the individual; b) extracting RNA from the blood sample; c) amplifying the RNA; d) generating expressed sequence tags (ESTs) from the amplified RNA product; e) detecting expression of genes in the ESTs, wherein the expression of the genes is associated with the effect of

3

the therapeutic treatment: and f) repeating steps a)-e), wherein the course of the therapeutic treatment is monitored by detecting the change of expression of the genes in the ESTs. Such a method may also be used for monitoring the onset of overt symptoms of a disease, wherein the expression of the genes is associated with the onset of the symptoms.

5

10

15

20

25

In still yet another embodiment of the present invention, there is provided a method for diagnosing a disease in a test subject, comprising the steps of:

a) generating a cDNA library for the disease from a whole blood sample from a normal subject; b) generating expressed sequence tag (EST) profile from the normal subject cDNA library; c) generating a cDNA library for the disease from a whole blood sample from a test subject; d) generating EST profile from the test subject cDNA library; and e) comparing the test subject EST profile to the normal subject EST profile, wherein if the test subject EST profile differs from the normal subject EST profile, the test subject might be diagnosed with the disease.

In still yet another embodiment of the present invention, there is provided a kit for diagnosing, prognosing or predicting a disease, comprising: a) genespecific primers; wherein the primers are designed in such a way that their sequences contain the opposing ends of two adjacent exons for the specific gene with the intron sequence excluded; and b) a carrier, wherein the carrier immobilizes the primer(s). Such a kit may be applied to a test subject whole blood sample to diagnose, prognose or predict a disease.

In yet another embodiment of the present invention, there is provided a kit for diagnosing, prognosing or predicting a disease, comprising: a) probes derived from a whole blood sample for a specific disease; and b) a carrier, wherein the carrier immobilizes the probes. Such a kit may be applied to a test subject whole blood sample to diagnose, prognose or predict a disease.

Furthermore, the present invention provides a cDNA library specific for a disease, wherein the cDNA library is generated from whole blood samples.

Other and further aspects, features, and advantages of the present invention will be apparent from the following description of the presently preferred embodiments of the invention. These embodiments are given for the purpose of disclosure.

5

10

15

20

25

BRIEF DESCRIPTION OF THE DRAWINGS

So that the matter in which the above-recited features, advantages and objects of the invention, as well as others which will become clear, are attained and can be understood in detail, more particular descriptions of the invention briefly summarized above may be had by reference to certain embodiments thereof which are illustrated in the appended drawings. These drawings form a part of the specification. It is to be noted, however, that the appended drawings illustrate preferred embodiments of the invention and therefore are not to be considered limiting in their scope not be considered to limit the scope of the invention.

Figure 1 shows the following RNA samples prepared from human blood; Figure 1A: Lane 1, Molecular weight marker; Lane 2, RT-PCR on APP gene; Lane 3, PCR on APP gene; Lane 4, RT-PCR on APC gene; Lane 5, PCR on APC gene; Figure 1B: Lanes 1 and 2, RT-PCR and PCR of βMyHC, respectively; Lanes 3 and 4, RT-PCR of βMyHC from RNA prepared from human fetal and human adult heart, respectively; Lane 5, Molecular weight marker.

Figure 2 shows quantitative RT-PCR analysis performed on RNA primer (5'samples extracted from a drop of blood. Forward GCCCTCTGGGGACCTGAC-3', SEQ ID No. 1) of exon 1 and reverse primer (5'-CCCACCTGCAGGTCCTCT-3", SEQ ID No. 2) of exons 1 and 2 of insulin gene. Blood samples of 4 normal subjects were assayed. Lanes 1, 3, 5 and 7 represent overnight "fasting" blood sample and lanes 2, 4, 6 and 8 represent "non-fasting" samples.

Figure 3 shows quantitative RT-PCR analysis performed on RNA samples extracted from a drop of blood. Lanes 1 and 2 represent normal healthy person and lane 3 represents late-onset diabetes (Type II) and lane 4 represents asymptomatic diabetes.

Figure 4 shows multiple RT-PCR assay in a drop of blood. Primers were derived from insulin gene (INS), zinc-finger protein gene (ZFP) and house-keeping gene (GADH). Lane 1 represents normal person. Lane 2 represents lateonset diabetes and lane 3 represents asymptomatic diabetes.

Figure 5 shows standardized levels of insulin gene (Figure 5A) and ZFP gene (Figure 5B) expressed in a drop of blood. The first three subjects were normal, second two subjects showed normal glucose tolerance, and the last subject had late onset diabetes type II. Figure 5C shows standardized levels of insulin gene expressed in each fractionated cell from whole blood.

Figure 6 shows the differential screening of human blood cell cDNA library with different cDNA probes of heart and brain tissue. Figure 6A shows blood cell cDNA probes vs. adult heart cDNA probes. Figure 6B shows blood cell cDNA probes vs. human brain cDNA probes.

Figure 7 graphically shows the 1,800 unique genes in human blood and in the human fetal heart grouped into seven cellular functions.

20

25

15

5

10

DETAILED DESCRIPTION OF THE INVENTION

In accordance with the present invention, there may be employed conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. See, e.g., Sambrook, Fritsch & Maniatis, "Molecular Cloning: A Laboratory Manual (1982); "DNA Cloning: A Practical Approach," Volumes I and II (D.N. Glover ed. 1985); "Oligonucleotide Synthesis" (M.J. Gait ed. 1984); "Nucleic Acid

Hybridization" [B.D. Hames & S.J. Higgins eds. (1985)]; "Transcription and Translation" [B.D. Hames & S.J. Higgins eds. (1984)]; "Animal Cell Culture" [R.I. Freshney, ed. (1986)]; "Immobilized Cells And Enzymes" [IRL Press, (1986)]; B. Perbal, "A Practical Guide To Molecular Cloning" (1984). Therefore, if appearing herein, the following terms shall have the definitions set out below.

5

10

15

20

25

A "cDNA" is defined as copy-DNA or complementary-DNA, and is a product of a reverse transcription reaction from an mRNA transcript. "RT-PCR" refers to reverse transcription polymerase chain reaction and results in production of cDNAs that are complementary to the mRNA template(s).

The term "oligonucleotide" is defined as a molecule comprised of two or more deoxyribonucleotides, preferably more than three. Its exact size will depend upon many factors which, in turn, depend upon the ultimate function and use of the oligonucleotide. The term "primer" as used herein refers to an oligonucleotide, whether occurring naturally as in a purified restriction digest or produced synthetically, which is capable of acting as a point of initiation of synthesis when placed under conditions in which synthesis of a primer extension product, which is complementary to a nucleic acid strand, is induced, i.e., in the presence of nucleotides and an inducing agent such as a DNA polymerase and at a suitable temperature and pH. The primer may be either single-stranded or double-stranded and must be sufficiently long to prime the synthesis of the desired extension product in the presence of the inducing agent. The exact length of the primer will depend upon many factors, including temperature, source of primer and the method used. For example, for diagnostic applications, depending on the complexity of the target sequence, the oligonucleotide primer typically contains 15-25 or more nucleotides, although it may contain fewer nucleotides. The factors involved in determining the appropriate length of primer are readily known to one of ordinary skill in the art.

As used herein, random sequence primers refer to a composition of primers of random sequence, i.e. not directed towards a specific sequence. These

sequences possess sufficient complementary to hybridize with a polynucleotide and the primer sequence need not reflect the exact sequence of the template.

"Restriction fragment length polymorphism" refers to variations in DNA sequence detected by variations in the length of DNA fragments generated by restriction endonuclease digestion.

5

10

15

20

25

A standard Northern blot assay can be used to ascertain the relative amounts of mRNA in a cell or tissue obtained from plant or other tissue, in accordance with conventional Northern hybridization techniques known to those persons of ordinary skill in the art. The Northern blot uses a hybridization probe, e.g. radiolabelled cDNA, either containing the full-length, single stranded DNA or a fragment of that DNA sequence at least 20 (preferably at least 30, more preferably at least 50, and most preferably at least 100 consecutive nucleotides in length). The DNA hybridization probe can be labelled by any of the many different methods known to those skilled in this art. The labels most commonly employed for these studies are radioactive elements, enzymes, chemicals which fluoresce when exposed to untraviolet light, and others. A number of fluorescent materials are known and can be utilized as labels. These include, for example, fluorescein, rhodamine, auramine, Texas Red, AMCA blue and Lucifer Yellow. A particular detecting material is antirabbit antibody prepared in goats and conjugated with fluorescein through an isothiocyanate. Proteins can also be labeled with a radioactive element or with an enzyme. The radioactive label can be detected by any of the currently available counting procedures. The preferred isotope may be selected from ³H, ¹⁴C, ³²P, ³⁵S, ³⁶Cl, ⁵¹Cr, ⁵⁷Co, ⁵⁸Co, ⁵⁹Fe, ⁹⁰Y, ¹²⁵I, ¹³¹I, and ¹⁸⁶Re. Enzyme labels are likewise useful, and can be detected by any of the presently utilized colorimetric, spectrophotometric. fluorospectrophotometric, amperometric gasometric techniques. The enzyme is conjugated to the selected particle by reaction with bridging molecules such as carbodiimides, diisocyanates, glutaraldehyde and the like. Many enzymes which can be used in these procedures are known and can be utilized.

The preferred are peroxidase, β -glucuronidase, β -D-glucosidase, β -D-galactosidase, urease, glucose oxidase plus peroxidase and alkaline phosphatase. U.S. Patent Nos. 3,654,090, 3,850,752, and 4,016,043 are referred to by way of example for their disclosure of alternate labeling material and methods.

As used herein, "individual" refers to human subjects as well as non-human subjects. The examples herein are not meant to limit the methodology of the present invention to human subjects only, as the instant methodology is useful in the fields of veterinary medicine, animal sciences and such.

5

10

15

20

25

In one embodiment of the present invention, there is provided a method for detecting expression of a gene in blood from a subject, comprising the steps of: a) quantifying RNA from a subject blood sample; and b) detecting expression of the gene in the quantified RNA, wherein the expression of the gene in quantified RNA indicates the expression of the gene in the subject blood. An example of the quantifying method is by mass spectrometry.

In another embodiment of the present invention, there is provided a method for detecting expression of one or more genes in blood from a subject, comprising the steps of: a) obtaining a subject blood sample; b) extracting RNA from the blood sample; c) amplifying the RNA; d) generating expressed sequence tags (ESTs) from the amplified RNA product; and e) detecting expression of the genes in the ESTs, wherein the expression of the genes in the ESTs indicates the expression of the genes in the subject blood. Preferably, the subject is a fetus, an embryo, a child, an adult or a non-human animal. The genes are non-cancer-associated and tissue-specific genes. Still preferably, the amplification is performed by RT-PCR using random sequence primers or gene-specific primers.

In still another embodiment of the present invention, there is provided a method for detecting expression of one or more genes in blood from a subject, comprising the steps of: a) obtaining a subject blood sample; b) extracting DNA fragments from the blood sample; c) amplifying the DNA fragments; and d) detecting

expression of the genes in the amplified DNA product, wherein the expression of the genes in the amplified DNA product indicates the expression of the genes in the subject blood.

5

10

15

20

25

In yet another embodiment of the present invention, there is provided a method for monitoring a course of a therapeutic treatment in an individual, comprising the steps of: a) obtaining a blood sample from the individual; b) extracting RNA from the blood sample; c) amplifying the RNA; d) generating expressed sequence tags (ESTs) from the amplified RNA product; e) detecting expression of genes in the ESTs, wherein the expression of the genes is associated with the effect of the therapeutic treatment; and f) repeating steps a)-e), wherein the course of the therapeutic treatment is monitored by detecting the change of expression of the genes in the ESTs. Such a method may also be used for monitoring the onset of overt symptoms of a disease, wherein the expression of the genes is associated with the onset of the symptoms. Preferably, the amplification is performed by RT-PCR, and the change of the expression of the genes in the ESTs is monitored by sequencing the ESTs and comparing the resulting sequences at various time points; or by performing single nucleotide polymorphism analysis and detecting the variation of a single nucleotide in the ESTs at various time points.

In still yet another embodiment of the present invention, there is provided a method for diagnosing a disease in a test subject, comprising the steps of:

a) generating a cDNA library for the disease from a whole blood sample from a normal subject; b) generating expressed sequence tag (EST) profile from the normal subject cDNA library; c) generating a cDNA library for the disease from a whole blood sample from a test subject; d) generating EST profile from the test subject cDNA library; and e) comparing the test subject EST profile to the normal subject EST profile, wherein if the test subject EST profile differs from the normal subject EST profile, the test subject might be diagnosed with the disease.

10

In still yet another embodiment of the present invention, there is provided a kit for diagnosing, prognosing or predicting a disease, comprising: a) genespecific primers; wherein the primers are designed in such a way that their sequences contain the opposing ends of two adjacent exons for the specific gene with the intron sequence excluded; and b) a carrier, wherein the carrier immobilizes the primer(s). Preferably, the gene-specific primers are selected from the group consisting of insulinspecific primers, atrial natriuretic factor-specific primers, zinc finger protein genespecific primers, beta-myosin heavy chain gene-specific primers, amyloid precurser protein gene-specific primers, and adenomatous polyposis-coli protein gene-specific primers. Further preferably, the gene-specific primers are selected from the group consisting of SEQ ID Nos. 1 and 2; and SEQ ID Nos. 5 and 6. Such a kit may be applied to a test subject whole blood sample to diagnose, prognose or predict a disease by detecting the quantitative expression levels of specific genes associated with the disease in the test subject and then comparing to the levels of same genes expressed in a normal subject. Such a kit may also be used for monitoring a course of therapeutic treatment or monitoring the onset of overt symptoms of a disease.

5

10

15

20

25

In yet another embodiment of the present invention, there is provided a kit for diagnosing, prognosing or predicting a disease, comprising: a) probes derived from a whole blood sample for a specific disease; and b) a carrier, wherein the carrier immobilizes the probes. Such a kit may be applied to a test subject whole blood sample to diagnose, prognose or predict a disease by detecting the quantitative expression levels of specific genes associated with the disease in the test subject and then comparing to the levels of same genes expressed in a normal subject. Such a kit may also be used for monitoring a course of therapeutic treatment or monitoring the onset of overt symptoms of a disease.

Furthermore, the present invention provides a cDNA library specific for a disease, wherein the cDNA library is generated from whole blood samples.

11

The following examples are given for the purpose of illustrating various embodiments of the invention and are not meant to limit the present invention in any fashion.

5

10

15

EXAMPLE 1

Construction of a cDNA library

RNA extracted from human tissues (including fetal heart, adult heart, liver, brain, prostate gland and whole blood) were used to construct unidirectional cDNA libraries. The first mammalian heart cDNA library was constructed as early as 1982. Since then, the methodology has been revised and optimal conditions have been developed for construction of human heart and hematopoietic progenitor cDNA libraries (Liew et al., 1984; Liew 1993, Claudio et al., 1998). Most of the novel genes which were identified by sequence annotation can now be obtained as full length transcripts.

EXAMPLE 2

Catalogue of blood cell ESTs

20 Random partial sequencing of expressed sequence tags (ESTs) of cDNA clones from the blood cell library was carried out to establish an EST database of blood. The known genes as derived from the ESTs were categorized into seven

major cellular functions (Hwang, Dempsey et al., 1997).

5

10

15

20

25

EXAMPLE 3

Differential screening of cDNA library

cDNA probes generated from transcripts of each tissue were used to hybridize the blood cell cDNA clones (Liew et al., 1997). The "positive" signals which were hybridized with ³²P-labelled cDNA probes were defined as genes which shared identity with blood and respective tissues. The "negative" spots which were not exposed to ³²P-labelled cDNA probes were considered to be blood-cell-enriched or low frequency transcripts.

EXAMPLE 4

Reverse transcriptase-polymerase chain reaction (RT-PCR) assay

RNA extracted from samples of human tissue was used for RT-PCR analysis (Jin et al. 1990). Three pairs of forward and reverse primers were designed for human cardiac beta-myosin heavy chain gene (βMyHC), amyloid precurser protein (APP) gene and adenomatous polyposis-coli protein (APC) gene. The PCR products were also subjected to automated DNA sequencing to verify the sequences as derived from the specific transcripts of blood.

EXAMPLE 5

Detection of tissue specific gene expression in human blood using RT-PCR

The beta-myosin heavy chain gene (βMyHC) transcript (mRNA) is known to be highly expressed in ventricles of the human heart. This sarcomeric protein is important for heart muscle contraction and its presence would not be expected in other non-muscle tissues and blood. In 1990, the gene for human cardiac

βMyHC was completely sequenced (Liew et al. 1990) and was comprised of 4 exons and 42 introns.

The method of reverse transcription polymerase chain reaction (RT-PCR) was used to determine whether this cardiac specific mRNA is also present in human blood. A pair of primers was designed; the forward primer (SEQ ID No. 3) was on the boundary of exons 21 and 22, and the reverse primer (SEQ ID No. 4) was on the boundary of exons 24 and 25. This region of mRNA is only present in β MyHC and is not found in the alpha-myosin heavy chain gene (α MyHC).

A blood sample was first treated with lysing buffer and then undergone centrifuge. The resulting pellets were further processed with RT-PCR. RT-PCR was performed using the total blood cell RNA as a template. A nested PCR product was generated and used for sequencing. The sequencing results were subjected to BLAST and the identity of exons 21 to 25 was confirmed to be from βMyHC (Figure 1A).

Using the same method just described, two other tissue specific genes - amyloid precursor protein (APP, forward primer, SEQ ID No. 7; reverse primer, SEQ ID No. 8) found in the brain and associated with Alzheimer's disease, and adenomatous polyposis coli protein (APC) found in the colon and rectum and associated with colorectal cancer (Groden *et al.* 1991; Santoro and Groden 1997) - were also detected in the RNA extracted from human blood (Figure 1B).

20

25

15

5

10

EXAMPLE 6

Multiple RT-PCR analysis on a drop of blood from a normal/diseased individual

A drop of blood was extracted to obtain RNA to carry out quantitative RT-PCR analysis. Specific primers for the insulin gene were designed: forward primer (5'-GCCCTCTGGGGACCTGAC-3', SEQ ID No. 1) of exon 1 and reverse primer (5'-CCCACCTGCAGGTCCTCT-3", SEQ ID No. 2) of exons 1 and 2 of insulin gene. Such reverse primer was obtained by deleting the intron between the

exons 1 and 2. Blood samples of 4 normal subjects were assayed. It was found that the insulin gene is expressed in the blood and the quantitative expression of the insulin gene in a drop of blood is influenced by fasting and non-fasting states of normal healthy subjects (Figure 2). This very low level of expression of the insulin gene reflects the phenotypic status of a person and strongly suggests that there is a physiological and pathological role for its expression, contrary to the basal or illegitimate theory of transcription suggested by Chelly *et al.* (1989) and Kimoto (1998).

5

10

15

20

25

Same quantitative RT-PCR analysis was performed using insulin specific primers on RNA samples extracted from a drop of blood from a normal healthy person, a person having late-onset diabetes (Type II) and a person having asymptomatic diabetes. It was found that the insulin gene is expressed differentially amongst subjects that are healthy, diagnosed as type II diabetic, and also in an asymptomatic preclinical patient (Figure 3).

Similarly, specific primers for the atrial natriuretic factor (ANF) gene were designed (forward primer, SEQ ID No. 5; reverse primer, SEQ ID No. 6) and RT-PCR analysis was performed on a drop of blood. ANF is known to be highly expressed in heart tissue biopsies and in the plasma of heart failure patients. However, atrial natriuretic factor was observed to be expressed in the blood and the expression of the atrial natriuretic factor gene is significantly higher in the blood of patients with heart failure as compared to the blood of a normal control patient.

Specific primers for the zinc finger protein gene (ZFP, forward primer, SEQ ID No. 9; reverse primer, SEQ ID No. 10) were also designed and RT-PCR analysis was performed on a drop of blood. ZFP is known to be high in heart tissue biopsies of cardiac hypertrophy and heart failure patients. In the present study, the expression of ZFP was observed in the blood as well as differential expression levels of ZFP amongst the normal, diabetic and asymptomatic preclinical subjects (Figure 4); although neither of the non-normal subjects has been specifically diagnosed as

suffering from cardiac hypertrophy and/or heart failure, the higher expression levels of the ZFP gene in their blood may indicate that these subjects are headed in that general direction.

It was hypothesized that a housekeeping gene such as glyceraldehyde dehydrogenase (GADH) which is required and highly expressed in all cells would not be differentially expressed in the blood of normal vs. disease subjects. This hypothesis was confirmed by RT-PCR using GADH specific primers (Figure 4). Thus, GADH is useful as an internal control.

Standardized levels of insulin gene or ZFP gene expressed in a drop of blood were estimated using a housekeeping gene as an internal control relative to insulin or ZFP expressed (Figures 5A & 5B). The levels of insulin gene expressed in each fractionated cell from whole blood were also standardized and shown in Figure 5C.

15 EXAMPLE 7

Human blood cell cDNA library

5

10

20

25

In order to further substantiate the present invention, differential screening of the human blood cell cDNA library was conducted. cDNA probes derived from human blood, adult heart or brain were respectively hybridized to the human blood cDNA library clones. As shown in Figure 7, more than 95% of the "positively" identified clones are identical between the blood and other tissue samples.

DNA sequencing of randomly selected clones from the human whole blood cell cDNA library was also performed. This allowed information regarding the cellular function of blood to be obtained concurrently with gene identification. More than 20,000 expressed sequence tags (ESTs) have been generated and characterized to date, 17.6% of which did not result in a statistically significant match to entries in the

GenBank databases and thus were designated as "Novel" ESTs. These results are summarized in Figure 7 together with the seven cellular functions related to percent distribution of known genes in blood and in the fetal heart.

From 20,000 ESTs, 1,800 have been identified as known genes which may not all appear in the hemapoietic system. For example, the insulin gene and the atrial natriuretic factor gene have not been detected in these 20,000 ESTs but their transcripts were detected in a drop of blood, strongly suggesting that all transcripts of the human genome can be detected by performing RT-PCR analysis on a drop of blood.

In addition, approximately 400 novel genes have been identified from the 20,000 ESTs characterized to date, and these will be subjected to full length sequencing and open reading frame alignment to reduce the actual number of novel ESTs prior to screening for disease markers.

Analysis of the approximately 6,283 ESTs which have known matches in the GenBank databases revealed that this dataset represents over 1,800 unique genes. These genes have been catalogued into seven cellular functions. Comparisons of this set of unique genes with ESTs derived from human brain, heart, lung and kidney demonstrated a greater than 50% overlap in expression (Table 1).

20 <u>TABLE 1</u>

5

10

15

25

Overlap of Genes Expressed in Blood *

| Tissues | ESTs** | Overlap in Blood |
|---------|---------|------------------|
| brain | 134,000 | 60% |
| heart | 65,000 | 59% |
| lung | 60,200 | 58% |
| kidney | 32,300 | 54% |

* Estimated from limited known genes of about 1,800 as derived from the database of 6,297 ESTs from human blood cell library.

** Obtained from the National Centre of Biotechnology Information (NCBI), U.S.A.

5

10

15

20

EXAMPLE 8

Blood cell ESTs

The results from the differential screening clearly indicate that the transcripts expressed in the whole blood are reflective of genes expressed in all cells and tissues of the body. More than 95% of detectable spots were identical from two different tissues. The remaining 5% of spots may represent cell- or tissue-specific transcripts; however, results obtained from partial sequencing to generate ESTs of these clones revealed most of them not to be cell- or tissue-specific transcripts. Therefore, the negative spots are postulated to be reflective of low abundance transcripts in the tissue from which the cDNA probes were derived.

An alternative approach that was employed to identify transcripts expressed at low levels is the large-scale generation of expressed sequence tags (ESTs). There is substantial evidence regarding the efficiency of this technology to detect previously characterized (known) and uncharacterized (unknown or novel) genes expressed in the cardiovascular system (Hwang & Dempsey *et al.*. 1997). In the present invention, 20,000 ESTs have been produced from a human blood cell cDNA library and resulted in the identification of approximately 1,800 unique known genes (Table 2)

25

In the most recent GenBank release, analysis of more than 300,000 ESTs in the database (dbESTs) generated more than 48,000 gene clusters which are thought to represent approximately 50% of the genes in the human genome. Only 4,800 of the dbESTs are blood-derived. In the present invention, 20,000 ESTs have

been obtained to date from a human blood cDNA library, which provides the world's most informative database with respect to blood cell transcripts. From the limited amount of information generated so far (i.e. 1,800 unique genes), it has already been determined that more than 50% of the transcripts are found in other cells or tissues of the human body (Table 2). Thus, it is expected that by increasing the number of ESTs generated, more genes will be identified that have an overlap in expression between the blood and other tissues. Furthermore, the transcripts for several genes which are known to have tissue-restricted patterns of expression (i.e. βMyHC, APP, APC, ANF, ZFP) have also been demonstrated to be present in blood.

5

10

15

20

Most recently, a cDNA library of human hematopoietic progenitor stem cells has also been constructed. From the limited set of 1,000 ESTs, there are at least 200 known genes that are shared with other tissue related genes (Claudio *et al.* 1998).

Table 2 demonstrates the expression of known genes of specific tissues in blood cells. Previously, only the presence of "housekeeping" genes would have been expected. Additionally, the presence of at least 25 of the currently known 500 genes corresponding to molecular drug targets was detected. These molecular drug targets are used in the treatment of a variety of diseases which involve inflammation, renal and cardiovascular function, neoplastic disease, immunomodulation and viral infection (Drews & Ryser, 1997). It is expected that additional novel ESTs will represent future molecular drug targets.

TABLE 2

Comparison of 1,800 Unique Genes Identified in the Blood Cell cDNA Library to Genes Previously Identified in Specific Tissues

| Gene Identification | No. of ESTs | Accession No. | Tissue Distribution | | | | | | | |
|---|----------------|------------------|---------------------|--------------|----------------|-----------------|--|----------|--------------|--|
| | | | ВГ | l Br | ТН | TK | Li | Lu | <u> </u> | |
| 100 kDa coactivator | 2 | U22055 | - | + | H | `` | | + | 1 | |
| 10kD protein (BC10) | 2 | AF053470 | <u> </u> | + | + | - | + | + | | |
| 14-3-3 epsilon | 2 | U54778 | | + | + | ╁ | 1 | + | _ | |
| 14-3-3 protein | 11 | U28964 | | + | - | - | + | <u> </u> | | |
| 15 kDa selenoprotein (SEP15) | 1 | AF051894 | | + | + | <u> </u> | | + | | |
| 1-phosphatidylinositol-4- phosphate 5-kinase isoform C | 1 | S78798 | | | | | | | | |
| 23 kD highly basic protein | 21 | X56932 | + | + | + | + | + | + | | |
| 2-5A-dependent RNase | 1 | L10381 | | | - | ├ | - | - | | |
| 2'-5'oligoadenylate synthetase 2 (OAS2) | 4 | M87284 | В | | | | _ | | | |
| 26S proteasome subunit 11 | 1 | AF086708 | | 1 | | † | | l | | |
| 36 kDa phosphothyrosine protein | 2 | AJ223280 | T | | + | | | | | |
| 3-7 gene product (non- exact 86%aa) | 1 | D64159 | | | | | | | | |
| 3-phosphoglycerate dehydrogenase (PGAD) | 1 | AF006043 | Т | + | + | | | + | | |
| 3-prime-phosphoadenosine 5-prime-phosphosulfate synthase 1 (PAPSS1) | 2 | U53447 | + | + | + | + | | + | | |
| 46kd mannose 6- phosphate receptor (MPR46) (low match) | . 1 | X56257 | | | | | | | | |
| 5-aminoimidazole-4- carboxamide ribonucleotide transformylase | 1 | D89976 | | | | | | | | |
| 5'-nucleotidase | 3 | D38524 | I | + | | | + | | | |
| 6-phosphofructo-2- kinase/fructose-2,6- biphosphatase 4 (PFKFB4) | 1 | D49818 | | + | | | | | | |
| 6-phosphofructo-2- kinase/fructose-2,6- bisphosphatase (PF2K) | 1 1 | AF041829 | | | | | | | | |
| 71 kd heat shock cognate protein hsc70 | 23 | Y00371 | | | | | | | | |
| 76 kDa membrane protein (P76) | 2 | U81006 | | + | + | + | + | + | | |
| 8-oxoguanine DNA glycosylase (OGG1) | 1 | U96710 | В | | | | + | + | | |
| a disintegrin and metalloprotease domain 10 (ADAM10) | 1 | AF009615 | Τ | | | | + | | | |
| a disintegrin and metalloprotease domain 8 (ADAM8) | 1 | D26579 | В | + | | | | | | |
| A kinase anchor protein 95 (AKAP95) | 2 | Y11997 | B, T activated | | + | | | + | | |
| A kinase anchor protein, 149kD (AKAP149) | 2 | X97335 | | + | + | + | | + | | |

| CT 7- PW | | | | | | | | | |
|---|--------------|----------|-----------|---|---|----|----|--------|------------------------|
| A4 differentiation- dependent protein (A4), triple LIM domain protein (LMO6), and synaptophysin (SYP); calcium channel alpha-1 | | U93305 | | | | | | | |
| subunit (CACNA1F) ABL and putative M8604 | 1 | U07561 | | | - | + | ╀ | +- | |
| Met protein Absent in melanoma 1 | 1 | U83115 | + | + | - | ╀ | +- | + | |
| (AlM1) accessory proteins | | Z31696 | | + | + | +- | ╄ | ┼ | - |
| BAP31/BAP29 (DXS1357E) | | | | | | | | | |
| acetyl-Coenzyme A acyltransferase (peroxisomal 3-oxoacyl- Coenzyme A thiolase) (ACAA) | 2 | X12966 | + | + | + | + | + | + | |
| acetyl-Coenzyme A transporter (ACATN) | 1 | D88152 | Tlymphoma | + | + | | | | |
| acidic 82 kDa protein | 4 | U15552 | | | | | | | |
| acidic protein rich in leucines (SSP29) | 1 | Y07969 | В | + | + | | + | + | |
| Aconitase 2, mitochondrial (ACO2) | 1 | U80040 | + | + | + | + | | + | |
| actin binding protein MAYVEN | 1 | AF059569 | | | | | | | · |
| actin, beta (ACTB) | 158 | X04098 | Т, В | + | + | | + | | - |
| actin, beta (ACTB) (non- exact, low match 73%) | 1 | M10277 | | | | | | | |
| actin, gamma (low score) | 1 | K00791 | | | | | | | |
| actin, gamma 1 (ACTG1) | 4 | X04098 | + | + | + | + | + | + | high in many libraries |
| actin-binding LIM protein (ABLIM) | 4 | D31883 | | + | + | + | | + | |
| Actinin, alpha 1 (ACTN1) | 8 | M95178 | | + | + | + | | + | |
| actinin, alpha 4 (ACTN4) | 1 | D89980 | | + | + | | + | | |
| activated p21cdc42Hs kinase (ACK) | 1 | L13738 | В | + | | | | + | |
| activated RNA polymerase II transcription cofactor 4 (PC4) | 1 | X79805 | + | + | + | + | | + | |
| activating transcription factor 1 (ATF1) | 1 | X55544 | | | + | | | | |
| activating transcription factor 2 (ATF2) | 1 | X15875 | | + | + | | + | | |
| activating transcription factor 4 (tax-responsive enhancer element B67) (ATF4) | 2 | M86842 | | | | | + | + | |
| active BCR-related gene (ABR) | 1 | U01147 | + | + | + | + | | + | |
| acyl-CoA oxidase (AOX) | 1 | U03254 | | | | | | | |
| acyl-Coenzyme A dehydrogenase, C-4 to C- 12 straight chain (ACADM) | 2 | M16827 | | | | | | | |
| acyl-Coenzyme A dehydrogenase, very long chain (ACADVL) | 3 | D43682 | + | + | + | + | + | + | |
| acyloxyacyl hydrolase (neutrophil) (AOAH) | 3 | M62840 | Т | | + | | + | + | |
| adaptin, delta (ADTD) | 2 | U91930 | | + | + | | + | | |
| adaptin, delta (ADTD) (non-exact 59%) | 1 | AC005328 | | | | | | | |
| adaptin, gamma (ADTG) | 1 | Y12226 | | + | + | + | | + | |
| adaptor complex sigma3B (AP3S3) | 2 | X99459 | | + | | + | | + | |
| adaptor protein p150 | 1 | Y08991 | | | | | | | |
| adducin 1 (alpha) (ADD1) | 2 | L07261 | | + | + | | + | \neg | |

| adducin 1 (alpha) (add1) | 3 | L29296 | + | + | + | + | Т | + | - | |
|---|----------|----------|--------------|----------|--------------|----------|---|----------|---------------|-----------------|
| adducin 3 (gamma) (ADD3) | 3 | U37122 | B, W | + | + | ╁ | + | + | | |
| adenine nucleotide | 2 | M57424 | | + | + | + | + | ╁ | | |
| translocator 2 (fibroblast) (ANT2) | | | | | | | | | | |
| adenine nucleotide translocator 2 (fibroblast) | 1 | J02683 | | | | 1 | | T | 1 | |
| (ANT2) (non-exact 81%) | | | | | | | | | | |
| adenine nucleotide | 1 | J02683 | <u> </u> | | | | | | | |
| translocator 2 (fibroblast) (ANT2) (non-exact, 79%) | <u>L</u> | | | | | | ļ | | | |
| adenine nucleotide translocator 2 (fibroblast) | 1 | J02683 | | | | | | | | |
| (ANT2) (non-exact, 86%) | | | İ | | | | | 1 | 1 | |
| adenine nucleotide translocator 3 (liver) | 3 | J03592 | | + | + | | + | + | | |
| (ANT3) | | | | Ì | | [| İ | | | |
| adenosine deaminase, RNA-specific (ADAR) | 6 | U18121 | | + | + | | + | | | |
| adenylate cyclase 3 | 2 | AF033861 | | + | + | + | + | + | | |
| (ADCY3) adenylate cyclase 7 | 1 | D25538 | | <u> </u> | | _ | | | | |
| (ADCY7) | | | | | İ | | | | | |
| adenylate kinase 2 (AK2) | 2 | U39945 | | + | + | | + | + | | |
| adenylate kinase 3 (AK3) (non-exact, 67%) | 1 | X60673 | | | | | | | | |
| adenylyl cyclase- associated protein (CAP) | 28 | M98474 | T T | | + | | + | | | |
| adipose differentiation- | 1 | X97324 | | _ | + | - | + | + | | |
| related protein; adipophilin (ADFP) | | | | | | | | | | |
| ADP-ribosylation factor 1 (ARF1) | 13 | M84326 | | + | + | | + | + | | |
| ADP-ribosylation factor 3 (ARF3) | 2 | M33384 | | + | + | | + | | | |
| ADP-ribosylation factor 4 (ARF4) | 1 | M36341 | Tlymphoma | + | + | | | + | | |
| ADP-ribosylation factor 5 (ARF5) | 1 | M57567 | | | + | + | + | + | | |
| ADP-ribosylation factor domain protein 1, 64kD (ARFD1) | 1 | L04510 | | + | | | | | | |
| ADP-ribosyltransferase (NAD+; poly (ADP-ribose) polymerase) (ADPRT) | 4 | M32721 | + | + | + | + | + | + | | |
| adrenergic, beta, receptor kinase 1 (ADRBK1) | 2 | X61157 | В | + | | | + | | | |
| adrenoleukodystrophy-like 1 (ALDL1) | 1 | AJ000327 | | | | | | | | |
| AE-binding protein 1 (AEBP1) (non-exact, 62%) | 1 | D86479 | | | | | | | | |
| AF-17 | 1 | U07932 | | | | | | | | |
| A-gamma-globin | 1 | V00514 | | | | | - | | | |
| A-gamma-globin | 1 | J00176 | | | | | | \dashv | | |
| (chromosome 11 allele) | 1 | U78027 | | | | | | | | |
| tyrosine kinase (ATK) | | | | | | | | 1 | | |
| ÁHNAK nucleoprotein (desmoyokin) (AHNAK) | 4 | M80899 | + | + | + | + | | + | | |
| alanyl (membrane) aminopeptidase | 1 | X13276 | | | + | | + | | | |
| (aminopeptidase N, | | | | | | | | | | |
| aminopeptidase M, | | | | | | | ' | | | |
| aminopeptidase, CD13, p150) (ANPEP) | | | | | | | • | | | |
| alcohol dehydrogenase 5 (class III), chi polypeptide (ADH5) | 1 | M29872 | | | | | | | | |
| aldehyde dehydrogenase | 1 | AF003341 | | + | | \dashv | + | + | | |
| 1, soluble (ALDH1) | | ! | | | | | | | | |

| aldehyde dehydrogenase 10 (fatty aldehyde dehydrogenase) (ALDH10) | 2 | U75286 | | | | | | | 100 |
|--|---|-------------|---|---|---|----------|---|---|-----|
| aldehyde reductase 1 (low Km aldose reductase) (ALDR1) | 3 | J04795 | В | + | + | + | + | | |
| aldo-keto reductase family 1, member A1 (aldehyde reductase) (AKR1A1) | 2 | J04794 | В | + | + | | + | | |
| aldo-keto reductase family 1, member C3 (3-alpha hydroxysteroid | 1 | D17793 | | + | + | + | | + | |
| dehydrogenase, type II) (AKR1C3) | | | | | | | | | |
| aldo-keto reductase family 7, member A2 (aflatoxin aldehyde reductase) (AKR7A2) | 1 | Y16675 | | + | + | | + | + | |
| aldolase A, fructose- bisphosphate (ALDOA) | 7 | X12447 | | + | + | | + | | |
| aldolase C, fructose- bisphosphate (ALDOC) | 2 | X05196 | | + | + | | + | | |
| alkaline phosphatase, liver/bone/kidney (ALPL) | 1 | 4502062 | | | | | | | |
| ALL-1 (=L04731;L04284 HRX) | 4 | Z69780 | | | | | | | |
| alpha mannosidase II isozyme | 1 | D55649 | | + | | | + | | |
| alpha thalassemia/mental retardation syndrome X- linked (ATRX) | 3 | U75653 | + | + | + | + | | + | |
| alpha-2 macroglobulin | 1 | Z11711 | | | | | | | |
| alpha-2-globin | 2 | V00516 | | | | | | | |
| alpha-2-macroglobulin receptor/lipoprotein receptor protein (A2MR/LRP) | 1 | U06985 | | | | | | | |
| alpha-polypeptide of N- acetyl-alpha- glucosaminidase (HEXA) | 1 | M13520 | | | | | | | |
| alpha-spectrin | 1 | X86901 | | 1 | | | | | |
| alpha-subunit of Gi2 a (GTP-binding signal transduction protein) | 1 | X07854 | | | | | | | |
| aminin receptor 1 (67kD); Ribosomal protein SA (LAMR1) | 2 | J03799 | T | + | + | | + | + | |
| aminolevulinate, delta-, dehydratase (ALAD) | 1 | X54467 | | + | | | | | |
| amino-terminal enhancer of split (AES) | 2 | X73358 | + | + | + | + | | + | |
| amino-terminal enhancer of split (AES) | 3 | U04241 | В | + | + | | + | + | |
| AMP deaminase isoform L (AMPD2) | 8 | M91029 | | + | | | | + | |
| amphiphysin (Stiff-Mann syndrome with breast cancer 128kD autoantigen) (AMPH) | 1 | U07616 | В | + | | | | + | |
| amphiphysin (Stiff-Mann syndrome with breast cancer 128kD autoantigen) (AMPH)(non-exact, 68%) | 1 | U07616 | | 7 | | | | | |
| amphiphysin (Stiff-Mann syndrome with breast cancer 128kD autoantigen) (AMPH)(non-exact, 68%) | 1 | U07616 | | | | | | | |
| amphiphysin II | 4 | U87558 | | + | + | \dashv | + | | |
| amphiphysin II (67%aa amphiphysin?) | 1 | AF068915 | | | | \dashv | | | |
| amphiphysin II (non-exact 69% aa) | 1 | AF001383 | | | | | | | |
| | | | | | | | | | |

| amphiphysin-like (AMPHL) | 1 | U68485 | | + | T + | T | П | Τ | |
|--|--------------|----------|--|---|----------|--------------|----|--|---------------------------------------|
| amphiphysin-like (AMPHL) | 1 | AF068918 | | | | <u> </u> | | | |
| (low match) AMY-1 | | D50692 | В. Т | | - | ┼ | + | - | |
| amyloid beta (A4) | 1 | L77864 | | + | + | + | ⊢ | + | |
| precursor protein-binding, family B, member 1 (Fe65) (APBB1) | | | | | | | | | |
| amyloid beta (A4) precursor-like protein 2 (APLP2) | 6 | L27631 | Tlymphoma | + | + | | + | + | |
| ankyrin 3, node of Ranvier (ankyrin G) (ANK) (non- exact, 50%) | 1 | U43965 | | | | | | | |
| annexin I (lipocortin I) (ANX1) | 1 | X05908 | | + | + | + | | + | |
| annexin II | 1 ··- | D28364 | | | <u> </u> | t | - | | |
| annexin II (lipocortin II; calpactin I, heavy polypeptide) (ANX2) | 7 | D00017 | + | + | + | + | + | + | high in many libraries |
| annexin IV (placental anticoagulant protein II) (ANX4) | 1 | M19383 | | + | + | + | + | + | |
| annexin V (endonexin II) (ANX5) | 2 | M21731 | | + | + | + | | + | |
| annexin V (endonexin II) (ANXV) | 1 | M19384 | | + | + | + | | + | · |
| annexin VI (p68) (ANX6) | 6 | Y00097 | | + | + | + | | + | · · · · · · · · · · · · · · · · · · · |
| annexin VII (synexin) (ANX7) | 1 | J04543 | | + | + | + | | + | |
| antigen identified by monoclonal antibodies 12E7, F21 and O13 (MIC2) | 2 | M16279 | | + | + | + | | + | |
| antigen identified by monoclonal antibodies 4F2, TRA1.10, TROP4, and T43 (MDU1) | 3 | J02939 | | + | + | + | + | + | |
| antigen TQ1 | 1 | | | | - | | ┢─ | | |
| anti-oxidant protein 2 (non- selenium glutathione peroxidase, acidic calcium- independent phospholipase A2) (KIAA0106) | 1 | D14662 | | + | + | + | + | + | |
| APEX nuclease (multifunctional DNA repair enzyme) (APEX) | 5 | X66133 | | + | + | | + | + | |
| Apolipoprotein L (APOL) (59%aa) | 1 | Z82215 | | | | | | | |
| apoptosis inhibitor 1 (API1) | 1 | L49431 | | + | + | + | + | + | |
| apoptosis inhibitor 4 | 1 | U75285 | B, W | + | + | | + | | |
| (survivin) (API4) apoptosis inhibitor 5 (API5) | 1 | U83857 | Tlymphoma | + | | | + | | |
| apoptosis specific protein (ASP) | <u>.</u> 1 | Y11588 | В | + | | | + | + | |
| apoptotic protease activating factor (APAF1) | 1 | AF013263 | В | + | + | | + | | |
| aquaporin 3 (AQP3) | <u>1</u> | AB001325 | | | | | + | | |
| aquaporin 9 (AQP9) | 7 | AB008775 | Tactivated | | | | + | - | |
| arachidonate 12- lipoxygenase (ALOX12) | 1 | M58704 | T | | | | + | + | |
| arachidonate 5- lipoxygenase-activating protein (ALOX5AP) | 3 | X52195 | + | + | | + | | + | |
| ariadne homolog (ARI) | 1 | AJ009771 | + | + | + | + | | + | |
| ariadne-2 (D. melanogaster) homolog (all-trans retinoic acid inducible RING finger) | 1 | AF099149 | + | + | + | + | | + | |
| (ARI2) | | | | | | | | | |

| ARP1 (actin-related protein 1, yeast) homolog A (centractin alpha) | 1 | X82206 | | + | | | + | | | | | |
|--|-------------|-----------|---------------------|---|----------|--------------|---|----------|--------------|---|---|----------|
| (ACTR1A) | | | | | | 1 | | 1 | ļ | | | |
| ARP2 (actin-related protein 2, yeast) homolog (ACTR2) | 9 | AF006082 | | + | + | | + | + | | | | |
| ARP2/3 protein compex subunit 34 (ARC34) | 5 | AF006085 | Tactivated, W | + | + | | + | | | | | _ |
| Arp2/3 protein compex subunit p41 (ARC41) | 6 | AF006084 | monocyte stimulated | + | + | | + | | | | | |
| Arp2/3 protein compex subunit p41 (ARC41)) (low match) | 1 | AF006084 | | | | | | | | | | |
| Arp2/3 protein complex subunit p16 (ARC16) | 20 | AF017807 | | + | + | | + | + | | | | _ |
| Arp2/3 protein complex subunit p20 (ARC20) | 2 | AF006087 | | + | + | <u> </u> | + | + | | | | _ |
| Arp2/3 protein complex subunit p21(ARC21) | 3 | AF006086 | w | | | t | + | + | | | | _ |
| ARP3 (actin-related protein 3, yeast) homolog (ACTR3) | 11 | AF006083 | w | | + | 一 | + | + | | | | - |
| arrestin, beta 2 (ARRB2) | 1 | AF106941 | B, T, W | + | + | ├ | + | - | | | | - |
| arsA (bacterial) arsenite | | AF047469 | B. T | + | <u> </u> | | + | <u> </u> | | | | _ |
| transporter, ATP-binding, homolog 1 (ASNA1) | • | 71 047403 | 5, 1 | Ť | | | * | | | | | |
| aryl hydrocarbon receptor nuclear translocator-like (ARNTL) | 2 | AF044288 | В | + | + | | + | | | | | _ |
| aryl hydrocarbon receptor- interacting protein (AIP) | 1 | U31913 | + | + | + | + | | + | | | | \dashv |
| arylsulfatase A (ARSA) | 1 | X52151 | Tactivated | + | | | + | \vdash | | | | \dashv |
| asialoglycoprotein receptor 2 (ASGR2) | 1 | M11025 | | | | - | + | + | | | | ᅱ |
| asparaginyl-tRNA synthetase (NARS) | 3 | D84273 | | + | + | - | + | _ | | | | ᅦ |
| aspartyl-tRNA synthetase (DARS) | 1 | J05032 | В | + | + | - | + | | | | | ㅓ |
| ataxia telangiectasia mutated (includes complementation groups A, C and D) (ATM) | 1 | U82828 | В, Т | | + | | + | | | | <u>- · </u> | |
| ataxin-2-like protein A2LP (A2LG) | 1 | AF034373 | B, T activated | + | + | | | + | | | | ᅦ |
| ATF6 | 1 | AF005887 | - GOLLVERCO | + | | | + | | | | | ┥ |
| ATP binding cassette transporter (ABCR) (non-exact 80%) | 1 | U88667 | | | | | | | | | | \dashv |
| ATP synthase (F1-ATPase) alpha subunit, | 1 - | X59066 | | | | | | | | - | · · · · · · · · · · · · · · · · · · · | \dashv |
| mitochondrial | | | | | | | | | | | | 1 |
| ATP synthase beta subunit gene | 1 | M19482 | | | | | | | | | | ٦ |
| ATP synthase, H+ transporting, mitochondrial F0 complex, subunit b, isoform 1 (ATP5F1) | 1 | X60221 | + | + | + | + | | + | | | | |
| ATP synthase, H+ transporting, mitochondrial F0 complex, subunit c (subunit 9), isoform 1 (ATP5G1) | 1 | X69907 | Tactivated | + | + | | + | + | | | | |
| ATP synthase, H+ transporting, mitochondrial F1 complex, alpha subunit, isoform 1, cardiac muscle (ATP5A1) | 3 | D14710 | | | | | | | | | | |
| ATP synthase, H+ transporting, mitochondrial F1 complex, alpha subunit, isoform 1, cardiac muscle (ATP5A1) (low match) | 1 | D14710 | | | | | | | | | | |

| | | | | | | | | • | C1/CA00/00003 |
|--|--------------------------------------|--|------------------------|----------|----------|-----|----------|--|----------------|
| ATP synthase, H+ | 2 | M27132 | | П | | T | | | · · |
| transporting, mitochondrial | ŀ | ļ | | | 1 | | 1 | | 1 |
| F1 complex, beta | | Ĭ | | J | | | Ì | ł | 1 |
| polypeptide (ATP5B) | | Ì | • | ì | | | | 1 | |
| ATP synthase, H+ | 1 | D16563 | W | + | + | + | + | | |
| transporting, mitochondrial | | | 1 | | 1 | | | | 1 |
| F1 complex, gamma | | | į. | 1 | 1 | ì | 1 | | l i |
| polypeptide 1 (ATP5C1) | 1 | | ŀ | 1 | 1 | 1 | 1 | | 1 |
| ATP synthase, H+ | 1 - | AF092124 | + | + | + | + | + | + | |
| transporting, mitochondrial | ' | 71 032 124 | T | * | * | | * | T | |
| F1F0, subunit g (ATP5JG) | ŀ | 1 | | | 1 | | l | | 1 |
| | | I I TO BE DO A | | | L | | <u> </u> | | |
| ATP/GTP-binding protein | 2 | U73524 | + | + | + | + | 1 | + | |
| (HEAB) | | | <u> </u> | ſ | 1 | | 1 | | 1 |
| ATPase, Ca++ | 5 | Z69881 | | + | T | | | | |
| transporting, ubiquitous | J | | | i | 1 | i i | 1 | | 1 |
| (ATP2A3) | 1 | | | ľ | 1 | 1 | i ' | 1 | 1 |
| ATPase, H+ transporting, | 2 | D89052 | + | + | + | + | | + | |
| lysosomal (vacuolar proton | _ | | | 1 | | i | 1 | i . | |
| pump) 21kD (ATP6F) | | | | i | ļ | | 1 | | ; |
| ATPase, H+ transporting, | 1 | X76228 | | + | + | + | _ | | |
| lysosomal (vacuolar proton | i ' | 170220 | | T | T | T | | + | |
| pump) 31kD (ATP6E) | ļ | • | İ | ļ | | Ī | | | |
| ATPase, H+ transporting, | 5 | VCO4E4 | | <u> </u> | ├ | ٠. | | | <u> </u> |
| lysosomal (vacuolar proton | i ³ | X69151 | | + | + | +_ | | + | |
| | I | | 1 | l | 1 | 1 | | l | į į |
| pump) 42kD; Vacuolar proton-ATPase. | | | | 1 | 1 | 1 | | | |
| | | | | | | 1 | | | 1 |
| subunit C; V-ATPase, | | } | | J |] | | 1 1 | | i l |
| subunit C (ATP6D) | | <u> </u> | | | 1 | ļ | ł i | 1 | · |
| ATPase, H+ transporting, | 3 | L09235 | | + | | + | | | |
| lysosomal (vacuolar proton | | | | | ľ | | | | 1 |
| pump), alpha polypeptide, | | • | } | | | 1 1 | i i | | ! |
| 70kD, isoform 1 (ATP6A1) | | | 1 | 1 | | | | | f I |
| ATPase, H+ transporting. | 6 | X62949 | + | + | + | + | - | + | |
| lysosomal (vacuolar proton | _ | | ł | ' | ' | 1 | | • | |
| pump), beta polypeptide, | | | | Į | l | | | |] |
| 56/58kD, isoform 2 | | | | İ | | 1 | | | |
| (ATP6B2) | | | | | ļ | | | | |
| ATPase, H+ transporting, | 2 | AF038954 | | | <u> </u> | | | | ļ |
| lysosomal (vacuolar proton | | AF030934 | + | + | + | + | | + | high in testis |
| pump), member J (ATP6J) | | | | l | 1 |] | | | l |
| ATPase, H+ transporting, | | | | L . | L | | | | |
| | 1 | D16469 | } | + " | + | + | | + | |
| lysosomal (vacuolar proton | | | i i | | İ | | | | |
| pump), subunit 1 (ATP6S1) | | | <u> </u> | l | l | | | | l |
| ATP-binding cassette 50 | 1 | AF027302 | + | + | + | + | | + | |
| (TNF-alpha stimulated) | | } | | | l | 1 | | | |
| (ABC50) | | j | | | ļ | | - 1 | | ! |
| ATP-binding cassette | 1 | AF047690 | | | | _ | \dashv | | |
| protein M-ABC1 | | | 1 | | | | | | l i |
| (mitochondrial) | | 1 | | | i i | | | | |
| ATP-dependent RNA | | | | | | 1 1 | 1 | | |
| | 7 | I A INTORAN | Lymphoma | | | | | | |
| | 1 | AJ010840 | Tlymphoma | | + | | + | | i |
| helicase | | | | | | | + | | |
| helicase autoantigen (Hs.75528) | 2 | L05425 | T lymphoma T activated | | + | | + | | |
| helicase autoantigen (Hs.75528) autoantigen (Hs.75528) | | | | | | | + | | |
| helicase autoantigen (Hs.75528) autoantigen (Hs.75528) (non-exact 84%) | | L05425 L05425 | | | | | + | | |
| helicase autoantigen (Hs.75528) autoantigen (Hs.75528) (non-exact 84%) | | L05425 | | + | | | + | + | |
| helicase autoantigen (Hs.75528) autoantigen (Hs.75528) (non-exact 84%) autoantigen (Hs.75682) | 1 | L05425 L05425 U17474 | T activated | + | | | + | + | |
| helicase autoantigen (Hs.75528) autoantigen (Hs.75528) (non-exact 84%) autoantigen (Hs.75682) autoantigen La/SS-B | 1 | L05425 L05425 U17474 Z35127 | T activated | + | | | + | + | |
| helicase autoantigen (Hs.75528) autoantigen (Hs.75528) (non-exact 84%) autoantigen (Hs.75682) | 1 | L05425 L05425 U17474 | T activated | + | | | + | + | |
| helicase autoantigen (Hs.75528) autoantigen (Hs.75528) (non-exact 84%) autoantigen (Hs.75682) autoantigen La/SS-B axin (AXIN1) | 1 1 1 | L05425 L05425 U17474 Z35127 AF009674 | T activated | | | | + | | |
| helicase autoantigen (Hs.75528) autoantigen (Hs.75528) (non-exact 84%) autoantigen (Hs.75682) autoantigen La/SS-B axin (AXIN1) axonemal dynein heavy | 2 1 1 | L05425 L05425 U17474 Z35127 | T activated | | | | • | + | |
| helicase autoantigen (Hs.75528) autoantigen (Hs.75528) (non-exact 84%) autoantigen (Hs.75682) autoantigen La/SS-B axin (AXIN1) axonemal dynein heavy chain (DNAH17) | 1 1 1 1 | L05425 L05425 U17474 Z35127 AF009674 AJ000522 | T activated | | | | + | | |
| helicase autoantigen (Hs.75528) autoantigen (Hs.75528) (non-exact 84%) autoantigen (Hs.75682) autoantigen La/SS-B axin (AXIN1) axonemal dynein heavy chain (DNAH17) BAI1-associated protein 3 | 1 1 1 | L05425 L05425 U17474 Z35127 AF009674 | T activated | | | | • | | |
| helicase autoantigen (Hs.75528) autoantigen (Hs.75528) (non-exact 84%) autoantigen (Hs.75682) autoantigen La/SS-B axin (AXIN1) axonemal dynein heavy chain (DNAH17) BAI1-associated protein 3 (BAIAP3) (non-exact 54%) | 1 1 1 1 | L05425 L05425 U17474 Z35127 AF009674 AJ000522 AB017111 | T activated | | | | • | | |
| helicase autoantigen (Hs.75528) autoantigen (Hs.75528) (non-exact 84%) autoantigen (Hs.75682) autoantigen (Hs.75682) autoantigen La/SS-B axin (AXIN1) axonemal dynein heavy chain (DNAH17) BAI1-associated protein 3 (BAIAP3) (non-exact 54%) basement membrane- | 1 1 1 1 | L05425 L05425 U17474 Z35127 AF009674 AJ000522 | T activated | | | | • | | |
| helicase autoantigen (Hs.75528) autoantigen (Hs.75528) (non-exact 84%) autoantigen (Hs.75682) autoantigen (Hs.75682) autoantigen La/SS-B axin (AXIN1) axonemal dynein heavy chain (DNAH17) BAI1-associated protein 3 (BAIAP3) (non-exact 54%) basement membrane- induced gene (ICB1) | 1 1 1 1 1 1 | L05425 L05425 U17474 Z35127 AF009674 AJ000522 AB017111 AF044896 | T activated | | | | • | | |
| helicase autoantigen (Hs.75528) autoantigen (Hs.75528) (non-exact 84%) autoantigen (Hs.75682) autoantigen (Hs.75682) autoantigen La/SS-B axin (AXIN1) axonemal dynein heavy chain (DNAH17) BAI1-associated protein 3 (BAIAP3) (non-exact 54%) basement membrane- induced gene (ICB1) basic leucine zipper | 1 1 1 1 | L05425 L05425 U17474 Z35127 AF009674 AJ000522 AB017111 | T activated | | | | + | | |
| helicase autoantigen (Hs.75528) autoantigen (Hs.75528) (non-exact 84%) autoantigen (Hs.75682) autoantigen (Hs.75682) autoantigen La/SS-B axin (AXIN1) axonemal dynein heavy chain (DNAH17) BAI1-associated protein 3 (BAIAP3) (non-exact 54%) basement membrane- induced gene (ICB1) basic leucine zipper nuclear factor 1 (JEM-1) | 1 1 1 1 1 1 | L05425 L05425 U17474 Z35127 AF009674 AJ000522 AB017111 AF044896 | T activated | | | | + | | |
| helicase autoantigen (Hs.75528) autoantigen (Hs.75528) (non-exact 84%) autoantigen (Hs.75682) autoantigen (Hs.75682) autoantigen La/SS-B axin (AXIN1) axonemal dynein heavy chain (DNAH17) BAI1-associated protein 3 (BAIAP3) (non-exact 54%) basement membrane- induced gene (ICB1) basic leucine zipper nuclear factor 1 (JEM-1) | 1 1 1 1 1 1 | L05425 L05425 U17474 Z35127 AF009674 AJ000522 AB017111 AF044896 | T activated | | | | + | | |
| helicase autoantigen (Hs.75528) autoantigen (Hs.75528) (non-exact 84%) autoantigen (Hs.75682) autoantigen (Hs.75682) autoantigen La/SS-B axin (AXIN1) axonemal dynein heavy chain (DNAH17) BAI1-associated protein 3 (BAIAP3) (non-exact 54%) basement membrane- induced gene (ICB1) basic leucine zipper nuclear factor 1 (JEM-1) (BLZF1) | 2 1 1 1 1 1 1 | L05425 L05425 U17474 Z35127 AF009674 AJ000522 AB017111 AF044896 U79751 | T activated B | + | + | | | + | |
| helicase autoantigen (Hs.75528) autoantigen (Hs.75528) (non-exact 84%) autoantigen (Hs.75682) autoantigen (Hs.75682) autoantigen La/SS-B axin (AXIN1) axonemal dynein heavy chain (DNAH17) BAI1-associated protein 3 (BAIAP3) (non-exact 54%) basement membrane- induced gene (ICB1) basic leucine zipper nuclear factor 1 (JEM-1) (BLZF1) basic transcription factor 3 | 1 1 1 1 1 1 | L05425 L05425 U17474 Z35127 AF009674 AJ000522 AB017111 AF044896 | T activated | | | + | + | | |
| helicase autoantigen (Hs.75528) autoantigen (Hs.75528) (non-exact 84%) autoantigen (Hs.75682) autoantigen (Hs.75682) autoantigen La/SS-B axin (AXIN1) axonemal dynein heavy chain (DNAH17) BAI1-associated protein 3 (BAIAP3) (non-exact 54%) basement membrane- induced gene (ICB1) basic leucine zipper nuclear factor 1 (JEM-1) (BLZF1) basic transcription factor 3 (BTF3) | 2 1 1 1 1 1 1 2 | L05425 L05425 U17474 Z35127 AF009674 AJ000522 AB017111 AF044896 U79751 | T activated B | + | + | + | + | + | |
| helicase autoantigen (Hs.75528) autoantigen (Hs.75528) (non-exact 84%) autoantigen (Hs.75682) autoantigen (Hs.75682) autoantigen La/SS-B axin (AXIN1) axonemal dynein heavy chain (DNAH17) BAI1-associated protein 3 (BAIAP3) (non-exact 54%) basement membrane- induced gene (ICB1) basic leucine zipper nuclear factor 1 (JEM-1) (BLZF1) basic transcription factor 3 (BTF3) basigin (BSG) | 2 1 1 1 1 1 1 2 | L05425 L05425 U17474 Z35127 AF009674 AJ000522 AB017111 AF044896 U79751 X74070 L10240 | T activated B | + | + | + | | + | |
| helicase autoantigen (Hs.75528) autoantigen (Hs.75528) (non-exact 84%) autoantigen (Hs.75682) autoantigen (Hs.75682) autoantigen La/SS-B axin (AXIN1) axonemal dynein heavy chain (DNAH17) BAI1-associated protein 3 (BAIAP3) (non-exact 54%) basement membrane- induced gene (ICB1) basic leucine zipper nuclear factor 1 (JEM-1) (BLZF1) basic transcription factor 3 (BTF3) | 2 1 1 1 1 1 1 2 | L05425 L05425 U17474 Z35127 AF009674 AJ000522 AB017111 AF044896 U79751 | T activated B | + | + | + | + | + | |

| WO 00/40/49 | | | | | | | | _ | 01.01100.0000 |
|---|----------|----------|-------------|---|---|----------|---|--|---------------------------------|
| B-cell CLU/lymphoma 6 (zinc finger protein 51) | 1 | U00115 | | + | + | | | | :• |
| (BCL6) B-cell translocation gene 1. | 1 | X61123 | | 4 | + | | | + | |
| anti-proliferative (BTG) | . | | <u></u> | 1 | _ | | | • | |
| BCL2/adenovirus E1B 19kD-interacting protein 2 (BNIP2) | 1 | U15173 | В | + | | | + | + | |
| BCL2/adenovirus E1B 19kD-interacting protein 3- like (BNIP3L) | 2 | AF067396 | | + | + | + | | + | |
| beclin 1 (coiled-coil, myosin-like BCL2- interacting protein) (BECN1) | 1 | AF077301 | В | + | + | | + | | |
| beta-1,2-N- acetylglucosaminyltransfer ase II (MGAT2) | 2 | U15128 | | | | | | | |
| beta-2-microglobulin (B2M) | 63 | S82297 | + | + | + | + | + | + | high in invasive prostate tumor |
| beta-hexosaminidase alpha chain (HEXA) | 1 | M16411 | | | _ | | | | prostate tunior |
| beta-tubulin | 7 | V00599 | + | + | + | + | + | + | high in many libraries |
| beta-tubulin (non-exact, 76%) | 1 | AF070561 | | | | | | | |
| beta-tubulin, pseudogene | 1 | J00315 | | 1 | | - | | | |
| BING4 | 1 | Z97184 | | + | | | | _ | |
| biotinidase (BTD) (non-eact 62%) | 1 | U03274 | | | | | | | |
| biotinidase (BTD) (non- exact 70%) | 1 | U03274 | - | 1 | | | | | |
| biotinidase (BTD) (non- exact, 56%) | 1 | U03274 | | | | | | - | |
| BIOTINIDASE PRECURSOR | 1 | P43251 | | | | | | | |
| biphenyl hydrolase-like (serine hydrolase) (BPHL) | 1 | X81372 | | + | | | + | | |
| bone marrow stromal cell antigen 1 (BST1) | 1 | D21878 | | | - | | + | | |
| box-dependent myc- interacting protein isoform BIN1-10 (BIN1) | 1 | AF043900 | | | | | | | |
| box-dependent myc- interacting protein isoform BIN1-10 (BIN1) (non-exact, 64%) | 1 | AF043900 | | | | | | | |
| brain my047 protein | 1 | AF063605 | T | + | + | | + | | |
| branched chain keto acid dehydrogenase E1, alpha polypeptide (maple syrup urine disease) (BCKDHA) | 3 | Z14093 | 7 | + | + | | + | | |
| BRCA1 associated protein- 1 (ubiquitin carboxy- terminal hydrolase) (BAP1) | 1 | D87462 | + | + | + | + | | | |
| BRCA1, Rho7 and vatl genes, and ipf35 | 1 | L78833 | | | | | | | |
| breakpoint cluster region protein, uterine leiomyoma, 1; barrier to autointegration factor (BCRP1) | 2 | AF044773 | | + | + | | | | |
| breakpoint cluster region protein, uterine leiomyoma, 2 (BCRP2) | 2 | AF044774 | | + | + | | + | + | |
| breast cancer anti-estrogen resistance 3 (BCAR3) (non-exact 73%) bromodomain-containing | 1 | U92715 | | | | | | | |
| protein, 140kD (peregrin) (BR140) | 2 | M91585 | | + | | | | | |
| Bruton's agammaglobulinemia tyrosine kinase (Btk) | 1 | U13424 | | | | | | | |

| Bruton's tyrosine kinase (BTK) | 1 | U78027 | | | | | | | |
|---|-----|------------------|-------------|----------|---|---|---|---|---|
| Bruton's tyrosine kinase (BTK), alpha-D- galactosidase A (GLA), L44-like ribosomal protein (L44L) and FTP3 (FTP3) | 1 . | U78027 | | | | | | | |
| BS4 | 1 | AF108083 | | | | | | | |
| BTG2 (BTG2) | 6 | Y09943 | + | + | + | + | | + | |
| BTK region clone ftp | 1 | U78027 | + | + | + | + | | + | |
| BTK region clone ftp-3 | 1 | U01923 | | + | + | | + | L | |
| BUB3 (budding uninhibited by benzimidazoles 3, yeast) homolog (BUB3) | 4 | AF053304 | + | + | + | + | | + | |
| butyrate response factor 1 (EGF-response factor 1) (BRF1) | 4 | X79067 | + | + | + | + | | + | |
| butyrophilin (BTF1) | 7 | U90543 | | + | + | | + | | |
| butyrophilin like receptor | 1 | AB020625.1 | | | | | | | |
| CAG repeat containing (CTG4A) CAGH32 | 2 | U80744 | | + | + | | | | |
| calcium channel, voltage- | 1 | U80743 M83566 | | + | + | | + | | |
| dependent, L type, alpha 1D subunit (CACNA1D) (low match) | • | 10105500 | | | | | | | · |
| calcium/calmodulin- dependent protein kinase (CaM kinase) II gamma (CAMK2G) | 1 | AF069765 | | + | + | + | | + | |
| calcium/calmodulin- dependent protein kinase kinase (KIAA0787) | | AF101264 | В | + | + | | + | | |
| calmodulin (=M19311) | 7 | D45887 | | <u> </u> | | | | | |
| calmodulin 1 (phosphorylase kinase, delta) (CALM1) | 6 | M27319 | В | + | + | | + | + | |
| calnexin (CANX) | 3 | M94859 | l | + | | | + | + | |
| calpain, large polypeptide L1 (CAPN1) | 5 | X04366 | | + | + | | + | + | |
| calpain, large polypeptide L2 (CANP2) | 5 | M23254 | | + | + | | | | |
| calpain, small polypeptide (CAPN4) | 1 | X04106 | | + | + | | + | + | |
| calpastatin (CAST) | 3 | D16217 | | | | | + | | |
| Calponin 2 | 2 | D83735 | | + | | + | | + | |
| calponin 2 (CNN2) | 1 | D83735 | В, Т | + | | | + | | |
| calponin 2 (CNN2) (low score) | 1 | D83735 | | | | | | | |
| calumenin (CALU) | 3 | AF013759 | В | | + | | + | + | |
| cAMP response element- binding protein CRE-Bpa (H_GS165L15.1) | 4 | L05912 | | | | | | | |
| cAMP-dependent protein kinase type II (Ht31) | 1 | M90360 | | | | | | | |
| canicular multispecific organic anion transporter (CMOAT2) | 1 | AF009670 | | | | + | + | + | |
| capping protein (actin filament) muscle Z-line, alpha 1 (CAPZA1) | 6 | U56637 | В, Т | | + | | | + | |
| capping protein (actin filament) muscle Z-line, alpha 2 (CAPZA2) | 2 | U03269 | В | + | + | | | | |
| capping protein (actin filament) muscle Z-line, beta (CAPZB) | 1 | U03271 | + | + | + | + | | + | |

| | | | | | | | | • | C1/CA00/00003 |
|--|-----|------------|--------------------|---------|--------|--|----------|--|---------------|
| capping protein (actin filament), gelsolin-like (CAPG) | 8 | M94345 | + | + | | + | | + | |
| carbamoyl-phosphate synthetase 2, aspartate transcarbamylase, and dihydroorotase (CAD) | 1 . | D78586 | + | + | + | + | | + | |
| carbonic anhydrase V, mitochondrial (CA5) | 1 | L19297 | | + | | | + | | |
| carboxypeptidase D (CPD) | 3 | U65090 | В | + | + | +- | \vdash | ├─ | |
| carnitine/acylcarnitine translocase (CACT) | 1 | Y10319 | | + | + | | + | | |
| Cas-Br-M (murine) | 2 | X57110 | | 1 | | | + | - | |
| transforming sequence (cbl) | | | | | | | | | |
| casein kinase 1, alpha 1 (CSNK1A1) | 1 | L37042 | + | + | + | + | | + | |
| casein kinase 2, alpha 1 polypeptide (CSNK2A1) | 2 | M55265 | В | + | | | + | + | |
| casein kinase I gamma 3L (CSNK1G3L) | 1 | AF049090.1 | | | | | | | |
| casein kinase II alpha subunit(=S72393) | 1 | X69951 | | | | | | | |
| CASP8 and FADD-like apoptosis regulator (CFLAR) | 4 | AF015450 | | + | + | + | + | + | |
| caspase 1, apoptosis- related cysteine protease (interleukin 1, beta, convertase) (CASP1) | 7 | U13697 | + | | | + | | | |
| caspase 10, apoptosis- related cysteine proteas (CASP10) | 1 | U60519 | B, T acti lymph | | Τ | | + | | |
| caspase 3, apoptosis- related cysteine protease (CASP3) | 3 | U13737 | B, T | + | + | + | + | | |
| caspase 4, apoptosis- related cysteine protease (CASP4) | 6 | U25804 | + | + | + | + | | + | |
| caspase 5, apoptosis- related cysteine protease (CASP5) | 1 | U28015 | | | + | ļ | | | |
| caspase 8, apoptosis- related cysteine protease (CASP8) | 2 | X98173 | | + | | + | | + | |
| caspase 9, apoptosis- related cysteine protease (CASP9) | 1 | U56390 | В | | | + | + | | |
| catalase (CAT) | 5 | X04076 | В | + | + | | + | | |
| catechol-O- methyltransferase (COMT) | 1 | M65213 | | + | + | | + | | |
| catenin (cadherin- associated protein), alpha 1 (102kD) (CTNNA1) | 6 | D14705 | | + | + | | | | |
| cathelicidin antimicrobial peptide (CAMP) | 1 | X89658 | В | | | | | | |
| cathepsin B (CTSB) | 4 | L16510 | | | + | | + | + | |
| cathepsin C (CTSC) | 3 | U79415 | | + | + | + | | + | |
| cathepsin D (lysosomal aspartyl protease) (CTSD) | 4 | M11233 | | + | + | | + | | |
| cathepsin E (CTSE) | 1 | J05036 | | | | | + | | |
| cathepsin G (CTSG) | 1 | M16117 | T, W | | + | | | | |
| cathepsin S (CTSS) | 34 | M86553 | B, Monocyt lym | e stimi | ulated | , T | + | + | |
| cathepsin W (lymphopain) (CTSW) | 4 | AF013611 | | | | | | + | |
| CBF1 interacting corepressor CIR (=U03644 recepin) | 1 | AF098297 | | | | | | | |

| | | _ | | | | | | | |
|--|-----|----------|------------|---|---|----------|----------|---|-------------|
| CCAAT/enhancer binding protein (C/EBP), alpha (CEBPA) | 3 | X87248 | | + | + | + | | + | |
| CCAAT/enhancer binding protein (C/EBP), delta (CEBPB) | 1 | S63168 | | | + | | + | + | |
| CCAAT-box-binding transcription factor (CBF2) | 2 | M37197 | T lymphoma | | | + | + | | |
| CCR5 receptor (CCR5) (non-exact?) | 1 | AF011504 | | - | | | | _ | |
| CD14 antigen (CD14) | 11 | M86511 | + | + | + | + | | + | |
| CD18 (=M95293) | 4 | X64071 | † | | | \top | <u> </u> | | |
| CD1C antigen, c polypeptide (CD1C) | 2 | M28827 | | | | | | + | |
| CD2 antigen (cytoplasmic tail)-binding protein 2 (CD2BP2) | 1 | AF104222 | | | | | | | |
| CD2 antigen (p50), sheep red blood cell receptor (CD2) | 4 | M14362 | + | | + | + | | + | |
| CD2 cytoplasmic tail- binding protein 1 (CD2BP1) | | AF038602 | | | | | + | | |
| CD20 antigen (CD20) | 1 | X12530 | | | | | | | |
| CD20 receptor (S7) | 1 | X07203 | | | | | | | |
| CD22 antigen (CD22) | 1 | U62631 | В | | | | | | |
| CD24 signal transducer | 1 | M58664 | | | | | | | |
| CD33 antigen (gp67) (CD33) | 1 | M23197 | | | | | + | | |
| CD33 antigen-like 2; OB binding protein-2 (CD33L2) (non-exact, 68%) | 1 | U71383 | | | | | | | |
| CD33L2 (61% aa) | 1 | D86359 | | | | | | | |
| CD36 antigen (collagen type I receptor, thrombospondin receptor) (CD36) | 7 | M98398 | T lymphoma | | + | | + | + | |
| CD37 antigen (CD37) | 5 | X14046 | + | + | | + | - | + | |
| CD38 alt | 1 - | D84277 | | | _ | ┢ | | ├ | |
| CD39 antigen (CD39) | 1 | U87967 | В | + | | \vdash | + | + | |
| CD3D antigen, delta polypeptide (TiT3 complex) (CD3D) | 1 | X03934 | | | + | + | | + | |
| CD3E antigen, epsilon polypeptide (TiT3 complex) (CD3E) | 1 | X03884 | + | | | + | | | |
| CD3G antigen, gamma polypeptide (TiT3 complex) (CD3G) | 2 | X06026 | W | | | | + | | |
| CD3Z antigen, zeta polypeptide (TiT3 complex) (CD3Z) | 2 | J04132 | + | | | + | | | |
| CD3-zeta (clone pBS NK1) | 1 | X55510 | | | | | | | |
| CD4 (low match) | 1 | S68043 | | | | | | | |
| CD4 antigen (p55) (CD4) | 4 | M12807 | | + | + | | + | | |
| CD44 antigen (homing function and Indian blood group system (CD44) | 6 | X56794 | W | | | | + | + | |
| CD48 antigen (B-cell membrane protein) (CD48) | 3 | X06341 | + | + | + | + | | + | |
| CD53 antigen (CD53) | 10 | L11670 | + | + | | + | | + | |
| CD53 antigen (CD53) (low match) | 1 | M60871 | | | | | | | |
| CD63 antigen (melanoma 1 antigen) (CD63) | 3 | M59907 | | | | | | | |
| CD68 antigen (CD68) | 2 | S57235 | | + | + | | + | + | |

WO 00/40749

PCT/CA00/00005

| CD74 antigen (invariant polypeptide of major histocompatibility complex, class II antigen-associated) | 72 | K01144 | + | + | + | + | + | + | high in many libraries |
|---|----|----------|--|--------------|--------------|--|--------------|--------------|------------------------|
| (CD74) | | | | | <u> </u> | <u> </u> | | | |
| CD79A antigen (immunoglobulin- associated alpha) (CD79A) | 2 | M80462 | | | + | | | | |
| CD79B antigen | 2 | M89957 | + | - | | | ├ | ┼ | |
| (immunoglobulin- associated beta) (CD79B) | | Mosso | | | 1 | | | | |
| CD8 antigen, alpha polypeptide (p32) (CD8A) | 2 | M27161 | + | | | + | I^- | + | |
| CD8 antigen, beta polypeptide 1 (p37) | 1 | X13445 | w | | ├─ | +- | <u> </u> | | |
| (CD8B1) | l | | 1 | l | 1 | | | ł | |
| CD81 antigen (target of antiproliferative antibody 1 (CD81) | 1 | M33680 | | + | + | | | + | |
| CD83 antigen (activated B | 1 | Q01151 | В | + | + | ┼── | | + | |
| lymphocytes, mmunoglobulin superfamily) (CD83) | | <u> </u> | | | | | | | |
| CD84 antigen (leukocyte | 1 | U82988 | | + | + | ┼— | | + | |
| antigen) (ČD84) | | | | • | * | 1 | 1 | • | |
| CD86 antigen | 1 | L25259 | 1 | + | <u> </u> | \vdash | ! | | |
| CD9 antigen (p24) (CD9) | 2 | M38690 | | | + | 1 | + | + | |
| CD97 antigen (CD97) | 12 | X84700 | + | + | | + | | | |
| CD97 antigen (CD97) (noin-exact 59%) | 1 | P48960 | | | | | | | |
| CD97 antigen (CD97) (non- exact 62%) | 1 | X94630 | + | + | | + | | | |
| CDC23 (cell division cycle 23, yeast, homolog) (CDC23) | 1 | AF053977 | | + | | | + | + | |
| CDC37 homolog | 1 | U63131 | В | + | + | ┼ | + | + | |
| Cdc42 effector protein 3 (CEP3) | 2 | AF104857 | В | + | + | - | + | | |
| CDC-like kinase (CLK) | 7 | L29219 | | + | + | + | | + | |
| CDC-like kinase 2 (CLK2) | 1 | AF023268 | В | + | + | \vdash | <u> </u> | | |
| CDW52 antigen (CAMPATH-1 antigen) (CDW52) | 13 | X15183 | Tactivated | + | + | | + | | |
| cell cycle progression restoration 8 protein(CPR8) | 1 | AF011794 | | | | | | | |
| cell division cycle 10 (homologous to CDC10 of S. cerevisiae) (CDC10) | 4 | S72008 | + | + | + | + | | + | |
| cell division cycle 20, S.cerevisiae homolog (CDC20) | 1 | U05340 | | + | + | + | | | |
| cell division cycle 25B (CDC25B) | 6 | Z68092 | + | + | + | + | | + | |
| cell division cycle 2-like 1 (PITSLRE proteins) (CDC2L1) (non-exact 42%) | 1 | AF067514 | | | | | | | |
| cell division cycle 42 (GTP- blnding protein, 25kD) (CDC42) | 5 | M35543 | + | + | + | + | | + | |
| cell division protein (non- exact 68%) | 1 | AF063015 | | | | | | | |
| CELL-CYCLE NUCLEAR AUTOANTIGEN SG2NA (S/G2 NUCLEAR ANTIGEN) | 1 | Q13033 | | | | | | | |
| centromere protein B (80kD) (CENPB) | 1 | X55039 | - | + | | | + | | |
| cep250 centrosome associated protein | 3 | AF022655 | В | + | | | + | | |
| associated biotein | | | L | | | | | | |

WO 00/40749

PCT/CA00/00005

| | | | | | | | | _ | C1/CA00/00003 |
|--|---|------------|------------------|----------|----------|--------------|----------|---|---------------|
| ceroid-lipofuscinosis, neuronal 2, late infantile (Jansky-Bielschowsky | 7 | AF017456 | + | + | + | + | + | + | high in bone |
| disease) (CLN2) c-fgr (=M63877 | 6 | X52206 | | ļ | | <u> </u> | ļ | | |
| nonreceptor protein- tyrosine kinase (fgr)) | | A32200 | | | | | | | |
| CGI-19 protein | 3 | AF132953.1 | | <u> </u> | \vdash | | \vdash | _ | |
| chaperonin containing TCP1, subunit 3 (gamma) (CCT3) | 1 | X74801 | | + | .+ | | | + | |
| chaperonin containing TCP1, subunit 4 (delta) (CCT4) | 1 | AF026291 | | + | + | | + | + | |
| chaperonin containing TCP1, subunit 6A (zeta 1) (CCT6A) | 4 | L27706 | В | + | + | | | | |
| chaperonin containing TCP1, subunit 7 (eta) (CCT7) | 4 | AF026292 | В | + | | | | + | |
| Chediák-Higashi syndrome 1 (CHS1) | 1 | U67615 | B, T lymphoma | + | + | | + | | |
| Chediak-Higashi syndrome 1 (CHS1) (low score) | 1 | U67615 | ,,,,,,,,,,,, | | | | | | |
| chemokine (C-C motif) receptor 2 (CCR2) | 4 | U03905 | | | <u> </u> | | | | |
| chemokine (C-C motif) receptor 4 (CCR4) (low match) (may contain repeat) | 1 | X85740 | | | | | | | |
| chemokine (C-C motif) receptor 7 (CCR7) | 6 | L31581 | | | | | _ | | |
| chemokine (C-X3-C) receptor 1 (CX3CR1) | 5 | U20350 | | + | | | | | |
| chemokine (C-X-C motif), receptor 4 (fusin) (CXCR4) | 5 | M99293 | + | + | + | + | | + | |
| chitinase 3-like 1 (cartilage glycoprotein-39) (CHI3L1) | 2 | M80927 | | + | | + | | + | |
| chitinase 3-like 2 (CHI3L2) | 2 | U49835 | | + | | + | | + | |
| chlonde channel 1 , skeletal muscle (CLCN1) | 1 | G18280 | | | | | | | |
| chloride channel 6 (CLCN6) | 1 | D28475 | | + | + | | | | |
| Chloride intracellular channel 1 (CLIC1) | 1 | U93205 | + | + | + | + | | + | |
| chondroitin sulfate proteoglycan 2 (versican) (CSPG2) | 5 | X15998 | | | + | | | | |
| chondroitin sulfate proteoglycan core protein | 2 | J02814 | | | + | | | + | |
| chromatin assembly factor 1 p48 subunit (CAF-1 P48 subunit) (retinoblastoma binding protein p48) (retinoblastoma-binding protein 4) (MSI1 protein homolog) | 1 | Q09028 | | | | - | | | |
| chromodomain helicase DNA binding protein 1 (CHD1) | 2 | AF006513 | | | | | | | |
| chromodomain helicase DNA binding protein 1-like (CHD1L) | 1 | AF054177 | | | | | | | |
| chromodomain helicase DNA binding protein 2 (CHD2) | 1 | AF006514 | В | + | + | | + | | |
| chromodomain helicase DNA binding protein 3 (CHD3) | 1 | AF006515 | | | | | | | |
| chromodomain helicase DNA binding protein 4 (CHD4) | 5 | X86691 | + | + | + | + | | + | |

| reading frame 7 (C1ORF7) | 1 | AF054176 | | | | | | | | | , | • |
|--|---|----------|--------------|---------------------|----------|----------|----------|----------|---------|---|-------------|----------|
| chromosome 1 specific transcript KIAA0493 | 1 | AB007962 | | | | | | | | | | |
| chromosome 17 open reading frame 1B (C17ORF1B) | 1 | AJ008112 | T | + | | | | | | | | |
| chromosome 4 open reading frame 1 (C4ORF1) | 1 | AF006621 | | + | + | + | | + | | | | |
| chromosome condensation 1-like (CHC1L) | 2 | AF060219 | | + | + | + | | + | | | | |
| chromosome X open reading frame 5 (CXORF5) | 1 | Y15164 | В | + | + | | + | | | | | |
| chromosome-associated polypeptide C(CAP-C) | 2 | AF092564 | В | + | + | | + | + | | | | |
| cig42 | 1 | AF026944 | | + | | \vdash | \vdash | | | | | _ |
| cig5 | 3 | AF026941 | - | + | | | - | - | l | | | \dashv |
| citrate synthase (CS) | 2 | AF047042 | В | + | + | | + | + | | | ·· | |
| class I major | 2 | U31372 | | <u> </u> | , | | <u> </u> | <u> </u> | ļ | | | |
| histocompatibility antigen (HLA-Cw3) | _ | | | | | | | | | | | |
| class I major histocompatibility antigen (HLA-Cw3) (low match) | 1 | U31372 | | | | | | | | , | | |
| clathrin assembly protein lymphoid myeloid leukemia (CALM) | 3 | U45976 | В | + | + | | | + | | | | |
| clathrin heavy chain | 1 | X55878 | | | | | _ | | | | | ᅥ |
| clathrin, heavy polypeptide- like 2 (CLTCL2) | 1 | D21260 | | | | | | | | | | |
| clathrin, light polypeptide (Lca) (CLTA) (low match) | 1 | M20472 | | | | | | | | | | |
| clathrin- | 3 | D63475 | | + | + | + | + | + | | | | |
| associated/assembly/adapt or protein, medium 1 (CLAPM1) | | | | | | | | | | | | |
| cleavage stimulation factor, 3' pre-RNA, subunit 2 64kD (CSTF2) (non-exact 82%) | 1 | M85085 | | | | | | | , | | | ٠, |
| cleavage stimulation factor, 3' pre-RNA, subunit 3, 77kD (CSTF3) | 1 | U15782 | В | + | + | | + | | | | | ; |
| cik3 | 1 | L29220 | В | + | + | | | | | | | |
| clone 23815 (Hs.82845) | 1 | U90916 | | + | + | | | + | | | | |
| clone 24592 mRNA sequence | 1 | D88378 | + | + | + | + | | + | | | | 1 |
| Clq/MBL/SPA receptor C1qR(p) () | 1 | U94333 | | | | | | | | | ····· | \dashv |
| clusterin (complement lysis inhibitor, SP-40,40, | 1 | M64722 | + | + | + | + | + | + | | | | \dashv |
| sulfated glycoprotein 2, testosterone-repressed prostate message 2, apolipoprotein J) (CLU) | | | | | | | | : | | | | |
| CMP-sialic acid transporter (CMPST) | 1 | D87969 | В | + | + | | | | | | | ᅦ |
| CMRF35 | 3 | X66171 | | + - 1 | | | | \dashv | | | | \dashv |
| c-myc oncogene containing | 1 | X54629 | | $\dagger = \dagger$ | | | | | | | | \dashv |
| coxIII coagulation factor II | 1 | M62424 | | + | + | | | + | | | | \dashv |
| (thrombin) receptor (F2R) coagulation factor V (proaccelerin, labile factor) | 1 | M14335 | | + | | + | + | | | | · | \dashv |
| (F5) coagulation factor XIII a | 3 | M21998 | | | | | | | | | | - |
| subunit coagulation factor XIII, A1 | 6 | M14354 | | + | + | + | | + | | | | \dashv |
| polypeptide (F13A1) coated vesicle membrane | 1 | X92098 | + | · | <u> </u> | + | + | + | | | | _ |
| protein (RNP24) | | | <u> </u> | \perp | | | | | | | | |

| coatomer protein complex, subunit alpha (COPA) | 5 | U24105 | Т | + | | | + | | 6 |
|---|----|----------|------|---|---|--------------|----------|---|---------------------|
| Cofilin 1 (non-muscle) (CFL1) | 13 | X95404 | + | + | + | + | + | + | high in fetal brain |
| cold inducible RNA-binding protein (CIRBP) | 7 | D78134 | | + | + | | | + | |
| cold shock domain protein A (CSDA) | 3 | X95325 | | + | + | | <u> </u> | | |
| collagen, type IX, alpha 2 (COL9A2) | 3 | AF019406 | В | 1 | | | | | |
| colony stimulating factor 1 receptor, formerly | 3 | X03663 | | + | | ļ | + | + | |
| McDonough feline sarcomal viral (v-fms) oncogene homolog (CSF1R) | | | | | | | | | |
| colony stimulating factor 2 receptor, beta, low-affinity (granulocyte-macrophage) (CSF2RB) | 5 | M59941 | | | | | | | |
| colony stimulating factor 2 receptor, beta, low-affinity (granulocyte-macrophage) (CSF2RB) (low match) | 1 | M59941 | | | | | | | |
| colony stimulating factor 3 receptor (granulocyte) (CSF3R) | 16 | X55720 | | + | | | | | |
| complement component 5 receptor 1 (C5a ligand) (C5R1) | 1 | M62505 | L | | | | | | |
| conserved gene amplified in osteosarcoma (OS4) | 2 | AF000152 | | + | + | + | | + | |
| COP9 (constitutive photomorphogenic, Arabidopsis, homolog) subunit 3 (COPS3) | 2 | AF031647 | | + | + | | | + | |
| COP9 homolog (HCOP9) | 2 | U51205 | В | + | + | + | + | + | |
| COPII protein, homolog of s. cerevisiae SEC23p (SEC23A) | 4 | X97064 | | + | + | | | | |
| copine I (CPNE1) | 2 | U83246 | В | + | + | | + | | |
| copine I (CPNE1) (low score) | 1 | U83246 | | | | | | | |
| coproporphyrinogen oxidase (coproporphyria, harderoporphyria) (CPO) | 1 | D16611 | | | + | | + | + | |
| core-binding factor, beta subunit (CBFB) | 1 | L20298 | | + | | | | | |
| coronin | 22 | X89109 | T, W | + | + | | + | | |
| coronin (low match) | 1 | U34690 | | | | | | | |
| coronin (non-exact, 71%) | 1 | X89109 | | | | | | | |
| cot (cancer Osaka thyroid) oncogene (COT) | 1 | D14497 | + | + | + | + | | + | |
| cryptochrome 1 (photolyase-like) (CRY1) | 1 | D84657 | | + | + | | | + | |
| CTD (carboxy-terminal domain, RNA polymerase II, polypeptide A) phosphatase, subunit 1 (CTDP1) | 1 | AF081287 | | + | + | + | | + | |
| C-terminal binding protein 1 (CTBP1) | 1 | U37408 | В | + | + | | + | | |
| C-terminal binding protein 2 (CTBP2) | 2 | AF016507 | | + | + | | + | | |
| CÙG triplet repeat, RNA- binding protein 1 (CUGBP1) | 3 | U63289 | | + | + | + | | + | |
| cullin 1 (CUL1) | 3 | U58087 | | + | + | + | | + | ` |
| cullin 3 (CUL3) | 2 | U58089 | | + | + | + | | + | |
| cut (Drosophila)-like 1 (CCAAT displacement protein) (CUTL1) | 1 | M74099 | В | + | | | | | |

| cyclin D2 (CCND2) | 2 | D13639 | | + | + | T # | T | + | i i |
|--|---|----------|------------------|---|-----|-----|-----|---|----------|
| cyclin D3 (CCND3) | 5 | M92287 | B, T lymphoma | | + | | + | | |
| cyclin G1 (CNNG1) | 1 | D78341 | В | + | + | | i – | + | |
| cyclin I | 3 | D50310 | В | + | | | + | | <u> </u> |
| cyclin T2 (CNNT2) | 1 | AF048732 | B, T lymphoma | В | | | | | |
| cyclin-dependent kinase 2 (CDK2) | 1 | X62071 | | | | | | | |
| cyclin-dependent kinase inhibitor (p27Kip1) | 1 | 576986 | | | | | | | |
| cyclin-dependent kinase inhibitor 1A (p21, Cip1) (CDKN1A) | 2 | S67388 | + | + | + | + | + | + | |
| CYP2D7-CYP2D6 intergenic region (partial) | 1 | X90926 | | | | | | | |
| cystatin B (stefin B) (CSTB) | 1 | L03558 | | | + | | + | + | |
| cysteine and glycine-rich protein 3 (cardiac LIM protein) (CSRP3) | 5 | L54057 | | | + | | | | |
| cytidine deaminase (CDA) | 2 | L27943 | | | | | + | | |
| cytochrome b | 1 | AF042500 | | | | | Ĺ | | |
| cytochrome b (CYTB) (isolate Aus5) | 1 | AF042518 | | | | | | | |
| cytochrome b(-245) beta chain N-terminal region (X- linked granulomatous disease gene) | 2 | X05895 | | | | | | | |
| cytochrome b-245, beta polypeptide (chronic granulomatous disease) (CYBB) | 2 | X04011 | + | | | + | | + | |
| cytochrome C | 1 | P00001 | | | | | | | |
| cytochrome c oxidase subunit IV (COX4) | 1 | U90915 | 1 | + | + | | + | + | |
| cytochrome c oxidase subunit Vb (COX5B) | 2 | M59250 | | | | | + | | |
| cytochrome c oxidase subunit VII-related protein (COX7RP) | 6 | AB007618 | + | + | + | + | | + | |
| cytokine suppressive anti- inflammatory drug binding protein 1 (p38 MAP kinase) (CSBP1) | 1 | L35263 | lymphocyte | + | + | | + | | |
| Cytoplasmic antiproteinase=38 kda intracellular serine proteinase inhibitor | 1 | S69272 | | | + | | | | |
| cytotoxic granule- associated RNA-binding protein p40-TIA-1 | 1 | S70114 | | | | | | | |
| D123 (D123) | 1 | D14878 | + | + | | + | | + | |
| D2-2 | 1 | AF019226 | | | | | | | |
| D38 | 1 | X74802 | | | | | | | |
| damage-specific DNA binding protein 1 (127kD) (DDB1) | 2 | AJ002955 | + | + | + " | + | + | + | |
| DCHT (low match) | 1 | AF017635 | | | | | | | |
| DEAD/H (Asp-Glu-Ala- Asp/His) box binding protein 1 (DDXBP1) | 1 | U78524 | | + | + | + | + | + | |
| DEAD/H (Asp-Glu-Ala- Asp/His) box polypeptide (72KD) (P72) | 2 | U59321 | T | + | + | | +. | + | |
| DEAD/H (Asp-Glu-Ala- Asp/His) box polypeptide 1 (DDX1) | 1 | X70649 | | + | + | | | + | |

| WO 00/40/49 | | | | | | | | r | C1/CA00/00005 |
|---|-------------|-------------|--|----------|----------|----------|----------|----------|---------------|
| DEAD/H (Asp-Glu-Ala- Asp/His) box polypeptide | 2 | AB001636 | | | | T | | | :- |
| 15 (DDX15) | | | | | <u> </u> | | | | |
| DEAD/H (Asp-Glu-Ala- Asp/His) box polypeptide | 2 | AB011149 | + | + - | + | + | | + | |
| 16 (DDX16) | <u></u> | | | | | | | | |
| DEAD/H (Asp-Glu-Ala- Asp/His) box polypeptide 3 | 3 | U50553 | + | + | + | + | | + | |
| (DDX3) | | | 1 | | ł | | | ł | |
| DEAD/H (Asp-Glu-Ala- | 37 | X15729 | + | + | + | + | | + | |
| Asp/His) box polypeptide 5 (RNA helicase, 68kD) | | | [| ŀ | 1 | ļ | ļ | 1 | |
| (DDX5) | <u>_</u> | | 1 | | 1 | | | l | |
| DEAD/H (Asp-Glu-Ala- Asp/His) box polypeptide 5 | 1 | AF015812 | | | | | | 1 | |
| (RNA helicase, 68kD) | | | | | 1 | 1 | Ì | İ | |
| (DDX5) (low match) | | | | | <u> </u> | | | <u> </u> | |
| DEAD/H (Asp-Glu-Ala- Asp/His) box polypeptide 6 | 2 | D17532 | + | + | | | | | |
| (RNA helicase, 54kD) | | | | | | | | | |
| (DDX6) DEAD/H (Asp-Glu-Ala- | | D50487 | <u> </u> | L. | ļ | | <u> </u> | ļ., | |
| Asp/His) box polypeptide 8 | ' | D50487 | | + | + | + | | + | |
| (RNA helicase, 54kD) | | | | | | ł | | | |
| (DDX8) DEAD/H (Asp-Glu-Ala- | 3 | L13848 | ++ | + | - | + | _ | + | |
| (Asp/His) box polypeptide 9 | | 210040 | | ` | l | ' | | * | |
| (RNA helicase A, nuclear DNA helicase II; | | | | 1 | ļ | İ | | | • |
| leukophysin) (DDX9) | | | | ļ | l | 1 | | ĺ | |
| DEAD/H (Asp-Glu-Ala- | 1 | AF000985 | | + | + | † | + | | |
| Asp/His) box polypeptide, Y chromosome (DBY) | | i | | | ł | | | | |
| Death associated protein 3 | 2 | X83544 | + | + | + | + | + | + | |
| (DAP3) death effector domain- | 1 | AF083236 | | | | | | | |
| containing protein (DEDD) | ' | AF003230 | | + | + | + | | + | |
| death-associated protein 6 (DAXX) | 2 | AF039136 | | + | + | + | | + | |
| dedicator of cyto-kinesis 2 | 4 | D86964 | + | + | | + | | + | |
| (DOCK2) defender against cell death | 1 | D45057 | | | | | | | |
| 1 (DAD1) | <u></u> | D15057 | | | + | | + | + | |
| Defensin, alpha 1, myeloid- related sequence (DEFA1) | 4 | L12690 | | | | + | + | + | |
| DEK gene (D6S231E) | | X64229 | В | - | + | \vdash | + | \vdash | |
| delta sleep inducing | 4 | Z50781 | + | + | + | + | | + | |
| peptide, immunoreactor (DSIPI) | | | | | | | | | |
| dendritic cell protein | 3 | AF064603 | + | + | + | + | | + | |
| deoxycytidine kinase | 1 | M60527 | <u> </u> | | | | | | |
| (DCK) deoxyribonuclease II, | | ABOOAETA | | | | | | | |
| lysosomal (DNASE2) | 3 | AB004574 | | | | | | | |
| ĎGS-I | 2 | L77566 | | + | | | | | |
| diacylglycerol kinase | 3 | D16440 | | | | | | | |
| diacylglycerol kinase alpha | 3 | AF064771 | | + | | \dashv | | | |
| (DAGK1) (clone 24) diacylgiycerol kinase alpha | 1 | AF064771 | | | | | | | |
| (DAGK1) (clone 24) (low match) | • | 711 00 4171 | | | | | | | |
| diaphanous (Drosophila, homolog) 1 (DIAPH1) | 1 | AF051782 | B, monocyte stimulated | + | + | \neg | + | + | |
| diaphorase (NADH) | 1 | Y09501 | + | + | + | + | + | + | |
| (cytochrome b-5 reductase) (DIA1) | | | | l | | | | | |
| differentiated Embryo | 1 | AB004066 | | + | | | + | + | |
| Chondrocyte expressed | Ť | |] | . | | | | | |
| gene 1 (DEC1) | | | <u> </u> | | | | | Ì | |

| differentiated Embryo Chondrocyte expressed gene 1 (DEC1) (low match) differentiation antigen CD20 DiGeorge syndrome critical region gene 2 (DGCR2) dihydrolipoamide dehydrogenase (E3 component of pyruvate dehydrogenase complex, 2-oxo-glutarate complex, branched chain keto acid dehydrogenase complex) (DLD) dihydrolipoamide S- acelyttransferase (E2 component of pyruvate dehydrogenase complex) (DLAT) dihydropyrimidinase-like 2 (DPYSL2) dinG gene 1 Y10571 dipthena toxin resistance protein required for diphthamide biosynthesis (Saccharomyces)-like 2 (DPH2L2) DJ-1 protein Dmx-like 1 (DMXL1) DNA (cytosine-5-)- methyttransferase 1 (DNMT1) DNA fragmentation factor, 40 kD, beta subunit A84076 + 23415 CD3415 CD3415 CD3415 CD3420 + 484076 + | | | + | + | · |
|--|-----|--------|---|----------|------------------------|
| differentiation antigen CD20 DiGeorge syndrome critical region gene 2 (DGCR2) dihydrolipoamide dehydrogenase (E3 component of pyruvate dehydrogenase complex, 2-oxo-glutarate complex, branched chain keto acid dehydrogenase complex) (DLD) dihydrolipoamide S- acetyltransferase (E2 component of pyruvate dehydrogenase complex) (DLAT) dihydrolipoamide S- acetyltransferase (E2 component of pyruvate dehydrogenase complex) (DLAT) dihydropyrimidinase-like 2 dinG gene 1 Y10571 diptheria toxin resistance 3 AF053003 B + protein required for dipthemide biosynthesis (Saccharomyces)-like 2 (DPH2L2) disintegrin-protease (nonexact 72%) DJ-1 protein 2 AF021819 + Dmx-like 1 (DMXL1) 1 AJ005821 + DMA (cytosine-5-)- methyltransferase 1 (DNMT1) DNA fragmentation factor, 40 kD, beta subunit (DFFB) DNA fragmentation factor, 45 kD, alpha subunit | | | + | | |
| DiGeorge syndrome critical region gene 2 (DGCR2) dihydrolipoamide dehydrogenase (E3 component of pyruvate dehydrogenase complex, 2-oxo-glutarate complex, branched chain keto acid dehydrogenase complex) (DLD) dihydrolipoamide S-acetyltransferase (E2 component of pyruvate dehydrogenase complex) (DLAT) dihydropyrimidinase-like 2 1 D78013 + component of pyruvate dehydrogenase complex) (DLAT) dihydropyrimidinase-like 2 1 D78013 + component of pyruvate dehydrogenase complex) (DLAT) dihydropyrimidinase-like 2 1 D78013 + component of pyruvate dehydrogenase complex) (DLAT) dipthena toxin resistance and protein required for dipthena toxin resistance protein required for dipththamide biosynthesis (Saccharomyces)-like 2 (DPH2L2) disintegrin-protease (nonexact 72%) DJ-1 protein 2 AF021819 + component of pyruvate dehydrogenase complex) (DLAT) DNA (cytosine-5-) and AF084019 + component of pyruvate dehydrogenase and pyruvate dehydrogenase complex) (DRAT1) DNA fragmentation factor, 1 AF084019 + component of pyruvate dehydrogenase complex) (DRA fragmentation factor, 2 U91985 T + component of pyruvate dehydrogenase complex) (DRA fragmentation factor, 2 U91985 T + component of pyruvate dehydrogenase complex) (DRA fragmentation factor, 2 U91985 T + component of pyruvate dehydrogenase complex) (DRA fragmentation factor, 2 U91985 T + component of pyruvate dehydrogenase complex) (DRA fragmentation factor, 2 U91985 T + component of pyruvate dehydrogenase complex) (DRA fragmentation factor, 2 U91985 T + component of pyruvate dehydrogenase complex) (DRA fragmentation factor, 2 U91985 T + component of pyruvate dehydrogenase complex) (DRA fragmentation factor, 2 U91985 T + component of pyruvate dehydrogenase complex) (DRA fragmentation factor, 2 U91985 T + component of pyruvate dehydrogenase complex (DRA fragmentation factor, 2 U91985 T + component of pyruvate dehydrogenase (DRA fragmentation factor, 2 U91985 T + component of pyruvate dehydrogenase (DRA fragmentation factor, 2 U91985 T + component of pyruvate dehydrogenase (DRA fragmentati | | | + | | |
| dihydrolipoamide dehydrogenase (E3 component of pyruvate dehydrogenase complex, 2-oxo-glutarate complex, branched chain keto acid dehydrogenase complex) (DLD) dihydrolipoamide S-acetyltransferase (E2 component of pyruvate dehydrogenase complex) (DLAT) dihydropyrimidinase-like 2 1 D78013 + (DPYSL2) dinG gene 1 Y10571 diptheria toxin resistance protein required for diphthamide biosynthesis (Saccharomyces)-like 2 (DPH2L2) disintegrin-protease (non-exact 72%) DJ-1 protein 2 AF021819 + Dmx-like 1 (DMXL1) 1 AJ005821 + DNA (cytosine-5-)-methyltransferase 1 (DNMT1) DNA fragmentation factor, 40 kD, beta subunit (DFFB) DNA fragmentation factor, 45 kD, alpha subunit | | | + | ╁┰ | 1 |
| acetyltransferase (E2 component of pyruvate dehydrogenase complex) (DLAT) dihydropyrimidinase-like 2 | 4 | | | | |
| (DPYSL2) dinG gene | | | + | | |
| dipthena toxin resistance protein required for diphthamide biosynthesis (Saccharomyces)-like 2 (DPH2L2) disintegrin-protease (non-exact 72%) DJ-1 protein 2 AF021819 + + Dmx-like 1 (DMXL1) 1 AJ005821 + DNA (cytosine-5-)- 3 X63692 I activated, hymphoma (DNMT1) DNA fragmentation factor, 40 kD, beta subunit (DFFB) DNA fragmentation factor, 45 kD, alpha subunit | + | | + | + | |
| protein required for diphthamide biosynthesis (Saccharomyces)-like 2 (DPH2L2) disintegrin-protease (nonexact 72%) DJ-1 protein DMA (cytosine-5-) methyltransferase 1 (DMMT1) DNA fragmentation factor, 40 kD, beta subunit (DFFB) DNA fragmentation factor, 45 kD, alpha subunit | | | | \vdash | |
| exact 72%) DJ-1 protein 2 AF021819 + + Dmx-like 1 (DMXL1) 1 AJ005821 + DNA (cytosine-5-)- methyltransferase 1 (DNMT1) DNA fragmentation factor, 40 kD, beta subunit (DFFB) DNA fragmentation factor, 45 kD, alpha subunit | + | | + | + | |
| Dmx-like 1 (DMXL1) DNA (cytosine-5-)- methyltransferase 1 (DNMT1) DNA fragmentation factor, 40 kD, beta subunit (DFFB) DNA fragmentation factor, 2 45 kD, alpha subunit | | | | | 7 |
| DNA (cytosine-5-)- methyltransferase 1 (DNMT1) DNA fragmentation factor, 40 kD, beta subunit (DFFB) DNA fragmentation factor, 45 kD, alpha subunit | + | + | | + | |
| methyltransferase 1 (DNMT1) DNA fragmentation factor, 1 AF064019 40 kD, beta subunit (DFFB) DNA fragmentation factor, 2 U91985 + 45 kD, alpha subunit | + | + | | | |
| 40 kD, beta subunit (DFFB) DNA fragmentation factor, 2 U91985 + + 45 kD, alpha subunit | | | + | + | |
| 45 kD, alpha subunit | | | | | |
| (DFFA) | + | | | + | |
| DNA mismatch repair 1 U17840 protein (hMLH1) | | | | | |
| DNA segment on 3 M64241 + + chromosome X (unique) 648 expressed sequence | + | + | + | + | high in many libraries |
| DNA segment, single copy 3 M73547 + probe LNS-CAI/LNS-CAII (deleted in polyposis (D5S346) | + | + | | + | |
| DNA-damage-inducible 1 L24498 transcript 1 (DDIT1) (low match) | | | | | |
| DnaJ protein 1 AJ001309 | 1 | | | | i . |
| , | 1 | | | | |
| (DOK2) | | \neg | | | |
| diphosphooligosaccharide- protein glycosyltransferase (DDOST) | + | + | + | + | activated T cell |
| mannosyltransferase polypeptide 1, catalytic subunit (DPM1) | + | | + | | |
| activation (immunoglobulin superfamily) (DORA) | | | + | | |
| adenoma DRA (low match) | 1 1 | | ヿ | \dashv | |
| D-type cyclin-interacting 1 AF082569 B protein 1 (DIP1) | | | | + | |
| DnaJ protein 1 AJ001309 docking protein 2, 56kD 1 AF034970 (DOK2) dolichyl- diphosphooligosaccharide- protein glycosyltransferase (DDOST) dolichyl-phosphate 1 D86198 T activated + mannosyltransferase polypeptide 1, catalytic subunit (DPM1) down-regulated by 1 AJ223183 activation (immunoglobulin superfamily) (DORA) down-regulated in 1 P40879 | | + | + | + | activated T cell |

| dual specificity phosphatase 1 (DUSP1) | 4 | X68277 | + | + | + | + | + | + | ř. |
|--|---|----------|---|---|---|---|---|---|----|
| dual specificity phosphatase 11 (RNA/RNP complex 1-interacting) (dusp11) | 1 | AF023917 | + | + | + | + | | + | |
| dual specificity phosphatase 3 (vaccinia virus phosphatase VH1- related) (DUSP3) | 1 | L05147 | | + | + | | + | + | |
| dual specificity phosphatase 6 (DUSP6) | 6 | X93920 | + | + | + | + | + | + | |
| dynactin 1 (p150, Glued (Drosophila) homolog) (DYTN1) | 3 | X98801 | | | | | | | |
| dynactin 1 (p150, Glued (Drosophila) homolog) (DYTN1) (low match) | 1 | X98801 | В | + | + | | | | |
| dynamin 2 (DNM2) | 1 | L36983 | • | | | | | | |
| dynamitin (dynactin complex 50 kD subunit) (DCTN-50) (non-exact 88%) | 1 | U50733 | | | | | | | |
| dynein, axonemal, heavy polypeptide 17-like (non- exact, 57%aa) | 1 | X99947 | | | | | | | |
| dynein, cytoplasmic, light intermediate polypeptide 2 (DNCLI2) | 1 | AF035812 | В | + | + | | | + | · |
| dynein, cytoplasmic, light intermediate polypeptide 2 (DNCLI2) (non-exact, 69%) dyskeratosis congenita 1, | | AF035812 | | | | | | | |
| dyskerin (DKC1) | 1 | U59151 | В | + | | | + | + | |
| dystonia 1, torsion (autosomal dominant) (DYT1) | 1 | AF007871 | | + | + | + | | + | |
| dystrobrevin, beta (DTNB) | 1 | AF022728 | | + | | | | | |
| dystrophia myotonica- containing WD repeat motif (DMWD) | 1 | L19267 | | + | + | | + | + | · |
| dystrophia myotonica- protein kinase (DMPK) | 1 | L08835 | + | + | + | | | + | |
| dystrophin (muscular dystrophy, Duchenne and Becker types) (DMD) (low match, 59%aa) | 1 | X14298 | | | | | | | |
| E18-55kDa-associated protein | 1 | AJ007509 | W | + | + | | + | + | |
| E2F transcription factor 3 (E2F3) | 2 | D38550 | | + | + | + | + | + | |
| E2F transcription factor 4, p107/p130-binding (E2F4) | 1 | X86096 | В | + | | | + | | |
| E2F transcription factor 5, p130-binding (E2F5) | 2 | U15642 | + | + | | + | | + | |
| E74-like factor 1 (ets domain transcription factor) (ELF1) | 1 | M82882 | В | | + | | + | + | |
| E74-like factor 4 (ets domain transcription factor) (ELF4) | 3 | U32645 | | + | + | | | + | |
| E74-like factor 4 (ets domain transcription factor) (ELF4) (non-exact, 71%) | 1 | U32645 | | | | | | | |
| early development regulator 2 (homolog of polyhomeotic 2) (EDR2) | 4 | U89278 | + | + | + | + | | + | |
| EBV induced G-protein coupled receptor (EBI2) | 1 | L08177 | W | | | | | | |
| ecotropic viral integration site 2B (EVI2B) | 3 | M60830 | | + | | + | | | |
| | | | | | | | | | |

| ectin, galactoside-binding, soluble, 1 (galectin 1) (LGALS1) | 1 | J04456 | | | | | | + | |
|--|----|------------|---|---|---|---|---|---|---|
| EGF-like-domain, multiple 4 (EGFL4) | 7. | AB011541 | | | | | | | |
| elF-2-associated p67 homolog | 3 | U13261 | В | + | | | | + | |
| elastin (supravalvular aortic stenosis, Williams-Beuren syndrome) (ELN) (low match) | 7 | M24782 | | + | + | | | | |
| elav-type RNA-binding protein (ETR-3) | 3 | U69546 | | | | | | | |
| electron-transfer- flavoprotein, alpha polypeptide (glutaric aciduria II) (ETFA) | 2 | J04058 | | + | | | | | |
| (SRF accessory protein 2) (ELK3) | 2 | Z36715 | | | + | | | + | |
| elongation factor 1-beta | 1 | L26404 | | | | | | | |
| elongation factor Ts (mitochondrial protein) | 1 | AF110399 | | | | | | | |
| elongation factor Tu- nuclear encoded mitochondrial | 1 | X84694 | | | _ | | | | |
| eMDC II protein | 1 | AJ242015.1 | | 1 | | | | | |
| ems1 sequence (mammary tumor and squamous cell carcinoma-associated (p80/85 src substrate) (EMS1) | 1 | M98343 | | + | + | | + | + | |
| endogenous retroviral element HC2 | 1 | Z70664 | | | | | | | |
| endosulfine alpha (ENSA) | 1 | X99906 | 1 | + | | | | | |
| endothelial differentiation, sphingolipid G-protein- coupled receptor, 1 (EDG1) | 2 | M31210 | | + | + | + | | + | |
| endothelial differentiation, sphingolipid G-protein- coupled receptor, 1 (EDG1) (low match 66%) | 1 | M31210 | | | | | | | |
| endothelial monocyte- activating polypeptide (EMAPII) | 1 | U10117 | + | + | + | + | | + | |
| enolase 1, (alpha) (ENO1) | 12 | M14328 | + | + | + | + | + | + | |
| enolase 2, (gamma, neuronal) (ENO2) | 1 | X51956 | | + | | | | | , |
| enolase-alpha | 1 | D28437 | | | | | | | |
| enoyl Coenzyme A hydratase 1, peroxisomal (ECH1) | 2 | U16660 | | - | | | | | |
| enoyl Coenzyme A hydratase, short chain, 1, mitochondrial (ECHS1) | 1 | D13900 | + | + | + | + | + | + | |
| ENOYL-COA HYDRATASE, MITOCHONDRIAL PRECURSOR (SHORT CHAIN ENOYL-COA HYDRATASE) (SCEH) (ENOYL-COA HYDRATASE 1) (low match, non-exact 56%) | 1 | P30084 | | | | | | | |
| epidermal growth factor receptor pathway substrate 15 (EPS15) | 2 | U07707 | | + | | + | | + | |

| EPIDIDYMAL SECRETORY PROTEIN E1 PRECURSOR (EPI-1) (HE1) (EPIDIDYMAL SECRETORY PROTEIN 14.6) (ESP14.6) | 2 | Q15668 | | | | | | | : |
|--|------------|----------|------|---|-------|--|--|-----------|------------------------------|
| epithelial membrane protein 3 (EM(P3) | 1 | U87947 | + | + | + | + | | + | |
| Epoxide hydrolase 1, microsomal (xenobiotic) (EPHX1) | 1 | L29766 | | | | | | | + only |
| ERCC2 (=L47234) | 1 | X52221 | 1 | 1 | | | _ | 1 | |
| ERF-2 | 3 | U07802 | + | + | + | + | | + | high in gall bladder |
| ERp28 protein | 1 | X94910 | + | + | + | + | _ | + | |
| erythrocyte membrane protein | 2 | M81635 | | | | | | | |
| erythroleukemic cells K562 | 2 | L25343 | † | | | | | | |
| EST (Hs.189509) | 2 | U24166 | 1 | 1 | | | | - | |
| estrogen receptor-related protein (hERRa1) | 1 | L38487 | | | - '.' | | | | |
| ESTS, Highly similar to ADENYLOSUCCINATE SYNTHETASE | 1 | X66503 | В, Т | + | + | | | | |
| ESTs, Moderately similar to cysteine-rich fibroblast growth factor receptor | | U28811 | + | + | + | + | | + | |
| ET binding factor 1 (SBF1) | 1 | U93181 | + | + | | | | + | |
| ets domain protein ERF | 1 | U15655 | + | + | + | + | | + | |
| eukaryotic translation elongation factor 1 alpha 1 (EEF1A1) | 326 | X03558 | T | + | + | | | + | |
| eukaryotic translation elongation factor 1 alpha 1 (EEF1A1) (low match) | <u>"</u> 1 | X03558 | | | | | | | |
| eukaryotic translation elongation factor 1 alpha 1 (EEF1A1) (low match) | 1 | X03558 | | | | | | | · |
| eukaryotic translation elongation factor 1 beta 2 (EEF1B2) | 5 | X60489 | + | + | + | + | | + | |
| eukaryotic translation elongation factor 1 delta (guanine nucleotide exchange protein) (EEF1D) | 1 | Z21507 | + | + | + | + | + | + | |
| eukaryotic translation elongation factor 1 gamma (EEF1G) | 31 | Z11531 | | | | | | | |
| eukaryotic translation elongation factor 2 (EEF2) | 2 | X51466 | | + | | | | + | |
| eukaryotic translation initiation factor 2, subunit 1 (alpha, 35kD) (EIF2S1) | 1 | J02645 | | | | | | | |
| eukaryotic translation initiation factor 2, subunit 2 (beta, 38kD) (EIF2S2) | 1 | M29536 | | | | | | | |
| eukaryotic translation initiation factor 2, subunit 3 (gamma, 52kD) (EIF2S3) | 3 | L19161 | | + | + | | | | |
| eukaryotic translation initiation factor 3, subunit 10 (theta, 150/170kD) (EIF3S10) | 2 | U78311 | | | | | | | |
| eukaryotic translation initiation factor 3, subunit 2 (beta, 36kD) (EIF3S2) | 3 | U36764 | + | + | + | + | + | + | high in white blood cells |
| eukaryotic translation initiation factor 3, subunit 3 (gamma, 40kD) (EIF3S3) | 6 | U54559 | + | + | + | + | | + | high in spleen |
| eukaryotic translation initiation factor 3, subunit 4 (delta, 44kD) (EIF3S4) | 9 | AF020833 | | + | + | + | | + | |
| | | | | | | | _ | _ | |

WO 00/40749

PCT/CA00/00005

| eukaryotic translation initiation factor 3, subunit 6 (48kD) (EIF3S6) | 4 | U94175 | + | 1 | + | + | | + | high in bladder :- |
|---|----|-------------|------|-----|---|---|----------|---|---|
| eukaryotic translation initiation factor 3, subunit 6 (EIF3S6) | 1 | U62962 | | + | + | + | | + | Highly represented (1.4833 pct) in library 36 human gall bladder |
| eukaryotic translation initiation factor 3, subunit 7 (zeta, 66/67kD) (EIF3S7) | 3 | U54558 | + | + | + | + | | + | |
| eukaryotic translation initiation factor 3, subunit 8, 110KD (EIF3S8) | | U46025 | + | + | + | + | + | + | high in testis |
| eukaryotic translation initiation factor 4 gamma, 1 (EIF4G) | 1 | AF012088 | | | | | | | |
| eukaryotic translation initiation factor 4 gamma, 1 (EIF4G) (low match) | 1 | AF012088 | | | | | | | |
| eukaryotic translation initiation factor 4 gamma, 1 (EIF4G1) | 2 | D12686 | | | | | | | |
| eukaryotic translation initiation factor 4 gamma, 2 (EIF4G2) | 6 | U73824 | + | + | + | + | + | + | |
| eukaryotic translation initiation factor 4 gamma, 2 (EIFG2) | 2 | U76111 | + | + | + | + | + | + | |
| eukaryotic translation initiation factor 4A, isoform 1 (EIF4A1) | 29 | D13748 | | | | | | | |
| eukaryotic translation initiation factor 4A, isoform 2 (EIF4A2) | 11 | D30655 | + | + | + | + | + | + | |
| eukaryotic translation initiation factor 4B (EIF4B) | 18 | X55733 | + | + | + | + | | + | |
| eukaryotic translation initiation factor 4E (EIF4E) | 1 | P06730 | | | | | | | |
| Eukaryotic translation initiation factor 4E binding protein 2 (EIF4EBP2) | 3 | L36056 | Т, В | + | | | + | + | |
| eukaryotic translation initiation factor 4H (EIF4H) | 2 | Q15056 | | | | | | | |
| eukaryotic translation initiation factor 5 (EIF5) | 2 | U49436 | + | + | + | + | + | + | |
| eukaryotic translation termination factor 1 (ETF1) | 2 | U90176 | + | + | + | + | | + | |
| EV12 protein | 1 | M55266 | - | + | | | | | |
| Ewing sarcoma breakpoint region 1 (EWSR1) | 1 | X66899 | + | + | + | + | | + | : |
| EWS/FLI1 activated transcript 2 homolog (EAT- | 2 | AF020264 | | | | | | | |
| EWS-E1A-F chimeric protein | 1 | U35622 | | | | | | | |
| excision repair cross- complementing rodent repair deficiency, | 1 | M28650 | + | + | + | + | | + | |
| complementation group 1 (includes overlapping antisense sequence) (ERCC1) | | 1 | | | | | | | |
| excision repair cross- complementing rodent repair deficiency, complementation group 5 (xeroderma pigmentosum, complementation group G ((Cockayne syndrome)) (ERCC5) | 1 | X69978 | | + | + | + | | + | |
| exostoses (multiple)-like 3 (EXTL3) | 1 | AF001690 | | + | + | + | \neg | + | |
| F11 | 1 | X77744 | | † † | | + | \dashv | | |
| | | | | | | | | | |

| WO 00/40/49 | | | | | | | | • | C1/CA00/00003 |
|---|----|----------|----|---|---|----------|---|---|------------------------|
| F1-ATPase beta subunit (F-1 beta) | 2 | X03559 | | | | | | | r, |
| Fanconi anaemia group A | 2 | Z83095 | | | | | | | |
| Fanconi anemia, | 1 | X99226 | + | + | + | + | | _ | |
| complementation group A (FANCA) | | | | | | | | | |
| far upstream element (FUSE) binding protein 1 (FUBP1) | 2 | U05040 | + | | + | | | + | |
| famesyl diphosphate | 1 | J05262 | + | + | + | + | | + | |
| synthase (famesyl pyrophosphate synthetase, dimethylallyltra nstransferase, geranyltranstransferase) (FDPS) | | | | | | | | | |
| farnesyl-diphosphate famesyltransferase 1 (FDFT1) | 2 | X69141 | + | + | + | + | + | + | |
| farnesyltransferase, CAAX box, beta (FNTB) | 2 | L00635 | | + | + | | | | |
| Fas ligand (gene and promoter region) | 1 | AF044583 | | | | | | | |
| Fas-ligand associated factor 1 | 1 | U70667 | | | | | | | |
| fatty-acid-Coenzyme A ligase, long-chain 1 (FACL1) | 4 | D10040 | + | + | + | + | + | + | |
| Fc fragment of IgA, receptor for (FCAR) | 1 | X54150 | | | | | | | |
| Fc fragment of IgE, high affinity I, receptor for; gamma polypeptide (FCER1G) | 1 | M33195 | + | + | + | + | | + | |
| Fc fragment of IgE, low affinity II, receptor for (CD23A) (FCER2) | 2 | X04772 | + | + | | | | | |
| Fc fragment of IgG, low affinity IIa, receptor for (CD32) | 6 | M31932 | + | + | + | + | + | + | |
| Fc fragment of IgG, low affinity IIa, receptor for (CD32) (FCGR2A) | 1 | X62572 | + | + | + | + | + | + | |
| Fc fragment of IgG, low affinity Illa, receptor for (CD16) (FCGR3A) | 34 | X07934 | + | + | + | + | | + | |
| Fc fragment of IgG, receptor, transporter, alpha (FCGRT) | 3 | U12255 | | + | + | + | + | + | high in many libraries |
| fc-fgr | 1 | Z13983 | | | | <u> </u> | | | |
| Fc-gamma-receptorIIIB (FCGR3B) | 2 | M90746 | | | | | | | |
| feline sarcoma (Snyder- Theilen) viral (v- fes)/Fujinami avian sarcoma (PRCII) viral (v- fps) oncogene homolog(FES) c-fes/fps) | 3 | X06292 | | | | | | | |
| female sterile homeotic- related gene 1 (mouse homolog) (FSRG1) | 2 | X96670 | + | + | + | + | | + | |
| femilin L-chain | 9 | Y09188 | | | | | | | |
| ferritin, heavy polypeptide 1 (FTH1) | 4 | M11146 | + | + | + | + | + | + | |
| fertilin alpha pseudogene | 1 | Y09232 | | | | | | | |
| fetal Alzheimer antigen (FALZ) | 2 | U05237 | | + | | | | | |
| fetal Ig heavy chain variable region | 1 | M34024 | | | | | | | |
| fibrillarin (FBL) | 1 | X56597 | + | + | + | + | + | + | |
| fibrinogen-like protein 2 (T49) | 3 | Z36531 | | | | + | | | |
| | | | 12 | | | | | | |

| tibroblast growth factor receptor 2 (bacteria- expressed kinase, keratinocyte growth factor | 1 | M35718 | + | + | + | + | + | + | |
|--|----------|----------|---|---|----------|---|---|---|---|
| receptor, craniofacial dysostosis 1, Crouzon syndrome) syndrome, | | | | | | | | | |
| Pfeiffer syndrome, Jackson-Weiss) (FGFR2) | | | | | 1 | | | | |
| ficolin (collager/fibrinogen domain-containing) 1 (FCN1) | 19 | D83920 | | | | + | | + | |
| filamin A, alpha (actin- binding protein-280) (FLNA) | 2 | X53416 | | | | | | | |
| filamin B, beta (actin- binding protein-278) (FLNB) | 1 | AF043045 | | + | + | | + | | |
| Finker-Biskis-Reilly murine sarcoma virus (FBR-MuSV) ubiquitously expressed (fox derived); ribosomal protein S30 (FAU) | 2 | X65923 | + | + | + | + | + | + | Highly represented in intraepithelial neoplasia and invasive prostate tumor |
| FK-506 binding protein | 1 | M80199 | + | + | + | + | | + | |
| FK506-binding protein 1A (12kD) (FKBP1A) | 2 | M34539 | | | <u> </u> | | | | |
| FK506-binding protein 1B (12.6 kD) (FKBP1B) | 1 | M92423 | | + | | + | | + | |
| FK506-binding protein 5 (FKBP5) | 4 | U71321 | | + | + | + | | + | |
| Flightless I (Drosophila) homolog (FLII) | 3 | U80184 | | + | | | | | · |
| Flightless I (Drosophila) homolog (FLII) (low match) | 1 | U80184 | | | | | | | |
| FLN29 (FLN29) | 2 | AB007447 | | + | | + | | + | |
| flotillin 2 (FLOT2) | 5 | M60922 | + | + | + | + | + | + | |
| folate receptor 2 (fetal) (FOLR2) forkhead (Drosophila) | 1 | AF000380 | | + | + | + | | + | |
| homolog (rhabdomyosarcoma) like 1 (FKHRL1) | 1 | AF032886 | + | + | | + | | + | |
| Formyl peptide receptor 1 (FPR1) | 9 | M60627 | + | + | + | + | | + | |
| formyl peptide receptor-like 1 (FPRL1) | 1 | M84562 | | | | | | | Found only in libraries from placenta |
| formyl peptide receptor-like 1 (FPRL1) (low score) | 1 | M84562 | | | | | | | |
| fragile X mental retardation 1 (FMR1) | 1 | L29074 | + | + | | + | | + | |
| fragile X mental reterdation, autosomal homolog 1 (FXR1) | 1 | U25165 | + | + | + | + | | | |
| Friend leukemia virus integration 1 (FLI1) | 3 | M93255 | + | + | | | | | |
| fructose-bisphosphatase 1 (FBP1) | 1 | D26054 | | | | + | | + | |
| FSHD-associated repeat DNA, proximal region | 1 | U85056 | | | | | | | |
| fucose-1-phosphate guanylyltransferase (FPGT) | 1 | AF017445 | | + | + | + | | | |
| full length insert cDNA clone ZA78A09 | | AF086122 | | | | | | | |
| full length insert cDNA YP07G10 | 1 | AF075061 | | | | | | | |
| fumarate hydratase (FH) | 1 | U59309 | | + | + | + | | + | |
| FUS (low match) | 1 | X99006 | | | | | | | |
| FYN-binding protein (FYB- 120/130) (FYB) | 16 | U93049 | | + | | + | | | |

| | | | | | | | | - | 31, 311 |
|---|-------------|----------|---------------------------------------|-----------|---|----------|----------|--|---------------------|
| G alpha interacting protein (GAIP) (low score) | 1 | X91809 | | | | | Γ | | :, |
| G protein beta subunit-like | 2 | D28398 | | 1 | | 1 | | | |
| G protein-coupled receptor | 1 | X81892 | | \dagger | - | + | ┢┈ | | |
| 64 (HE6) (non-exact 59%) G protein-coupled receptor | 2 | L16862 | + | + | + | _ | <u> </u> | + | |
| kinase 6 (GPRK6) G1 to S phase transition 1 | | | , , , , , , , , , , , , , , , , , , , | | | ļ | <u> </u> | | |
| (GSPT1) | 2 | X17644 | | + | + | + | + | + | |
| GA-binding protein transcription factor, beta subunit 2 (47kD) (GABPB2) | | D13316 | | + | + | + | + | + | |
| galactose-1-phosphate uridylyltransferase (GALT) | 2 | M60091 | | | | | | | |
| galactosidase, beta 1 (GLB1) | 3 | M27508 | | + | | | + | + | |
| galactosyltransferase (=X13223 N- acetylglucosamide-(beta 1- 4)-galactosyltransferase) | | M13701 | | | | | | | |
| galectin-9 isoform | 1 | AB006782 | + | | | + | | + | |
| gamma2-adaptin (G2AD) | 1 | AF068706 | + | + | | + | | + | |
| gamma-actin | 2 | M37130 | | | | | | | |
| gamma-aminobutyric acid (GABA) B receptor 1 (GABBR1) | 2 | AJ012187 | | + | + | | | + | |
| GATA-binding protein 2 (GATA2) | 1 | M68891 | | | | + | | + | |
| GATA-binding protein 3 (GATA3) | 1 | M69106 | | | + | + | | + | |
| GCN5 (general control of amino-acid synthesis, yeast, homolog)-like 1 (GCN5L1) | 3 | D64007 | + | + | + | + | | + | |
| GDP dissociation inhibitor | 1 | D45021 | + | + | + | + | | + | high in adult brain |
| GDP dissociation inhibitor 2 (GCI2) | 4 | Y13286 | | | | | | | |
| GDS-related protein (HKE1.5) | 4 | U68142 | + | + | + | + | | + | |
| gelsolin (amyloidosis, Finnish type) (GSN) | 3 | X04412 | | + | + | + | + | + | |
| general transcription factor | 4 | Y14946 | + | + | + | + | + | + | |
| general transcription factor III, i, pseudogene 1 (GTF2IP1) | 1 | AF038968 | + | + | + | + | + | + | high in fetal brain |
| general transcription factor IIF, polypeptide 1 (74kD subunit) (GTF2F1) | 4 | X64037 | + | + | + | + | | + | |
| general transcription factor IIH, polypeptide 3 (34kD subunit) (GTF2H3) | 2 | Z30093 | В, Т | | | | | | |
| general transcription factor IIH, polypeptide 4 (52kD subunit) (GTF2H4) | 3 | Y07595 | | + | | + | | + | |
| general transcription factor IIIA (GTF3A) | 1 | U14134 | + | + | | + | | + | |
| general transcription factor IIIC, polypeptide 1 (alpha subunit, 220kD) (GTF3C1) | 1 | U02619 | | + | | + | | | |
| general transcription factor IIIC, polypeptide 2 (beta subunit, 110kD) (GTF3C2) | 3 | D13636 | + | + | + | + | + | + | |
| germline immunoglobulin heavy chain (IGHV@) | 1 | L06612 | | | | | | | |
| germline immunoglobulin heavy chain, variabl region | 1 | X92236 | | | | | | | |
| germline immunoglobulin heavy chain, variable | 1 | X92343 | | | | | | | |
| region, (21-2) | | | | 1 | | | | l | |
| 109.01.; (2 2) | | L— | | | | | | L | · |

| WO 00/40/49 | | | | | | | | P | CT/CAUU/UUUU5 |
|--|-------------|----------|---------------------------------------|-----------|---|--|---|---|---------------|
| GLE1 (yeast homolog)-like, RNA export mediator (GLE1L) | 1 | AF058922 | | + | + | | | | ŗ. |
| glia maturation factor, beta (GMFB) | 1 - | AB001106 | + | + | | + | | + | |
| glioma-associated oncogene homolog (zinc finger protein) (GLI) | 1 | X07384 | | | | | | | |
| glioma-associated oncogene homolog (zinc finger protein) (GLI) (low score) | 1 | X07384 | | | | | | | |
| globin, alpha 2 | 1 | V00516 | | + | | | - | _ | |
| glucocorticoid receptor (=M69104) | 1 | M32284 | · · · · · · · · · · · · · · · · · · · | | | | | | |
| glucocorticoid receptor (GRL) | 2 | U80947 | + | + | + | + | | + | |
| glucos phosphate isomerase (CONTAINS LARGE REPEAT) | 1 | L09105 | | | | | | | |
| glucosamine (N-acetyl)-6- sulfatase (Sanfilippo disease IIID) (GNS) | 1 | Z12173 | + | | | | | | |
| glucosamine (N-acetyl)-6- sulfatase (Sanfilippo disease IIID) (GNS) (non- exact 56%) | 1 | Z12173 | | | | | | | |
| glucose transporter-like protein-III (GLUT3) | 1 | M20681 | | + | + | + | + | + | |
| glucose transporter-like protein-III (GLUT3) (low match) | 1 | M20681 | | | | | | | |
| glucosidase, alpha; acid (Pompe disease, glycogen storage disease type II) (GAA) | 1 | Y00839 | + | + | | + | | + | |
| glucosidase, beta; acid (includes glucosylceramidase) (GBA) | 1 | K02920 | + | + | + | + | | + | |
| glutamate dehydrogenase | 1 | M20867 | | + | + | + | + | + | |
| glutamate-ammonia ligase (glutamine synthase) (GLUL) | 12 | X59834 | + | + | + | + | | + | |
| glutamate-ammonia ligase (glutamine synthase) (GLUL) (low score) | 1 | Y00387 | | | | | | | |
| glutamate-cysteine ligase (gamma-glutamylcysteine synthetase), catalytic (72.8kD) (GLCLC) | 1 | M90656 | | | | + | | | |
| glutamine cyclotransferase | 1 | X71125 | | + | + | | | | |
| glutamine-fructose-6- phosphate transaminase 1 (GFPT1) | 1 | M90516 | | + | | + | | | |
| glutaminyl-tRNA synthetase | 1 | X72396 | | | | | | | |
| giutaminyl-tRNA synthetase (QARS) | 6 | X76013 | + | + | + | + | | + | |
| glutamyl-prolyl-tRNA synthetase (EPRS) | 1 | X54326 | | | | | | | |
| glutathione peroxidase 1 (GPX1) | 2 | M21304 | + | + | + | + | + | + | |
| glutathione peroxidase 4 (phospholipid | 1 | X71973 | + | + | + | + | | + | |
| hydroperoxidase) (GPX4) glutathione S-transferase pi (GSTP1) | | U30897 | | + | + | + | + | + | |
| glutathione S-transferase subunit 13 homolog | 1 | AF070657 | <u> </u> | + | | | | | |
| glyceraldehyde-3- phosphate dehydrogenase (GAPD) | 12 | J02642 | | | | | + | | |
| (One o) | | 1 | | لـــــــا | | لــــا | | | |

| glycogenin (GYG) | 1 | U31525 | | + | + | T + | | T + | |
|--|-----|----------|-------------|---|--|--|--------------|-----|------------------|
| glycophorin C (Gerbich | 1 | X12496 | | + | + | + | \vdash | + | |
| blood group) (GYPC) glycoprotein M68 (GPM68) | -1- | U45955 | | + | + | ┼ | - | ┼ | |
| glycyl-tRNA synthetase | | U09587 | | + | + | ++ | - | + | |
| (GARS) | | | | | | | | | |
| glyoxalase i (lactoyl glutathione lyase) (GLYI) | 1 | L07837 | + | + | + | + | | + | |
| golgi autoantigen, golgin subfamily a, 1 (GOLGA1) | 1 | U51587 | | + | | + | | | |
| golgi autoantigen, golgin subfamily a, 2 (GOLGA2) (non-exact, 70%) | 1 | L06147 | | | | | | | |
| golgi autoantigen, golgin subfamily a, 4 (GOLGA4) | 1 | U31906 | | | | | | | |
| golgi autoantigen, golgin subfamily b, macrogolgin (with transmembrane signal), 1 (GOLGB1) | 1 | X75304 | | + | + | + | | + | |
| gp25L2 protein | 4 | X90872 | | | † | 一 | | | |
| grancalcin | 8 | M81637 | | + | + | + | | | |
| granulin (GRN) | 16 | X62320 | + | + | + | + | <u> </u> | + | |
| granulin (GRN) (low match) | 1 | X62320 | | | - | <u> </u> | | H | |
| Granulysin (NKG5) | 5 | M85276 | + ' | | | | | + | |
| granzyme A (granzyme 1, cytotoxic T-lymphocyte-associated serine esterase 3) (GZMA) | 1 | M18737 | + | + | + | + | | + | |
| GRB2-related adaptor protein (GRAP) | 1 | U52518 | Tonly | 1 | | | | | |
| Grb2-related adaptor protein 2 (GRAP2) | 1 | AF090456 | T | | | | + | | |
| GRO1 oncogene (melanoma growth stimulating activity, alpha) (GRO1) | 1 | X54489 | | | | + | | + | |
| growth arrest and DNA- damage-inducible gene (GADD153) | 1 | \$40706 | | | | | | | |
| growth arrest-specific 7 (GAS7) | 4 | AB007854 | | + | + | | | | |
| growth factor receptor- bound protein 2 (GRB2) | 1 | X62852 | В | + | | | + | + | |
| GS1 (protein of unknown function) | 1 | M86934 | | + | + | + | | | |
| GS3955 | 4 | D87119 | | + | + | + | - | + | |
| GTP binding protein 1 | 1 | U87964 | | + | + | + | | - | |
| (GTPBP1) GTP binding protein similar to S. cerevisiae HBS1 (HBS1) | 1 | U87791 | | + | + | + | | + | |
| GTPase activating protein- like (GAPL) | 1 | AB011110 | | + | + | + | | + | high fetal brain |
| GTP-binding protein (low match) | 1 | Z49068 | | | | | | - | |
| GTP-binding protein G(K), alpha subunit (=G(I) ALPHA-3)(=GTP-binding regulatory protein Gi alpha- 3 chain) | 1 | P08754 | | | | | | | |
| Gu protein (GURDB) | 2 | U41387 | + | | + | + | | + | |
| guanine nucleotide binding protein | 1 | | | | | | | | |
| guanine nucleotide binding protein (G protein), alpha inhibiting activity polypeptide 2 (GNAI2) | 4 | J03004 | + | + | + | + | | + | |

| | | | | | | | | - | |
|---|--------------|----------|-------------|--------------|----------|--------------|--------------|--|--------------------------------|
| guanine nucleotide binding protein (G protein), alpha inhibiting activity | 7 | M20597 | + | + | + | + | | + | " |
| polypeptide 3 (GNAI3) guanine nucleotide binding | 2 | X04409 | | ـ | | ↓ | | ļ., | |
| protein (G protein), alpha stimulating activity polypeptide 1 (GNAS1) | 2 | X04409 | В, Т | + | | | + | + | |
| guanine nucleotide binding protein (G protein), alpha transducing activity | 1 | Z18859 | | | | | | | |
| polypeptide 2 (GNAT2) | | AE047050 | | | <u> </u> | | <u> </u> | L. | ļ |
| guanine nucleotide binding protein (G protein), beta 5 (GNB5) | 2 | AF017656 | | + | + | + | | + | |
| guanine nucleotide binding protein (G protein), beta polypeptide 1 (GNB1) | 5 | M36430 | + | + | + | + | + | + | |
| guanine nucleotide binding protein (G protein), q polypeptide (GNAQ) | 2 | AF011496 | | + | + | +. | | | |
| guanine nucleotide binding protein-like 1 (GNL1) | 1 | L25665 | + | + | + | + | _ | + | |
| guanine nucleotide | 1 | L13857 | + | + | + | + | | | |
| exchange factor guanine nucleotide | | X15610 | + | +- | + | + | <u> </u> | + | |
| regulatory factor (LFP40) | , | 1170700 | | | | <u> </u> | ļ | <u>L</u> | ļ |
| guanine nucleotide regulatory factor (LFP40) | 1 | U72206 | + | + | + | + | | + | |
| GŬANINE NUCLEOTIDE- BINDING PROTEIN BETA | 1 | P25388 | ··· | | | | | | |
| SUBUNIT-LIKE PROTEIN 12.3 (P205) (RECEPTOR OF ACTIVATED PROTEIN KINASE C 1) (RACK1) | | | | | | } | | | ! |
| GUANINE- MONOPHOSPHATE SYNTHETASE (GMPS) | 1 | U10860 | | | + | | | | |
| guanosine monophosphate reductase (GMPR) (non- exact, 72%) | 1 | M24470 | | | | | | | |
| guanosine-diphosphatase like protein | 7 | AF016032 | | † | | | | | |
| guanylate binding protein 1, interferon-inducible, 67kD (GBP1) | 2 | M55542 | | + | + | + | + | + | |
| guanylate binding protein 2, interferon-inducible (GBP2) | 6 | M55543 | + | + | + | + | | + | |
| H2A histone family, member C (H2AFC) | 1 | Z83742 | | | | | | | |
| H2A histone family, member Y (H2AY) | 2 | AF041483 | + | + | + | + | | + | |
| H2B histone family, member L (H2BFL) | 2 | Z80783 | + | + | + | + | + | + | high in adrenal gland tumor |
| h2-calponin | 1 | D86059 | | 1 | | | | | |
| H-2K binding factor-2 | 1 | L08904 | | + | + | + | | + | |
| H3 histone family, member K (H3FK) | 1 | Z83735 | | | | | | | |
| H3 histone, family 3A (H3F3A) | 7 | M11353 | + | + | + | + | | + | high in ovary |
| H3 histone, family 3B (H3.3B) (H3F3B) | 15 | Z48950 | + | + | + | + | | + | high in endothelial cells |
| hbc647 | 1 | U68494 | | + | + | + | + | | |
| heat shock 27kD protein 1 (HSPB1) | 1 | U12404 | | + | + | | +. | + | |
| heat shock 40kD protein 1 (HSPF1) | 4 | D85429 | + | + | + | + | + | + | high in testis |
| heat shock 60kD protein 1 (chaperonin) (HSPD1) | 3 | M22382 | + | + | + | + | + | + | |
| heat shock 70kD protein 1 (HSPA1A) | 7 | M59828 | + | + | + | + | + | + | high in activated T cells |
| | | | | | | | | | |

| heat shock 70kD protein 5 (glucose-regulated protein, 78kD) (HSPA5) | 13 | X87949 | | + | + | | + | | r. |
|---|----------|--------------------|-------------|------------|----------|----------|---|---|-------------------------|
| heat shock 70kD protein 6 (HSP70B') (HSPA6) | 4 | X51757 | + | + | + | | | | |
| heat shock 70kD protein 9B (mortalin-2) (HSPA9B) | 2 | L15189 | | + | + | + | + | + | |
| HEAT SHOCK COGNATE 71 KD PROTEIN | 1 | P11142 | | | | | | | |
| heat shock factor binding protein 1 (HSBP1) | 2 | AF068754 | | | | | | | |
| heat shock protein 90 | 13 | M27024 | + | + | + | + | + | + | high in many libraries |
| heat shock protein, DNAJ- like 2 (HSJ2) | 1 | D13388 | | + | + | | + | + | |
| Hect (homologous to the E6-AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1 (HERC1) | 1 | U50078 | | + | + | + | | | |
| hect domain and RLD 2 (HERC2) | 1 | AB002391 | + | + | + | + | | + | |
| helicase-like protein (HLP) | 1 | X98378 | + | + | | + | | + | |
| helix-loop-helix protein HE47 (E2A) | 1 | M65214 | | | ····· | | | + | |
| hematopoietic cell-specific Lyn substrate 1 (HCLS1) | 18 | X16663 | + | | + | + | | + | |
| heme oxygenase (decycling) 1 (HMOX1) | 1 | X06985 | | + | | + | + | + | |
| HEMOGĽÓBÍN ALPHÁ CHAIN | 1 | P19015 | | | | | | | |
| hemoglobin beta (beta globin) hemoglobin, alpha 1 | 5 301 | AF117710 V00491 | | | + | | + | + | |
| (HBA1) hemoglobin, alpha 1 | 301 | V00491 V00491 | | | | | | _ | |
| (HBA1) (low match) hemoglobin, alpha 1 (low | 1 | V00491 V00493 | | | | <u> </u> | | | |
| match) hemoglobin, alpha 1 (non- | 1 | J00153 | | | | | | | |
| exact, 76%) hemoglobin, alpha 1 (non- | 1 | V00493 | | ļ <u> </u> | | | _ | | |
| exact, 82%) hemoglobin, beta (HBB) | 129 | V00497 | + | | + | + | + | + | high in many libraries |
| , , , | | | т | 1 - | | | | _ | ingir iir many iibranes |
| hemoglobin, beta (HBB) (low match) | | V00497 | | | | | | | |
| hemoglobin, beta (HBB) (low match) hemokine (C-X-C motif), | 1 | L48220 D10924 | + | + | + | + | | + | |
| receptor 4 (fusin) (CXCR4) | 5 | M16591 | | Ť | T | + | | + | |
| (HCK) | | | | | | | | | |
| hepatitis C-associated microtubular aggregate protein p44 | 2 | D28908 | | | | | | | |
| hepatoma-derived growth factor | 1 | D16431 | + | + | + | + | | + | |
| Hermansky-Pudlak syndrome (HPS) | 2 | U65676 | | | | | | | |
| HERV-E integrase (non- exact 76%aa) | 1 | AF026246 | | | | | | | |
| heterogeneous nuclear protein similar to rat helix destabilizing protein (FBRNP) | 2 | S63912 | | + | + | + | | + | |
| heterogeneous nuclear ribonucleoprotein (C1/C2) (HNRPC) | 4 | M16342 | | | | | | | |
| heterogeneous nuclear ribonucleoprotein A/B (HNRPAB) | 1 | M65028 | + | + | + | + | + | + | |

| *************************************** | | | | | | | | • ` | 31,0,100,0000 |
|---|----|------------|---------------------------------------|---|----------|--------------|----------|-----|---|
| heterogeneous nuclear ribonucleoprotein A1 (HNRPA1) | 20 | X12671 | + | + | + | + | + | + | High in alveolar :- rhabdomyosarcoma |
| heterogeneous nuclear ribonucleoprotein A2/B1 (HNRPA2B1) | 3 | M29064 | + | + | + | + | + | + | High in activated T cell, fetal brain |
| heterogeneous nuclear ribonucleoprotein D (hnRNP D) | 2 | D55673 | + | + | + | + | + | + | |
| heterogeneous nuclear ribonucleoprotein D-like (HNRPDL) | 5 | D89092 | + | + | + | + | + | + | |
| heterogeneous nuclear ribonucleoprotein F (HNRPF) | 1 | L28010 | + | + | + | + | | + | |
| heterogeneous nuclear ribonucleoprotein F (HNRPF) (83%) | 1 | L28010 | | | | | | | |
| heterogeneous nuclear ribonucleoprotein G (HNRPG) | 2 | Z23064 | | + | + | + | | + | |
| heterogeneous nuclear ribonucleoprotein H (HNRPH) (FTP-3) | 3 | P55795 | | | | | | | |
| heterogeneous nuclear ribonucleoprotein H (HNRPH) (low match) | 1 | P31943 | | | | | | | |
| heterogeneous nuclear ribonucleoprotein H1 (H) (HNRPH1) | 2 | L22009 | + | + | + | + | | + | |
| heterogeneous nuclear ribonucleoprotein K (HNRPK) | 21 | S74678 | + | + | + | + | + | + | |
| heterogeneous nuclear ribonucleoprotein R (HNRPR) | 1 | AF000364 | | + | + | + | + | + | |
| heterogeneous nuclear ribonucleoprotein U (scaffold attachment factor A) (HNRPU) | 3 | X65488 | + | + | + | + | + | + | |
| hexokinase 1 (HK1) | 2 | X66957 | | + | + | + | \vdash | + | |
| hexokinase 2 (HK2) | 3 | Z46376 | + | + | + | + | ├ | + | |
| hexokinase 3 (HK3) | 2 | U51333 | | | | ┼─ | | ├ | |
| hexosaminidase A (alpha | 1 | S62047 | · | | <u> </u> | | ļ | | |
| polypeptide) (HEXA | • | 302047 | | | | | | | |
| HGMP07I gene for olfactory receptor | 2 | U76377 | | | | | | | |
| High density lipoprotein | 2 | M64098 | + | + | + | + | + | + | |
| binding protein (HDLBP) high-mobility group | | V40-04 | | | | | | | |
| (nonhistone chromosomal) | 5 | X12597 | + | + | + | + | + | + | |
| high-mobility group (nonhistone chromosomal) protein 1 (HMG1) (non- exact 60%) | 1 | D63874 | | | | | | | |
| High-mobility group (nonhistone chromosomal) protein 17 (HMG17) | 2 | M12623 | + | + | + | + | | + | |
| high-mobility group (nonhistone chromosomal) protein 2 (HMG2) | 2 | M83665 | + | + | + | + | + | + | |
| high-mobility group (nonhistone chromosomal) protein isoforms I and Y | 2 | L17131 | + | + | + | | + | + | |
| high-risk humanpapilloma viruses E6 oncoproteins targeted protein E6TP1 beta (=A8007900 KIAA0440) | 1 | AF090990.1 | , , , , , , , , , , , , , , , , , , , | | | | | | |
| histidine ammonia-lyase | 1 | D16626 | | | | , only | | | |

WO 00/40749

PCT/CA00/00005

| histidyl-tRNA synthetase | 1 | Z11518 | , , , , , , , , , , , , , , , , , , , | | | | | | |
|---|--|------------|---|----------------|--------------|--|--|----------------|---------------------------------------|
| (HARS) | 2 | | + | | + | + | + | + | " |
| histocompatibility antigen (HLA-Cw3), class I | 1 | U31372 | | | | | Ţ | | |
| histone deacetylase 1 (HDAC) | 4 | U50079 | + | + | + | + | | + | |
| histone deacetylase 1 (HDAC1) | 2 | D50405 | + | + | + | + | | + | |
| histone deacetylase 5 (NY- CO-9) | 1 | AF039691 | | + | + | | | | |
| HK2 gene for hexokinase II | 1 | Z46362 | † | | | 1 | | | |
| HL9 monocyte inhibitory receptor precursor | 2 | U91928 | | † · · · · | _ | + | | | |
| HLA class I heavy chain (HLA-Cw*1701) | 1 | | | 1 | | | | | |
| HLA class I locus C heavy chain | 1 | X58536 | | | | | | | |
| HLA class II SB 4-beta | 1 | X03022 | | | | | <u> </u> | | |
| HLA class III region containing NOTCH4 gene | 1 | U89335 | + | + | + | + | <u> </u> | + | |
| HLA-A | 7 | Z72423 | <u> </u> | + | _ | +- | | - | · · · · · · · · · · · · · · · · · · · |
| HLA-A | 2 | AJ006020 | | + | | \vdash | | - | |
| HLA-A*7402 | 1 | AJ223060 | | +- | - | | | - | - |
| HLA-A11 | | U02934 | - | + | - | ├ | \vdash | \vdash | |
| HLA-B | 2 | X75953 | | - | | +- | - | \vdash | |
| ньа-в | - | X83401 | | | <u> </u> | - | ├ | | |
| HLA-B | 1 | X78426 | | | ļ | ⊢ | l | | |
| HLA-B associated | | Z37166 | | +- | + | + | + | + | <u> </u> |
| transcript-1 (D6S81E) | | | <u> </u> | * | * | | * | | |
| HLA-B associated transcript-2 (D6S51E) | 2 | M33509 | + | + | + | + | | | |
| HLA-B*1529 | 4 | D44501 | | | | | | | |
| HLA-Bw72 antigen | 119 | L09736 | + | + | + | + | + | + | high in many libraries |
| HLA-C gene (HLA- Cw*0701 allele) | 1 | D83957 | | | | | | | |
| HLA-Cw*0701 | 9 | Z46810 | | | | | | | |
| HLA-Cw*0801 | 1 | D64151 | | | | | | | |
| HLA-Cw*1203 | 1 | D64146 | | | | | | | • |
| HLA-DC classII histocompatibility antigens alpha-chain (=K01160) | 2 | X00370 | | | | | | | |
| HLA-DR alpha-chain | 17 | M60333 | + | + | + | + | + | + | high in spleen |
| HLA-F (leukocyte antigen F) | 3 | X17093 | | | + | + | | + | |
| HMG box containing protein 1 | 3 | AF019214 | | | | | | | |
| hMLH1 (=U83845) | 1 | AB017806.1 | | | | | | | |
| Hmob33 | 3 | Y14155 | | | | | | | |
| HMT1 (nnRNP methyltransferase, S. cerevisiae)-like 1 (HRMT1L1) | 2 | U80213 | + | + | + | + | | + | |
| hnRNP C1/C2 | 2 | D28382 | | | | | | | |
| homeobox (=X58250 Mouse homeo box protein, put. transcription factor involved in embryogenesis and hematopoiesis) | 1 | M60721 | | | | | | | |
| homeobox protein (HLX1) (=M60721) | | U14326 | | | | | | | |
| homeodomain-interacting protein kinase 3 (HIPK3) | 1 | AF004849 | + | | + | + | | + | |
| homolog of Drosophila past (PAST) | | AF001434 | + | + | + | + | | + | |
| homolog of yeast (S. cerevisiae) ufd2 (UFD2) | 3 | D50916 | | + | + | + | | + | |

50

| AB015344 | | | | | | | | | • • | 31,01100,0000 | • |
|--|---|---|------------|-------------|---|---|--|--|--|---------------|---|
| HRX.His protein (eAPT) 1 | HPV16 E1 protein binding protein | 1 | U96131 | | + | + | | | + | | |
| (=AF010403 ALR) shock cognate protein shock cognate protein HSPC012 1 | | | | | + | + | | | + | | |
| shock cognate protein HSPC012 | (=AF010403 ALR) | | Y08836 | | | | | | | | |
| HSPC021 | shock cognate protein | 3 | Y00371 | | | | | | | | |
| HSPext3p | | 1 | AF077036.1 | | | | \vdash | | 1- | | |
| Name | HSPC021 | 1 | AF077207.1 | | | | | 1 | | | |
| HU-K4 1 080644 | HsPex13p | 1 | U71374 | | + | | | | <u> </u> | | |
| RIUNKI 1 Y12059 + + + + + + + + + + + + + + + + + + + | htra2-beta-2 | 1 | U87836 | + | + | + | + | | + | | |
| HUNKI 1 Y12059 + + + + + + H HUNTING H | HU-K4 | 1 | U60644 | | | | | | | | |
| AF-049528 Truntingtin-interacting protein HYPA/FBP11 Truntingtin-interacting protein HYPA/FBP11 Truntingtin-interacting protein HYPA/FBP11 Truntingtingtingtingtingtingtingtingtingtin | hunc18b2 | 1 | U63533 | | + | + | + | | + | | |
| protein HYPA/FBP11 (HYPS41) | HUNKI | 1 | Y12059 | + | + | | + | + | + | | |
| hydroxyacyl-Coenzyme A dehydrogenasea/3-ketoacyl-Coenzyme A hitolase/enoyl-Coenzyme A hydratase (tifunctional protein), alpha subunit (HADHA) hydroxyacyl-Coenzyme A hydratase (tifunctional protein), alpha subunit (HADHA) hydroxyacyl-Coenzyme A hydratase (tifunctional protein), beta subunit (HADHB) hydroxysteroid (17-beta) dehydrogenase 1 (HSD1781) hypothetical protein 1 (HSD1781) hypothetical protein 1 (HSD1781) hypothetical protein 1 (HSD1781) hypothetical protein 1 (HSD1781) hypothetical protein 1 (HSD1781) hypothetical protein 1 (HSD1781) hypothetical protein 1 (HSD1781) hypothetical protein 1 (HSD1781) hypothetical protein (ICIn9878K, ZA8.1 chromosome 8) hypothetical protein (ICIn9878K, ZA8.1 chromosome 8) hypothetical protein (ICIn9878K, ZA8.1 chromosome 8) hypothetical protein (ICIn9878K, ZA8.1 chromosome 8) hypothetical protein (ICIn9878K, ZA8.1 chromosome 8) hypothetical protein (ICIn9878K, ZA8.1 chromosome 8) hypothetical protein 1 AL022238 (ICIn9878K, ZA8.1 chromosome 8) hypothetical protein 2 AL031432 (ICIn9878K, ZA8.1 chromosome 8) hypothetical protein 2 AL031432 (ICIn9878K, ZA8.1 chromosome 8) hypothetical protein 2 AL031432 (ICIN9878K, ZA8.1 chromosome 8) hypothetical protein 2 AL031432 (ICIN9878K, ZA8.1 chromosome 8) hypothetical protein 2 AL031432 (ICIN9878K, ZA8.1 chromosome 8) hypothetical protein 3 AL031432 (ICIN9878K, ZA8.1 chromosome 8) hypothetical protein 3 AL031432 (ICIN9878K, ZA8.1 chromosome 8) hypothetical protein 3 AL031432 (ICIN9888K, ZA8.1 chromosome 8) hypothetical protein 3 AL031432 (ICIN988K, ZA8.1 chromosome 8) hypothetical protein 3 AL031432 (ICIN988K, ZA8.1 chromosome 8) hypothetical protein 3 AL031432 (ICIN988K, ZA8.1 chromosome 8) hypothetical protein 3 AL031432 (ICIN988K, ZA8.1 chromosome 8) hypothetical protein 4 AL031432 (ICIN988K, ZA8.1 chromosome 8) hypothetical protein 4 AL031432 (ICIN988K, ZA8.1 chromosome 8) hypothetical protein 4 AL031432 (ICIN988K, ZA8.1 chromosome 8) hypothetical protein 4 AL031432 (ICIN988K, ZA8.1 chromosome 8) hypothetical protein 4 AL031444 (I | huntingtin-interacting protein HYPA/FBP11 (HYPA) | | | | | | | | | | |
| dehydrogenase/3-ketoacyl-Coenzyme A htholase/enoyl-Coenzyme A htholase/enoyl-Coenzyme A htholase/enoyl-Coenzyme A hydratase (infunctional protein), alpha subunit (HADHA) hydroxyacyl-Coenzyme A dehydrogenase/3-ketoacyl-Coenzyme A htholase/enoyl-Coenzyme A htholase/enoyl-Coenzyme A hydratase (infunctional protein), beta subunit (HADHB) hydroxysteroid (17-beta) dehydrogenase 1 (HSD1781) hydroxysteroid (17-beta) dehydrogenase 1 (HSD1781) hypothetical protein 1 (HSD1781) hypothetical protein 1 (HSD1781) hypothetical protein 1 (CIT8875K, ZA8 1 chromosome 8) hypothetical protein (clone 1 Z70222 (CRFp50762480). hypothetical protein (clone 1 Z70222 (CRFp50762480). hypothetical protein (clone 1 AL022238 (dJ104257A7.1) hypothetical protein (dJ1042K10.4) (non-exact 76%) hypothetical protein (dJ1042K10.4) (non-exact 76%) hypothetical protein (dJ1042K17.1.1) 2 AL031432 (dJ1042K17.1.1) 2 AL031432 (dJ1042K17.1.1) 2 AL03853 (dJ104753P8.2) hypothetical protein 1 AL050131.1 (DKF2p5861111) hypothetical protein 1 AL050729 hypot | | i | | | | | | | | | |
| dehydrogenase/3-ketoacyl- Coenzyme A thiolase/enoyl-Coenzyme A hydratase (trifunctional protein), beta subunit ((HADHB) hydroxysteroid (17-beta) dehydrogenase 1 (HSD17B1) hypothetical protein (HSD17B1) hypothetical protein (AL008729) (JC57A7.2) hypothetical protein (CIT987SK 2AB 1 chromosome 8) hypothetical protein (clone (CRFp507G2490). hypothetical protein (Al1042K10.4) (non-exact 78%) hypothetical protein (Jd465N24.1 similar to predicted yeast and worm proteins) hypothetical protein (Jd47SAP9.2) hypothetical protein (Jd47SAP9.2) hypothetical protein (JC57A7.2) hypothetical protein (JC57A7.2) hypothetical protein (JC57A7.2) hypothetical protein (JC57A7.2) hypothetical protein (JC57A7.2) hypothetical protein (JC57A7.2) hypothetical protein (JC57A7.2) hypothetical protein (JC57A7.2) hypothetical protein (JC57A7.2) hypothetical protein (JC67A7.2) hypothetical protein (JC67A7.2) hypothetical protein (JC67A7.2) hypothetical protein (JC67A7.2) hypothetical protein (JC67A7.2) hypothetical protein (JC67A7.2) hypothetical protein (JC67A7.2) hypothetical protein (JC67A7.2) hypothetical protein (JC67A7.2) hypothetical protein | dehydrogenase/3-ketoacyl- Coenzyme A thiolase/enoyl-Coenzyme A hydratase (trifunctional protein), alpha subunit (HADHA) | 1 | U04627 | | + | + | | + | | | |
| dehydrogenase 1 (HSD17B1) hypothetical protein 1 (AL008729) (JJ257A7.2) hypothetical protein 1 (CIT9875K_2A8_1 chromosome 8) hypothetical protein (clone 1 24840) hypothetical protein (clone 1 270222 ICRFp507G2490). hypothetical protein 1 (J1042K10.4) (non-exact 76%) hypothetical protein 2 (J465N24.1 similar to proteins) hypothetical protein 2 (J485N24.1 similar to proteins) hypothetical protein 2 (J485N27.1.1) hypothetical protein 2 (J485N37.1.1) hypothetical protein 1 (J48777.1.1) hypothetical protein 2 (J5753P9.2) hypothetical protein 1 hypothetical protein (LTH 3' region) | hydroxyacyl-Coenzyme A dehydrogenase/3-ketoacyl- Coenzyme A thiolase/enoyl-Coenzyme A hydratase (trifunctional protein), beta subunit (HADHB) | 1 | D16481 | + | + | + | + | | + | | |
| hypothetical protein (AL008729) (J257A7.2) hypothetical protein (CiT9875K_2A8_1 chromosome 8) hypothetical protein (Cione 24640) hypothetical protein (Cione 1 CRFp50762490). hypothetical protein (Cione 1 AL022238 (CJ1042K10.4) (non-exact 76%) hypothetical protein (CJ0486N24.1 similar to predicted yeast and worm proteins) hypothetical protein (CJ48717.1.1) hypothetical protein (CJ48777.1.1) hypothetical protein (CJ783P9.2) hypothetical protein (DKFZp588i111) hypothetical protein (DKFZp588i111) hypothetical protein (DKFZp588i111) hypothetical protein (DKFZp588i111) hypothetical protein (DKFZp588i111) hypothetical protein (DKFZp586i1111) hypothetical protein (DKFZp586i1111) hypothetical protein (DKFZp586i1111) hypothetical protein (DKFZp586i1111) hypothetical protein (DKFZp586i1111) hypothetical protein (DKFZp586i111111111111111111111111111111111111 | dehydrogenase 1 (HSD17B1) | | U34879 | | + | | | + | | | , |
| (AL008729) (dJ257A7.2) hypothetical protein (Cione defation of the protein (Cirip875K, 2A8, 1 chromosome 8) hypothetical protein (clone defation of the protein (clone defation of the protein (clone defation of the protein (clone defation of the protein (clone defation of the protein (clone defation of the protein defation of the protein (dJ1042K10.4) (non-exact f8%) hypothetical protein defation defation of the protein defation of the | I | · | | | | | | | | | |
| hypothetical protein (CIT987SK_2A8_1 chromosome 8) hypothetical protein (clone 24840) hypothetical protein (clone 1 270222 (CRFp507G2490). hypothetical protein (dJ1042K10.4) (non-exact 76%) hypothetical protein (dJ465N24.1 similar to proteins) hypothetical protein 2 AL031432 (dJ465N24.1 similar to proteins) hypothetical protein 2 AL008730 (dJ487J7.1.1) hypothetical protein 3 AL023653 (dJ753P9.2) hypothetical protein 4 AL050131.1 (DKFZp586i111) hypothetical protein 5 AL008729 (J257A7.2) hypothetical protein 6 AB007900 KKIAA0440) (=AF026504 R.norvegicus SPA-1 like protein (L1H 3' region) | (AL008729) (dJ257A7.2) | 1 | | | | | | | | | |
| hypothetical protein (clone 24640) hypothetical protein (clone 1 270222 | hypothetical protein (CIT987SK_2A8_1 | 1 | U96629 | | | | | | | | |
| CRFp507G2490). hypothetical protein 1 | hypothetical protein (clone 24640) | 1 | AF055004 | | | | | | | | |
| hypothetical protein (dJ1042K10.4) (non-exact 76%) hypothetical protein 2 AL031432 (dJ465N24.1 similar to predicted yeast and worm proteins) hypothetical protein 2 AL008730 (dJ487J7.1.1) hypothetical protein 2 AL023653 (dJ753P9.2) hypothetical protein 1 AL050131.1 (DKFZp586i111) hypothetical protein 1 AL008729 (J257A7.2) hypothetical protein 1 AB007900 (KIAA0440) (=AF026504 R. norvegicus SPA-1 like protein) hypothetical protein (L1H 3' 1 region) | hypothetical protein (clone ICRFp507G2490). | 1 | Z70222 | - | | | | | | | |
| (dJ465N24.1 similar to predicted yeast and worm proteins) hypothetical protein (dJ487/7.1.1) hypothetical protein (dJ753P9.2) hypothetical protein (DKFZp586i111) hypothetical protein (J2 AL023653 (dJ753P9.2) hypothetical protein (DKFZp586i111) hypothetical protein (J257A7.2) hypothetical protein (J257A7.2) hypothetical protein (KIAA0440) (=AF026504 R. norvegicus SPA-1 like protein) hypothetical protein (L1H 3' 1 region) | hypothetical protein (dJ1042K10.4) (non-exact 76%) | 1 | AL022238 | | | | | | - | | |
| (dJ487J7.1.1) hypothetical protein (dJ753P9.2) hypothetical protein (DKFZp586i111) hypothetical protein (J257A7.2) hypothetical protein (J257A7.2) hypothetical protein (KIAA0440) (=AF026504 R.norvegicus SPA-1 like protein) hypothetical protein (L1H 3' 1 region) | (dJ465N24.1 similar to predicted yeast and worm proteins) | _ | AL031432 | | | | | | | | |
| (dJ753P9.2) hypothetical protein | (dJ487J7.1.1) | _ | AL008730 | | | | | | | | |
| (DKFZp586l111) hypothetical protein (J257A7.2) hypothetical protein (KIAA0440) (=AF026504 R.norvegicus SPA-1 like protein) hypothetical protein (L1H 3' 1 region) | (dJ753P9.2) | | AL023653 | | | | | | | | |
| (J257A7.2) hypothetical protein 1 AB007900 (KIAA0440) (=AF026504 R.norvegicus SPA-1 like protein) hypothetical protein (L1H 3' 1 region) | hypothetical protein (DKFZp586I111) | | AL050131.1 | - | | | | | | | |
| (KÍAA0440) (=AF026504 R.norvegicus SPA-1 like protein) hypothetical protein (L1H 3' 1 region) | (J257A7.2) | | AL008729 | | | | | | | | |
| region) | hypothetical protein (KIAA0440) (=AF026504 R.norvegicus SPA-1 like protein) | 1 | AB007900 | | | | | | | | |
| hypothetical protein (S184) 1 P49756 | hypothetical protein (L1H 3' region) | | | | | | | | | | |
| | hypothetical protein (S164) | 1 | P49756 | | | | | | | | |

PCT/CA00/00005

WO 00/40749

| hypothetical protein (similar to thrombospondin) (non- exact 56%) | 1 | AF109907 | | | | | | | 10 |
|---|----------|------------------|----------|--------------|----------|--|--------------|--------------|------------------------|
| hypothetical protein 3 | 1: | | | | | | | 1 | |
| hypothetical protein B (HSU47926) (non-exact, 56%) | 1 | U47926 | | | | | | | |
| hypothetical protein from BCRA2 region (CG005) | 3 | U50532 | + | + | + | + | | + | |
| hypoxia-inducible factor 1, alpha subunit (basic helix- loop-helix transcription factor) (HIF1A) | 1 | AF050115 | | | | | | | |
| la-associated invariant gamma-chain (clones lambda-y (1,2,3)) | 1 | M13555 | | | | | | | |
| iduronate 2-sulfatase (Hunter syndrome) (IDS) | 2 | M58342 | + | + | + | + | _ | + | |
| Ig heavy chain V region (=D11016) | 1 | L20779 | | | | | | | |
| Ig heavy chain variable region | 2 | M34024 | | | <u> </u> | | | | |
| lig heavy chain variable region (VH4DJ) (clone IT14.4) | 1 | 275378 | | | | | | | |
| Ig heavy chain variable region (VH4DJ) (clone T22.18) | 1 | 275392 | | | | | | | |
| lg J chain | 7 | M12378 | | | - | \vdash | | | |
| lg kappa | 1 | S49007 | | | | 1 | | | |
| IG kappa light chain variable region A20 | 1 | X63398 | | | | | | | |
| Ig kappa light chain, V- and J-region (=X59315) | 1 | D90158 | | | | | | | |
| lg lambda light chain variable region (26- 34ITIIF120) | 1 | Z85052 | | | | | | | |
| Ig mu-chain VDJ4-region | 1 | M16949 | <u> </u> | | | | | | |
| Ig rearranged anti-myelin kappa-chain (V-J4-region, hybridoma AE6-5) | 1 | M29469 | | | | | | | |
| Ig rearranged H-chain mRNA V-region | 2 | M97920 | | | | | | | |
| Ig rearranged light-chain V region (=D90158) | 1 | M74020 | | | | | | | |
| IGF-II mRNA-binding protein 3 (KOC1) (non- exact, 75%) | 1 | U97188 | + | + | + | | | | |
| IgG Fc binding protein (FC(GAMMA)BP) | 1 | D84239 | + | + | | + | | + | |
| lgG heavy chain variable region (vH26) | 1 | M83136 | | | | | | | |
| IgM heavy chain (C mu, membrane exons) | 1 | X14939 | | | | | | | |
| IkB kinase-beta (IKK-beta) | 1 | AF029684 | | | | | | | |
| IL-1 receptor type II IL2-inducible T-cell kinase | 1 2 | U14177 S65186 | | | | | | | |
| (ITK) | 1 | M62831 | + | ļ | + | + | | + | |
| (ETR101) | <u>'</u> | D87018 | | - | | <u> </u> | | <u> </u> | |
| (lambda) | · | Y08915 | В, Т | + | + | | + | ļ | |
| binding protein 1 (IGBP1) immunoglobulin C (mu) and | | X57331 | J., 1 | ' | | | <u> </u> | - | |
| C (delta) heavy chain (=K02878) | | | | | | | | | |
| immunoglobulin G Fc receptor IIIB | 1 | Z46223 | | | | | | | |
| immunoglobulin gamma 3 (Gm marker) (IGHG3) | 3 | Y14737 | + | | | + | | + | high in many libraries |
| | | | | | | | | | |

| WO 00/40749 | | | | | | | | PC | T/CA00/00005 |
|--|---|----------|---|----|---|---|----------|----|--------------|
| immunoglobulin gamma heavy chain variable region | 1 | Z66542 | | | | | | | 7. |
| (=X61011) immunoglobulin heavy | 1 | X62109 | | 11 | | | \dashv | | |
| chain (VI-3B) immunoglobulin heavy | 1 | X86356 | | | | | | | |
| chain J region immunoglobulin heavy | 2 | X86355 | | | | | | | |
| chain J region, B1 haplotype | | Arnenane | | 1 | | | | | |
| immunoglobulin heavy chain variable region (IGH) (clone 21u-48) | 1 | AF062126 | | | | | | | |
| immunoglobulin heavy chain variable region (IGH) (clone 23u-1) | 1 | AF062212 | | | į | | | | |
| immunoglobulin heavy chain variable region V1-18 (IGHV@) (=X60503) | 2 | M99641 | | | | | | | |
| immunoglobulin heavy chain variable region V3-43 (IGHV@) | 2 | M99672 | | | | | | | |
| immunoglobulin heavy chain variable region V3-7 (IGHV@) | 3 | M99649 | | | | | | | |
| immunoglobulin IgH heavy chain Fd fragment | 1 | U07986 | | | | | | | |
| immunoglobulin kappa light chain | 1 | X58081 | | | | | | | |
| immunoglobulin kappa light chain V-segment A27 | 1 | X12686 | | | | | | | |
| immunoglobulin light chain | 1 | D86990 | | | | | | | |
| immunoglobulin light chain (low match) | 1 | D86996 | | | | | | | |
| immunoglobulin light chain variable region (lambda IIIb subgroup) from IgM rheumatoid factor | 1 | L29157 | | | | | | | |
| immunoglobulin M heavy chain V region=anti-lipid A antibody | 1 | \$50735 | | | | | | | |
| immunoglobulin mu (IGHM) | 9 | X57086 | + | + | | + | | + | |
| immunoglobulin mu binding protein 2 (IGHMBP2) | 1 | L24544 | | + | | | + | | |
| immunoglobulin superfamily, member 2 (IGSF2) | 1 | 233642 | | | | | | | |
| immunoglobulin VH mRNA (487 bp) (=M99652 immunoglobulin heavy chain variable region V3-11 (IGHV@)) | 1 | X61013 | | | | | | | |
| imogen 38 (IMOGEN38) | 1 | Z68747 | | + | + | + | | + | |
| IMP (inosine monophosphate) dehydrogenase 1 (IMPDH1) | 1 | J05272 | + | + | + | + | | | |
| IMP (inosine monophosphate) dehydrogenase 2 (IMPDH2) | 2 | L39210 | + | + | + | + | | + | |
| inc finger protein 151 (pHZ-67) (ZNF151) | | Y09723 | + | + | + | + | | + | |
| inc finger protein, C2H2, rapidly turned over (ZNF20) | 1 | AF011573 | | + | + | | | | |
| inducible poly(A)-binding protein (IPABP) | 1 | U33818 | + | + | + | + | | + | |
| inducible poly(Á)-binding protein (IPABP) (low match) | 1 | U33818 | | | | | | | |

| | | | | | | | | • | C1/CA00/00003 |
|---|----|----------|---|---|---|---|---|---|---------------|
| inducible protein (Hs.80313) | 2 | L47738 | + | + | + | + | | + | :- |
| inhibitor of DNA binding 2, dominant negative helix- loop-helix protein (ID2) | 4 | M97796 | + | + | + | + | + | + | |
| inhibitor of kappa light polypeptide gene enhancer in B-cells, kinase complex- associated protein ((IKBKAP) | 2 | AF044195 | | | | | | | |
| inositol 1,3,4-trisphosphate | 1 | U51336 | + | + | + | + | + | + | |
| inositol 1,4,5 trisphosphate receptor type 1 (ITPR1) | 1 | U23850 | | + | + | + | | | |
| inositol 1,4,5-trisphosphate 3-kinase B (ITPKB) | 2 | X57206 | В | + | + | | + | | |
| inositol monophosphatase | 1 | 538980 | | | | | - | | |
| inositol polyphosphate-5- phosphatase, 145kD (INPP5D) | 2 | U84400 | + | + | + | + | | + | |
| Ins(1,3,4,5)P4-binding protein | 1 | X89399 | | + | | | | + | |
| insulin-like growth factor 2 receptor (IGF2R) | 5 | Y00285 | + | + | + | + | | + | |
| integral membrane protein 1 (ITM1) | 1 | L38961 | | | + | + | | + | |
| integral membrane protein 2C (ITM2C) | 1 | AF038953 | Ţ | | + | | + | + | |
| integral membrane protein Tmp21-I (p23) | 3 | U61734 | + | + | + | + | + | + | |
| integrin beta 4 binding protein (ITGB4BP) | 2 | AF047433 | | | + | | | + | |
| integrin, alpha 2b (platelet glycoprotein IIb of IIb/IIIa complex, antigen CD41B) (ITGA2B) | 3 | M34480 | | + | | | + | | |
| integrin, alpha 5 (fibronectin receptor, alpha polypeptide) (ITGA5) | 4 | X06256 | + | + | + | | + | + | |
| integrin, alpha L (antigen CD11A (p180), lymphocyte function-associated antigen 1; alpha polypeptide) ((iTGAL) | 6 | Y00796 | | | | | | | |
| integrin, alpha M (complement componentreceptor 3, alpha; also known as CD11b (p170), macrophage antigen alpha polypeptide) (ITGAM) | | M18044 | | | | | | | |
| integrin, alpha X (antigen CD11C (p150), alpha polypeptide) (ITGAX) | | M81695 | + | + | | | | + | |
| integrin, beta 1 (fibronectin receptor, beta polypeptide, antigen CD29 includes MDF2 MSK12) (ITGB1) | 2 | X07979 | | | | | | | |
| integrin, beta 2 (antigen CD18 (p95), lymphocyte function-associated antigen 1; macrophage antigen 1 (mac-1) beta subunit) (ITGB2) | 32 | M15395 | + | + | | + | | + | |
| integrin, beta 7 (ITGB7) | 1 | M68892 | + | | | | | | |
| Integrin-linked kinase (ILK) | 1 | U40282 | + | + | + | + | | + | |
| intercellular adhesion molecule 1 (CD54), human rhinovirus receptor (ICAM1) | 1 | J03132 | + | | | + | + | + | |
| intercellular adhesion molecule 2 (ICAM2) | 1 | X15606 | + | + | + | + | | + | |

| intercellular adhesion molecule 3 (ICAM3) | 6 | X69819 | + | | | | | + | :- |
|--|----|----------|-----------|--------|------|----------|----------|----------|---|
| intercellular adhesion molecule 4, Landsteiner- Wiener blood group (ICAM4) | | L27670 | | | | | | + | |
| Interferon consensus sequence binding protein 1 (ICSBP1) | 1 | M91196 | W, 1 | lymp | homa | | <u> </u> | | |
| Interferon consensus sequence binding protein 1 (ICSBP1) (low match) | 1 | M91196 | | | | | | | |
| interferon regulatory factor 2 (IRF2) | 4 | X15949 | + | + | + | + | | | |
| interferon regulatory factor1 (IRF1) | 4 | L05072 | + | + | + | + | | + | |
| interferon regulatory factor5 (IRF5) | 1 | U51127 | + | + | | + | | | |
| interferon, gamma- inducible protein 16 (IFI16) | 2 | M63838 | + | + | + | + | | + | |
| interferon, gamma- inducible protein 30 (IFI30) | 9 | J03909 | + | + | | + | | + | |
| INTERFERON-INDUCED GUANYLATE-BINDING PROTEIN 1 (GUANINE NUCLEOTIDE-BINDING PROTEIN 1) (non-exact 62%) | 1 | P32455 | | | | | | | |
| interferon-induced protein 17 (IFI17) | 3 | X84958 | | + | + | + | | + | |
| interferon-induced protein 54 (IFI54) | 5 | M14660 | | | | | | | |
| interferon-inducible (1-8D) | 5 | X57351 | T | | + | | + | + | |
| interferon-inducible (1-8U) | 1 | X57352 | | L. | + | <u> </u> | + | + | |
| developmental regulator 1 (IFRD1) | 5 | Y10313 | | + | + | | | + | |
| interferon-stimulated transcription factor 3, gamma (48kD) (ISGF3G) | 2 | M87503 | | + | | + | | + | · |
| interleukin 1 receptor, type II (IL1R2) | 1 | U64094 | | | | + | | | |
| Interleukin 10 receptor, beta (I.10RB) | 1 | U08988 | Tactivate | ed | + | | | + | |
| interleukin 12 receptor, beta 1 (IL12RB1) | 2 | U03187 | + | | | | | <u> </u> | only found in T cell |
| interleukin 13 receptor, alpha 1 (IL13RA1) | 2 | Y09328 | | + | + | + | + | + | |
| interleukin 16 (lymphocyte chemoattractant factor) (IL16) | 6 | U82972 | | + | | | | | |
| interleukin 18 receptor 1 (IL18R1) | 1 | U43672 | | | | | | | |
| interleukin 2 receptor, beta (IL2RB) | 9 | M26062 | | | | | | | |
| interleukin 2 receptor, gamma (severe combined immunodeficiency) (IL2RG) | 6 | D11086 | + | | + | | | + | |
| interleukin 4 receptor (IL4R) | 3 | X52425 | + | + | | + | | + | |
| interleukin 6 receptor (IL6R) | 5 | X12830 | | + | | | · | + | |
| interleukin 6 signal transducer (gp130, oncostatin M receptor) (IL6ST) | .1 | M57230 | | | | | | | |
| interleukin 7 receptor (IL7R) | 14 | M29696 | + · | | | | | + | |
| interleukin 7 receptor (IL7R) (low match) | 1 | AF043123 | | | | | | | |
| interleukin 8 (IL8) | 8 | Y00787 | + | | + | | + | | High in activated T cells, bone and pancreatic islets |

55

| interleukin 8 receptor alpha (IL8RA) | 11 | L19591 | | | | | | | |
|--|-----|----------|-----------------|---|---|----------|----------|----------|--|
| interleukin 8 receptor, beta | 14 | M94582 | | | | | | | |
| interleukin enhancer | 3 | U10323 | + | + | + | + | + | + | high in uterus |
| binding factor 2, 45kD (ILF2) | - | | | | | | | | |
| interleukin enhancer binding factor 3, 90kD (ILF3) | 2 | U10324 | | | | | | | |
| interleukin-1 receptor- | 2 | L76191 | | + | + | + | | + | |
| associated kinase 1 (IRAK1) | 2 | 270151 | | | • | | | | |
| interleukin-1 receptor- | 1 | U52112 | | 1 | | | | <u> </u> | |
| associated kinase 1 (low match) | | | | | | | | | |
| interleukin-10 receptor, alpha (IL10RA) | 5 | U00672 | + | + | + | + | | | |
| interleukin-11 receptor, | 7 | Z38102 | | + | + | | | | |
| alpha (IL11RA) | | | | | | | <u> </u> | | |
| INTERLEUKIN-14 PRECURSOR (IL-14) (HIGH MOLECULAR WEIGHT B-CELL GROWTH FACTOR) (HMW-BCGF) (non-exact 46%) | 1 | P40222 | | | | | | | |
| intestinal carboxylesterase; | 1 | U60553 | · ······· · · · | + | | \vdash | + | | |
| liver carboxylesterase-2 (ICE) | | | | | | | | | |
| inversin protein (non-exact 52%) | 1 | AF084367 | | | | | | | |
| IQ motif containing GTPase activating protein 1 (IQGAP1) | 6 | L33075 | | | | | | | |
| IQ motif containing GTPase activating protein 2 (IQGAP2) | · 1 | U51903 | | + | | + | | | |
| isocitrate dehydrogenase 1 (NADP+), soluble (IDH1) | 1 | AF020038 | + | + | + | + | + | + | |
| isocitrate dehydrogenase 2 (NADP+), mitochondrial (IDH2) | 2 | X69433 | + | + | + | + | + | + | |
| isocitrate dehydrogenase 3 (NAD+) alpha (IDH3A) | 2 | U07681 | | | + | | | | |
| isocitrate dehydrogenase 3 (NAD+) gamma (IDH3G) | 1 | Z68907 | + | + | + | + | | + | |
| isolate Aus3 cytochrome b (CYTB) | 1 | AF042516 | | | | | | | |
| isolate TzCCR5-179 CCR5 receptor (CCR5) | 1 | AF011524 | | | | | | | |
| isopentenyl-diphosphate delta isomerase (IDI1) | 5 | X17025 | + | + | + | + | | + | |
| Janus kinase 1 (a protein tyrosine kinase) (JAK1) | 4 | M64174 | + | + | + | + | | + | |
| Janus kinase 2 (a protein tyrosine kinase) (JAK2) | 1 | AF005216 | | | | | | | |
| Jk-recombination signal binding protein (RBPJK) | 2 | L07876 | | | | | | | |
| JM1 protein | 1 | AJ005890 | | + | | + | | | |
| jumonji (mouse) homolog (JMJ) | 1 | U57592 | | + | + | + | | + | |
| jun D proto-oncogene (JUND) | 1 | X51346 | + | + | + | + | | + | |
| jun dimerization protein | 1 | AF111167 | | | | | | | only found in germ |
| junction plakoglobin (JUP) | 1 | M23410 | | + | + | + | | + | |
| | | | | | | | | | |

| WG 00/40/49 | | | | | | | | | |
|--|-------------|----------|-------------|---|---|---|---|----------|---|
| kangai 1 (suppression of tumorigenicity 6, prostate; CD82 antigen (R2 | 1 | U20770 | + | + | + | + | + | + | ; · · · · · · · · · · · · · · · · · · · |
| leukocyte antigen, antigen detected by monoclonal and antibody IA4)) (KAI1) | · | | | | | | | | |
| karyopherin (importin) beta 1 (KPNB1) | 2 | L39793 | + | + | + | + | + | + | |
| karyopherin (importin) beta 2 (KPNB2) | 1 | U72395 | + | + | + | + | | | |
| karyopherin alpha 1 (importin alpha 5) (KPNA1) | 1 | S75295 | + | + | + | | + | | |
| karyophenn alpha 2 (RAG cohort 1, importin alpha 1) (DPNA2) | | U09559 | | | | | | | |
| karyopherin alpha 3 (importin alpha 4) (KPNA3) | 1 | D89618 | | + | | | + | | |
| karyopherin alpha 4 (KPNA4) | 1 | M17887 | | + | + | | | | |
| Katanin (80 kDa) (KAT) | 1 | AF052432 | | + | + | + | | + | |
| KE03 protein | 2 | AF064604 | | | | | | | |
| Kelch-like ECH-associated protein 1 (KIAA0132) (66%aa) | 1 | D50922 | | | | | | | |
| Keratin 8 (KRT8) | 1 | X74929 | | + | + | + | + | + | |
| ketohexokinase (fructokinase) (KHK) | 1 | X78678 | | + | | + | + | | |
| KIAA0001 (KIAA0001) (72% aa) | 1 | Q15391 | | | | | | | |
| KIAA0001 (KIAA0001) (76% aa) | 1 | Q15391 | | | | | | | |
| KIAA0001 (KIAA0001) (non-exact 72%) | 1 | Q15391 | | _ | + | + | | + | |
| KIAA0002 (KIAA0002) | 5 | D13627 | | · | | | | | |
| KIAA0005 (KIAA0005) | 4 | D13630 | | + | + | + | | + | |
| KIAA0010 (KIAA0010) | 1 | D13635 | | + | | | | + | |
| KIAA0016 (KIAA0016) | 1 | D13641 | + | + | + | + | | + | |
| KIAA0017 (KIAA0017) | 2 | D87686 | | | | | | | |
| KIAA0022 (KIAA0022) | 2 | D14664 | | + | + | + | | <u> </u> | |
| KIAA0023 (KIAA0023) | 1 | D14689 | | + | | | | | |
| KIAA0024 (KIAA0024) | 1 | D14694 | + | + | + | + | | + | |
| KIAA0025 (KIAA0025) | 1 | D14695 | | + | + | + | + | + | |
| KIAA0026 (KIAA0026) | 2 | D14812 | | + | + | + | | + | |
| KIAA0027 | 1 | D25217 | | + | | | | | |
| KIAA0032 (KIAA0032) | 2 | D25215 | | + | + | + | | | |
| KIAA0040 (KIAA0040) | 1 | D25539 | + | + | + | + | | + | |
| KIAA0050 (KIAA0050) | 4 | D26069 | | | | | | | |
| KIAA0053 (KIAA0053) | 17 | D29642 | + | | + | + | | | |
| KIAA0057 (KIAA0057) | 1 | D31762 | + | + | + | + | + | + | high in fetal lung |
| KIAA0058 (KIAA0058) | 11 | D31767 | + | | + | + | | + | |
| KIAA0063 (KIAA0063) | 3 | D31884 | + | + | + | + | | + | |
| KIAA0064 (KIAA0064) | 1 | D31764 | + | + | + | + | | + | |
| KIAA0066 | 1 | D31886 | + | + | + | + | | + | |
| KIAA0068 | 1 | D38549 | | + | + | + | + | + | |
| KIAA0073 | 3 | D38552 | | + | + | + | | + | |
| KIAA0081 | 2 | D42039 | | + | | + | | + | |
| KIAA0084 | 2 | D42043 | + | + | + | + | | + | |
| KIAA0085 | 26 | U30498 | + | + | + | + | + | + | |
| KIAA0088 | 3 | D42041 | + | + | + | + | + | + | |
| KIAA0090 | 2 | D42044 | + | + | + | + | + | + | |
| KIAA0092 (KIAA0092) | 1 | D42054 | | + | + | + | | + | |
| | | | | | | | | | |

| KIAA0094 | 3 | D42084 | | | + | + | | | 5- |
|--------------------------|--------------|----------|--|--|--|--|--|--|--------------|
| KIAA0095 (KIAA0095) | 1 | D42085 | | | | | | | |
| KIAA0096 | 1: | D43636 | + | + | + | + | | + | |
| KIAA0097 (KIAA0097) | | X92474 | | + | + | - | + | | |
| KIAA0099 (KIAA0099) | 3 | D43951 | + | + | + | + | + | + | |
| KIAA0102 (KIAA0102) | - 2 | D14658 | | + | | + | + | + | |
| KIAA0105 | 1 | D14661 | В | + | | | + | + | - |
| KIAA0120 | | P37802 | | | | - | | | |
| KIAA0120 (non-exact, | | M83106 | | | | - | | _ | |
| (non-exact, | • | 14163100 | | | | | | | |
| KIAA0121 (KIAA0121) | 1 | D50911 | + | + | + | + | | + | |
| KIAA0123 | 1 | D21064 | | + | + | + | | + | |
| KIAA0128 | 1 | D50918 | + | + | + | + | | + | |
| KIAA0129 (KIAA0129) | 1 | D50919 | + | + | + | + | | | |
| KIAA0130 (KIAA0130) | 1 | AF055995 | | + | + | + | | | |
| KIAA0136 | 2 | D50926 | | † | | | | | |
| KIAA0137 (KIAA0137) | 1 | AB004885 | | + | + | + | | + | |
| KIAA0140 (KIAA0140) | 1 | D50930 | + | + | | + | | + | |
| KIAA0141 (KIAA0141) | 3 | D50931 | | † | \vdash | Г | | \vdash | |
| KIAA0144 (KIAA0144) | 3 | D63478 | + | + | + | + | 1 | + | |
| KIAA0144 (KIAA0144) (low | | D63478 | | | <u> </u> | | | | |
| match) | | | | | | <u> </u> | | | |
| KIAA0144 (non-exact 61%) | 1 | Q14157 | | <u> </u> | | ļ | <u> </u> | | |
| KIAA0144 (non-exact 65%) | 1 | Q14157 | | | | <u> </u> | | <u> </u> | |
| KIAA0146 | 2 | D63480 | | + | + | + | | + | |
| KIAA0148 (KIAA0148) | 1 | D63482 | | + | | | | + | |
| KIAA0154 | 2 | D63876 | + | <u> </u> | + | + | | + | |
| KIAA0156 | 1 | D63879 | | + | + | + | | + | |
| KIAA0160 | 2 | D63881 | | | | Ĺ. | | | <u> </u> |
| KIAA0161 (KIAA0161) | 1 | D79983 | + | + | | + | | | |
| KIAA0164 (KIAA0164) | 3 | D79986 | | | | | | | |
| KIAA0167 (KIAA0167) | 1 | D79989 | | + | | 1 | | | |
| KIAA0168 (KIAA0168) | 3 | D79990 | | + | + | + | | + | |
| KIAA0169 | 3 | D79991 | | | | | | | |
| KIAA0171 (KIAA0171) | 3 | D79993 | | + | + | + | | + | |
| KIAA0174 (KIAA0174) | 7 | D79996 | + | + | + | + | | + | |
| KIAA0179 | 2 | D80001 | | + | + | + | | + | |
| KIAA0181 | 1 | D80003 | | + | + | + | | + | |
| KIAA0183 | 4 | D80005 | + | + | + | + | + | + | |
| KIAA0184 | 1 | D80006 | + | + | + | + | | + | |
| KIAA0191 (72% aa) | 1 | D83776 | | | | \top | T | | 1 |
| KIAA0191 (non-exact 77%) | 1 | + | | | | | | | |
| KIAA0193 (KIAA0193) | 1 | D83777 | + | + | + | + | | + | |
| KIAA0200 (KIAA0200) | 1 | D83785 | | + | + | + | T | + | |
| KIAA0210 (KIAA0210) | 3 | D86965 | | +- | + | | <u> </u> | 1- | |
| KIAA0217 | 2 | D86971 | + | + | + | + | 1 | + | |
| KIAA0219 | 2 | U77700 | | + | + | + | \vdash | + | |
| KIAA0222 (KIAA0222) | 1 | D86975 | | + | ┼── | \vdash | | + | |
| KIAA0223 | 2 | D86976 | | + | +- | + | + | +- | |
| KIAA0229 | 1 | D86982 | + | ++ | 1 | +- | ╁ | + | |
| KIAA0232 (KIAA0232) | <u> </u> | D86985 | | ++ | ++ | + | +- | + | |
| KIAA0233 (KIAA0233) | | D87071 | <u></u> | + | + | + | ┼ | +- | |
| KIAA0235 | 2 | D87078 | + | + | + | + | \vdash | - | |
| KIAA0239 | 1 | D87076 | + + | + | | ╁ | \vdash | + | |
| וווראטנטט | <u> </u> | 201010 | <u> </u> | <u> </u> | <u> </u> | | 1 | ــــــــــــــــــــــــــــــــــــــ | <u> </u> |

| KIAA0239 (non-exact 80%) KIAA0240 KIAA0242 KIAA0248 KIAA0249 (KIAA0249) KIAA0253 KIAA0254 (KIAA0254) KIAA0255 (KIAA0255) KIAA0262 (KIAA0262) KIAA0263 (KIAA0263) KIAA0264 | 1 1 4 2 3 5 1 4 3 1 | D87076 D87077 D87684 D87435 D87436 D87442 D87443 D87444 D87451 | + + + | + + + + + | + + + | + + + | + | ++ | |
|---|--|--|--|--------------|-------|-------|---|----|-------------|
| KIAA0242 KIAA0248 KIAA0249 (KIAA0249) KIAA0253 KIAA0254 (KIAA0254) KIAA0255(KIAA0255) KIAA0262 (KIAA0262) KIAA0263 (KIAA0263) | 4 2 3 5 1 4 3 | D87684 D87435 D87436 D87442 D87443 D87444 | + | + | + | + | + | | |
| KIAA0248 KIAA0249 (KIAA0249) KIAA0253 KIAA0254 (KIAA0254) KIAA0255(KIAA0255) KIAA0262 (KIAA0262) KIAA0263 (KIAA0263) | 2 3 5 1 4 3 | D87435 D87436 D87442 D87443 D87444 | + | + | + | + | + | | |
| KIAA0249 (KIAA0249) KIAA0253 KIAA0254 (KIAA0254) KIAA0255(KIAA0255) KIAA0262 (KIAA0262) KIAA0263 (KIAA0263) | 3 5 1 4 3 | D87436 D87442 D87443 D87444 | | + | | | | + | |
| KIAA0253 KIAA0254 (KIAA0254) KIAA0255(KIAA0255) KIAA0262 (KIAA0262) KIAA0263 (KIAA0263) | 5 1 4 3 1 | D87442 D87443 D87444 | | L | + | + | | | |
| KIAA0254 (KIAA0254) KIAA0255(KIAA0255) KIAA0262 (KIAA0262) KIAA0263 (KIAA0263) | 1 4 3 | D87443 D87444 | + | + | | | İ | + | |
| KIAA0255(KIAA0255) KIAA0262 (KIAA0262) KIAA0263 (KIAA0263) | 4 3 1 | D87444 | | | + | + | + | + | |
| KIAA0262 (KIAA0262) KIAA0263 (KIAA0263) | 3 | | | + | + | + | | | |
| KIAA0263 (KIAA0263) | 1 | D87451 | | + | + | + | | + | |
| | • | | + | + | + | + | | + | |
| KIAA0264 | 3 | D87452 | + | + | + | + | | + | |
| | - | D87453 | | + | + | + | | + | |
| KIAA0268 | 1 | D87742 | + | + | | + | | + | |
| KIAA0269 | 1 | Q92558 | | - | | | | | |
| KIAA0275 (KIAA0275) | 13 | D87465 | + | + | | + | | + | |
| KIAA0304 (KIAA0304) | 2 | AB002302 | + | + | + | + | + | + | |
| KIAA0308 | 2 | AB002306 | | + | + | | | + | |
| KIAA0310 (KIAA0310) | 1 | AB002308 | ~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~ | + | + | + | | + | |
| KIAA0314 (=U96635 | 3 | AB002312 | | | | | | | |
| M.musculus ubiquitin | | | | | | | | | |
| protein ligase Nedd-4) KIAA0315 (KIAA0315) | 4 | AB002313 | | + | + | + | + | + | |
| KIAA0325 (=L08505 | - 2 | AB002313 | | <u> </u> | | , | | | |
| R.norvegicus cytoplasmic dynein heavy chain (MAP 1C)) | 2 | AB002323 | | | | | | | |
| KIÁA0329 (KIAA0329) | 1 | AB002327 | | + | + | + | | + | |
| KIAA0330 | 1 | AB002328 | + | + | + | | | + | |
| KIAA0332 | 1 | AB002330 | | + | + | + | | + | |
| KIAA0333 | 2 | AB002331 | | + | + | + | + | + | |
| KIAA0336 (KIAA0336) | 3 | AB002334 | + | + | + | + | | + | |
| KIAA0336 (KIAA0336) (low match) | 1 | AB002334 | | | | | | | |
| KIAA0342 (KIAA0342) | 1 | AB002340 | | + | + | | | + | |
| KIAA0344 (KIAA0344) | 2 | AB002342 | | | | + | | + | |
| KIAA0354 (KIAA0354) | 1 | AB002352 | + | + | + | + | | + | |
| KIAA0365 (KIAA0365) | 3 | AB002363 | + | + | + | + | + | + | |
| KIAA0370 | 6 | AB002368 | | + | + | + | + | + | |
| KIAA0372 (KIAA0372) | 1 | AB002370 | | | | | | | |
| KIAA0373 (KIAA0373) | | AB002371 | | + | | + | | | |
| KIAA0375 (KIAA0375) | 1 | AB002373 | | + | | + | | | |
| KIAA0377 (KIAA0377) | 1 | AB002375 | | + | | + | + | | |
| KIAA0379 | 1 | AB002377 | | | | + | | | |
| KIAA0379 (non-exact, 65%) | 1 | AB002377 | *************************************** | | | | | | |
| KIAA0380 (KIAA0380) | 1 | AB002378 | + | + | | + | | + | |
| KIAA0380 (KIAA0380) (60%aa) | 1 | AB002378 | <u></u> _ | | | | | | |
| KIAA0382 (KIAA0382) | 2 | AB002380 | | + | + | + | | + | |
| KIAA0383 | 1 | AB002381 | | | | | | | |
| KIAA0386 (KIAA0386) | 5 | AB002384 | | | | | | | |
| KIAA0392 | 1 | AB002390 | · | | | | | | |
| KIAA0397 (KIAA0397) | 4 | AB007857 | | + | + | + | + | + | |
| KIAA0403 | 3 | AB007863 | | | | | | | |
| KIAA0404 | 1 | AB007864 | | + | | + | | | |
| KIAA0409 | 1 | AB007869 | | + | | + | | | |
| KIAA0421 | 1 | AB007881 | + | + | + | | | + | |
| KIAA0424 (non-exact 82%) | 7 | AB007884 | | 1 | | | | | |

| KIAA0428 (KIAA0428) | 9 | Y13829 | | T | 1 | I | | Т | |
|---------------------|--|----------------------|---------------------------------------|--|--------------|--|---------------|--|--|
| KIAA0429 (KIAA0429) | 2 | AB007889 | + | + | + | + | † | + | |
| KIAA0430 (KIAA0430) | 2 | AB007890 | • | + | | | \vdash | ╁ | only in ovary |
| KIAA0432 (KIAA0432) | 2 | U86753 | | + | + | ╁╌ | + | + | |
| KIAA0435 (KIAA0435) | | AB007895 | | | \vdash | + | ╁┈ | + | |
| KIAA0438 (KIAA0438) | 1 1 | AB007898 | | + | + | ╁ | + | + | |
| KIAA0447 (KIAA0447) | 3 | AB007916 | + | + | + | + | ╁ | + | |
| KIAA0449 | 1 | AB007918 | | + | - | - | ┼ | + | <u> </u> |
| KIAA0456 | | AB007925 | | + | + | + | +- | + | |
| KIAA0458 (KIAA0458) | | AB007927 | | + | ┼ | \vdash | ┼ | ┼ | <u> </u> |
| KIAA0462 | 1 1 | AB007931 | + | + | + | + | ├ | + | |
| KIAA0465 | 1 | AB007934 | | ++ | + | + | + | + | |
| KIAA0476 (KIAA0476) | 1 | AB007945 | | ++- | + | + | | | |
| KIAA0489 | | AB007958 | | + - | <u> </u> | <u> </u> | ├ | ├ ─ | |
| KIAA0494 (KIAA0494) | 1 - | AB007963 | + | + | + | + | | + | |
| KIAA0515 | | AB011087 | + | + | + | + | ├— | + | |
| KIAA0521 | 3 | AB011093 | | + | <u> </u> | <u> </u> | ├— | + | |
| KIAA0525 | 1 1 | AB011097 | | + | ├ | + | - | <u> </u> | |
| KIAA0530 | 1 1 | AB011097 | | + | + | + | | - | ļ |
| KIAA0532 | | AB011102 | + | ++ | + | + | ļ | + | |
| KIAA0537 (KIAA0537) | | AB011109 | тт | | <u> </u> | ĻŤ | ـــــ | <u> </u> | |
| KIAA0540 | | AB011109 AB011112 | + | ++ | + | <u> </u> | <u> </u> | ļ., | |
| KIAA0543 | | AB011112 | | ↓ <u> </u> | 1 | + | - | + | |
| KIAA0544 | | 11 | <u> </u> | | + | + | <u> </u> | + | |
| KIAA0549 | ! | AB011116 | | + | + | + | <u>L</u> | + | |
| KIAA0551 | 2 | AB011121 | | + | + | + | <u> </u> | + | |
| KIAA0554 | 2 | AB011123 | | + | | | <u> </u> | + | |
| KIAA0561 | 8 | AB011126 | | + | + | + | | + | |
| | 1 | AB011133 | | + | | + | <u> </u> | <u> </u> | · |
| KIAA0562 (KIAA0562) | | AB011134 | | <u> </u> | | | | L | |
| KIAA0563 (KIAA0563) | 1 | AB011135 | | | | | <u> </u> | | |
| KIAA0569 (KIAA0569) | 2 | AB011141 | | + | + | + | | + | |
| KIAA0571 (KIAA0571) | 2 | AB011143 | | + | + | + | l | | |
| KIAA0573 | 1 | AB011145 | | + | | + | | + | |
| KIAA0576 | 1 | AB011148 | | | | | | | |
| KIAA0580 | 1 | AB011152 | | | | | | | |
| KIAA0584 | 1 | AB011156 | | + | | | | | |
| KIAA0592 | 3 | AB011164 | + | + | + | + | | + | |
| KIAA0596 | 1 | AB011168 | | + | + | | | | |
| KIAA0598 (KIAA0598) | 1 | AB011170 | | + | + | + | | | |
| KIAA0608 | 1 | AB011180 | | | + | + | | | |
| KIAA0614 | 2 | AB014514 | + | + | + | + | | + | |
| KIAA0615 (KIAA0615) | 1 | AB014515 | | | | | | | |
| KIAA0621 | 1 | AB014521 | | + | + | | | + | |
| KIAA0648 | 1 | AB014548 | | + | + | + | | + | |
| KIAA0652 (KIAA0652) | 1 | AB014552 | + | + | + | + | | + | |
| KIAA0668 | 1 | AB014568 | · · · · · · · · · · · · · · · · · · · | | | | | | |
| KIAA0669 | 1 | AB014569 | | | | | | | |
| KIAA0671 (KIAA0671) | 1 | AB014571 | | | + | + | | + | |
| KIAA0675 (KIAA0675) | 1 | AB014575 | | + | | + | + | | |
| KIAA0676 | 1 | AB014576 | | + | + | + | | + | |
| KIAA0677 (KIAA0677) | 2 | AB014577 | | + | + | + | + | + | |
| KIAA0678 | 1 | AB014578 | + | + | + | + | | + | |
| KIAA0679 | 6 | AB014579 | | + | + | + | | + | |
| | 1 | L | | | | | | لــــا | |

| [] T T T T T T T T T T T T T T T T T T T | | | | | | | | | C1/CA00/00003 |
|--|------|------------|--------------|--|---|--|--|----------------|---------------------------------------|
| KIAA0680 (KIAA0680) | 1 | AB014580 | | | | | | | :, |
| KIAA0692 | 1 | AB014592 | + | + | + | + | | + | |
| KIAA0697 | 1. | AB014597 | | | | | | | |
| KIAA0699 | 1 | AB014599 | + | + | + | + | | + | |
| KIAA0700 | 1 | AB014600 | | + | + | + | | + | |
| KIAA0737 (KIAA0737) | 3 | AF014837 | + | + | + | + | | + | |
| KIAA0748 (KIAA0748) | 2 | AB018291 | | + | | | | | |
| KIAA0763 (KIAA0763) | 2 | AB018306 | + | + | + | + | | + | |
| KIAA0769 (KIAA0769) | 2 | AB018312 | | + | + | + | | + | |
| KIAA0782 | 1 | AB018325 | + | + | | + | 1 | † | high in BPH stroma |
| KIAA0796 | 1 | AB018339 | | + | + | + | | + | |
| KIAA0798 (KIAA0798) | 1 | AB018341 | | 1 | | ! | \vdash | | |
| KIAA0823 | 1 | AB020630 | · | | | | \vdash | \vdash | |
| KIAA0854 | 1 | AB020661 | + | + | + | + | — | + | |
| KIAA0856 | 1 | AB020663 | <u> </u> | + | + | + | | + | |
| KIAA0860 | 1 | AB020667 | | + | | + | | ╁ | |
| KIAA0862 | 1 | AF054828 | | + | + | + | - | - | |
| KIAA0871 (non-exact 88%) | 1 | AB020678 | | + | | | \vdash | | |
| KIAA0873 | 1 | AB020680 | | + | + | + | ├─ | + | |
| KIAA0892 | 1 | AB020699 | + | + | + | + | _ | + | |
| KIAA0906 | 1 | AB020713 | + | + | + | + | _ | + | · |
| KIAA0991 | 1 | AB023208.1 | <u> </u> | - | | - | _ | | · · · · · · · · · · · · · · · · · · · |
| killer cell lectin-like | 1 | U11276 | | | + | + | - | + | |
| receptor subfamily B, member 1 (KLRB1) | · | | | | | • | | | |
| killer cell lectin-like receptor subfamily C, member 4 (KLRC4) | 1 | U96846 | | | | | | | |
| kinectin 1 (kinesin receptor) (KTN1) | 1 | D13629 | | | | | | | |
| kinesin family member 5B (KIF5B) | 2 | X65873 | | + | + | + | | | |
| kinesin-like DNA binding protein | 1 | AB017430 | + | + | + | + | | + | |
| Krueppel-related DNA- binding protein (TF6) (low match) | 1 | M61869 | | | | | | | |
| Kruppel related gene (clone pHKR1RS) | 1 | M20675 | | | | | | | |
| Kruppel-like zinc finger | 3 | U51869 | + | + | + | + | + | + | |
| protein Zf9 | | | | | | | | | i |
| Kruppel-like zinc finger protein Zf9 (non-exact 76%) | 1 | U44975 | | + | + | | + | + | |
| kruppel-type zinc finger protein, ZK1 | 1 | AB011414.1 | | | | | | | |
| L apoferritin | 3 | X03742 | | | | | | | |
| lactate dehydrogenase A (LDHA) | 3 | X02152 | | + | + | + | + | + | |
| lactate dehydrogenase A (LDHA) (non-exact, 81%) | 1 | X02152 | | | | | | | |
| lactate dehydrogenase B (LDHB) | 6 | X13794 | + | + | + | + | + | + | high in fetal lung fibrablast |
| lactotransfernin (LTF) | 1 | U07643 | + | |] | + | | + | high in bone marrow |
| laminin binding protein (low score) | 1 20 | D28372 | - | | | | | | |
| laminin receptor 1 (67kD); Ribosomal protein SA (LAMR1) | 20 | X15005 | + | + | + | + | + | + | high in many libraries |
| laminin receptor homolog (3' region) | 1 | S35960 | | | | | | | |
| laminin, gamma 1 (formerly LAMB2) (LAMC1) | 2 | J03202 | + | + | + | | | + | |

| WO 00/40749 | | | | | | | | | CI/CA00/00003 |
|---|---|----------|-------------|----------|---|----------|---|---|-----------------------|
| latent transforming growth factor beta binding protein 1 (LTBP1) | 2 | M34057 | | + | + | + | | + | :- |
| LAZ3/BCL6 (=Z79582;D28522/4) | 1 | Z79581 | | | | | | | |
| LDLC | 2 | Z34975 | + | + | + | + | | + | |
| lecithin-cholesterol acyltransferase (LCAT) (non-exact, 66%) | 1 | M17959 | | | | | | | |
| lectin, galactoside-binding, soluble, 2 (galectin 2) (LGALS2) | 1 | M87842 | | | | + | | | |
| lectin, galactoside-binding, soluble, 3 binding protein (galectin 6 binding protein) (LGALS3BP) | 1 | L13210 | + | + | + | + | | + | |
| leucine rich repeat (in FLII) interacting protein 1 (LRRFIP1) | 5 | AJ223075 | + | + | + | + | + | + | |
| leucocyte immunoglobulin- like receptor-5 (LIR-5) | 2 | AF072099 | | | | + | | | |
| leucocyte immunoglobulin- like receptor-6a (LIR-6) | 7 | AF025530 | | | | | | | |
| leucocyte immunoglobulin- like receptor-7 (LIR-7) | 2 | U82275 | | + | | | | | only found in CNS |
| leukemia virus receptor 1 (GLVR1) | 1 | L20859 | + | + | + | + | | + | |
| leukocyte adhesion protein p150,95 alpha subunit | 1 | M29484 | | | | | | | |
| leukocyte antigen, HLA-A2 | 3 | Y13267 | | | | | | | |
| leukocyte immunoglobulin- like receptor (MIR-10) | 3 | AF025528 | | + | | | | | |
| leukocyte tyrosine kinase (LTK) | 1 | X60702 | + | | | | | | found only in blood |
| leukocyte-associated lg- like receptor 1 (LIAR1) | 3 | AF013249 | | | | + | | | |
| leukotriene A4 hydrolase (LTA4H) | 6 | J03459 | + | + | + | + | + | + | |
| leupaxin (LDPL) | 2 | AF062075 | + | <u> </u> | l | + | | + | |
| ligase I, DNA, ATP- dependent (LIG1) | 1 | M36067 | В, Т | + | + | | | + | |
| LIM and SH3 protein 1 (LASP1) | 2 | X82456 | + | + | + | + | + | + | |
| LIM domain kinase 2 (LIMK2) | 2 | AC002073 | + | + | + | + | | + | |
| line-1-protein | 1 | | | 1 | | <u> </u> | | | |
| Line-1 repeat mRNA with 2 open reading frames | 1 | U93566 | + | + | + | + | + | + | |
| Line-1 repeat with 2 open reading frames | 1 | M22332 | + | + | + | + | + | + | high in gastric tumor |
| LINE-1 REVERSE TRANSCRIPTASE HOMOLOG | 1 | P08547 | | | _ | | | | |
| lipase A, lysosomal acid, cholesterol esterase (Wolman disease) (LIPA) | 4 | X76488 | + | + | + | + | | + | |
| lipase, hormone-sensitive (LIPE) | 1 | L11706 | + | + | | | | + | |
| LMP7 | 1 | L11045 | | | | | | | |
| Lon protease-like protein (LONP) | 2 | X74215 | + | + | + | + | · | + | |
| low density lipoprotein- related protein 1 (alpha-2- macroglobulin receptor) (LRP1) | 2 | AF058414 | | | | | + | | only in liver |
| low density lipoprotein- related protein-associated protein 1 (alpha-2- macroglobulin receptor- associated protein 1) (LRPAP1) | 1 | M63959 | | + | + | | + | + | |

WO 00/40749

PCT/CA00/00005

| low density lipoprotein- related protein-associated protein 1 (alpha-2- | 1 | M63959 | | | | | | | |
|---|----|----------|---|--|------------|----------|--|-------|-------|
| macroglobulin receptor- | | | | | | 1 | | | |
| associated protein 1) (LRPAP1) (non-exact, 75%) | | | | | 1 | | | | |
| low-affinity Fc-gamma receptor IIA | 1 | L08107 | | | - | \vdash | +- | _ | |
| LPS-induced TNF-alpha factor (PIG7) | 9 | AF010312 | + | + | + | + | + | + | |
| Lst-1 | 1 | U00921 | + | + | + | + | | + | |
| L-type amino acid transporter subunit LAT1 | 1 | AF104032 | | | ļ <u>.</u> | 1 | \vdash | t | |
| lung resistance-related protein (LRP) | 1 | X79882 | + | + | + | + | | + | |
| Lymphocyte antigen 75 (LY75) | 1 | AF011333 | В | | - | | | | |
| lymphocyte antigen 9 (LY9) | 2 | L42621 | | \dagger | | | \vdash | - | · |
| lymphocyte antigen HLA- B*4402 and HLA-B*5101 | 2 | L42345 | | | | | | | |
| lymphocyte cytosolic protein 1 (L-plastin) (LCP1) | 42 | J02923 | | <u> </u> | | - | | | |
| lymphocyte cytosolic protein 2 (SH2 domain- | 4 | U20158 | | <u> </u> | lymp | hon | a, T | activ | vated |
| containing leukocyte protein of 76kD) (LCP2) | | | | | | | | | |
| lymphocyte glycoprotein | 2 | X04391 | + | | + | | T | Γ- | |
| lymphocyte-specific protein 1 (LSP1) | 16 | M33552 | + | + | + | + | _ | + | |
| lymphocyte-specific protein tyrosine kinase (LCK) | 7 | M36881 | | + | | _ | - | + | |
| lymphoid phosphatase LyP1 | 1 | AF001847 | | | | | | | |
| lymphoid-restricted membrane protein (LRMP) | 4 | U10485 | + | | + | + | | | |
| lymphoid-specific SP100 homolog (LYSP100-A) | 1 | U36500 | | | | | | + | |
| lymphoma proprotein convertase (LPC) | 2 | U33849 | + | + | + | + | | + | |
| PROTECTIVE PROTEIN | 1 | P10619 | | | | | | | - , |
| PRECURSOR (CATHEPSIN A) | | | | | | | | | |
| (CARBOXYPEPTIDASE C) | | 10.4400 | | | | | | | |
| lysosomal-associated membrane protein 1 (LAMP1) | 1 | J04182 | + | + | + | + | + | + | |
| Lysosomal-associated | 1 | J04183 | | + | + | + | + | + | |
| membrane protein 2 (LAMP2) | | | | | | | | | |
| lysozyme (renal amyloidosis) (LYZ) | 39 | M19045 | + | + | + | + | | + | |
| lysyl-tRNA synthetase (KARS) | 2 | D32053 | + | + | + | + | | + | |
| M phase phosphoprotein 10 (U3 small nucleolar | 1 | X98494 | | | | | | | |
| ribonucleoprotein) (MPP- 10) | | | | | ļ | | | | |
| M1-type and M2-type pyruvate kinase | 2 | X56494 | | | | | | | |
| m6A methyltransferase (MT-A70) | 7 | AF014837 | + | + | | + | | | |
| mab-21 (C. elegans)-like 1 (MAB21L1) | 1 | U38810 | | + | + | + | | + | |
| MacMarcks | 1 | X70326 | + | + | + | + | + | + | |
| macrophage-associated antigen (MM130) | 1 | Z22968 | | + | + | + | | + | |

| | | | | | | | | - ' | 21/CA00/00003 |
|--|-------------|--|----------------|--|--|----------|--|------------|--|
| MADS box transcription enhancer factor 2, | 1 | U49020 | | T + | + | + | | + | |
| polypeptide A (myocyte | | | | 1 | | | ŀ | | |
| enhancer factor 2A) | | | | 1 | | 1 | ŀ | 1 | |
| (MEF2A) | | 1 | | | | 1 | 1 | | |
| MADS box transcription | 1 | L08895 | · - | + | + | + | \vdash | + | |
| enhancer factor 2, | | | | 1 | | | 1 | 1 | 1 |
| polypeptide C (myocyte | | | | 1 | Į | | l | | |
| enhancer factor 2C) (MEF2C) | | | | | ļ | | į. | 1 | ļ i |
| major cytoplasmic tRNA- | | X17516 | | - | | ├ | ├ | ├ | |
| Val(IAC) (=M33940) | • | 7 | | | | | | | l |
| major histocompatibility | 1 | M95531 | | + | | | | - | |
| complex class I beta chain | | | | | | | | | 1 1 |
| (HLA-B) | | 700010 | | | | <u> </u> | <u> </u> | <u> </u> | |
| major histocompatibility complex, class I, A (HLA-A) | 41 | Z93949 | + | + | + | + | İ | + | high in villous |
| major histocompatibility | | Z72422 | | | | ! | | — | adenoma |
| complex, class I, A (HLA-A) | • | 212422 | | 1 | | ١. | | | 1 |
| (low match) | | 1 | | | | | l | 1 | [|
| major histocompatibility | 82 | M24097 | + | + | + | + | + | + | |
| complex, class I, C (HAL- | | | | į. | | ł | | | 1 |
| (C) | | - Danasa | | | <u> </u> | <u> </u> | L | L . | ļ |
| complex, class I, E (HLA-E) | 77 | M20022 | + | + | + | + | 1 | + | |
| major histocompatibility | 2 | U15085 | + | + | + | + | ├ | + | |
| complex, class II, DM | _ | 0.0000 | • | ' | • | • | | • | |
| BETA (HLA-DMB) | | | | | | | Ī | 1 | • |
| major histocompatibility | 10 | M57466 | + | + | + | + | | + | |
| complex, class II, DP beta | | | | | | | l | | |
| 1 (HLA-DPB1) | 9 | V00522 | | + | + | + | | | |
| complex, class II, DR beta | 9 | V00522 | + | T . | _ | • | l | + | |
| 1 (HLA-DRB1) | | | | | | | | | |
| Major histocompatibility | 2 | M24070 | | + | + | _ | + | + | |
| complex, class II, Y box- | | | | | | | l | | |
| binding protein I; DNA- | | | | 1 | | | 1 | 1 | |
| binding protein B (YB1) malate dehydrogenase 1, | 1 | D55654 | + | + | + | + | + | + | |
| NAD (soluble) (mdh1) | • | 000004 | • | | • | 1 | " | 1 | |
| malate dehydrogenase 1. | 3 | D55654 | | + | + | 1 | + | + | |
| NAD (soluble) (MDH1) | | | | | | l | 1 | | |
| malonyl-CoA | 2 | AF097832 | | | | | | | |
| decarboxylase precursor | | A C 0 4 0 0 0 0 | | | | | | | |
| maltase-glucoamylase (mg) | 1 | AF016833 | | | | + | | | |
| manic fringe (Drosophila) | 1 | U94352 | + | ++ | +- | + | | + | |
| homolog (MFNG) | • | 037332 | • | | | ' | | ļ ' | |
| mannose phosphate | 1 | X76057 | | ++ | + | + | | + | |
| isomerase (MPI) | | | | ⊥ | | <u> </u> | L | <u> </u> | |
| mannose phosphate | 2 | X76057 | | + | + | + | | + | |
| isomerase (mpi) | | VECAPA | | 1 | | Ь_ | <u> </u> | <u> </u> | |
| mannose-6-phosphate receptor (cation | 3 | X56253 | | + | + | 1 | + | + | |
| dependent) (M6PR) | | | | | | | | | |
| mannose-P-dolichol | 1 | AF038961 | | + | + | + | - | + | |
| utilitzation defect 1 | • | | | | | | | | |
| (MPDU1) | | | | لــــــــــــــــــــــــــــــــــــــ | | L | | | |
| mannosidase, alpha B, lysosomal (MANB) | 1 | U60885 | | + | | + | + | + | |
| mannosyl (alpha-1,3-)- | | M55621 | + | ++ | + | + | + | + | |
| glycoprotein beta-1,2-N- | I | 1VIJJ02 I | ₹ | * | 7 | 🗂 | 7 | * | |
| acetylglucosaminyltransfer | | | | | | | | l | |
| ase (MGAT1) | | <u> </u> | | ⊥ ∣ | | | | | |
| map 4q35 repeat region | 1 | AF064849 | | | | | | | |
| MAP kinase-interacting | 2 | AB000409 | | + | + | + | + | + | |
| serine/threonine kinase 1 | | | | | | | | | |
| (MKNK1) | | | | $oldsymbol{ol}}}}}}}}}}}}}}}}}}$ | | | | | |
| MAP/ERK kinase kinase 3 (MEKK3) | 5 | U78876 | | + | | | | | |
| MAP/ERK kinase kinase 5 | 1 | D84476 | | + | + | | + | | |
| (MEKK5) | ı | 2047/0 | | - | т , | | * | | |
| <u> </u> | | <u> </u> | | <u> </u> | | | | L | <u></u> |

| | | | | | | | | - ' | | |
|---|----------|--|--|--------------|--|--------------|--|--------------|----------------|-------|
| MAP/microtubule affinity- regulating kinase 3 (MARK3) | 4 | M80359 | | 1 | + | | | + | | |
| Marenostrin protein | 1 | Y14441 | | | | ${f 	o}$ | | 1 | | |
| MASL1 | 1 | AB016816 | | + | | | | | - | |
| MAX dimerization protein | 3 | L06895 | | + | | + | | + | | |
| (MAD) MaxiK potassium channel | | AF025046 | | | | <u> </u> | <u> </u> | <u> </u> | | |
| beta subunit | 1 | AF035046 | | | | | Ì | |] | |
| MBP-2 for MHC binding protein 2 | 1 | X65644 | | + | + | + | | + | | |
| Meis (mouse) homolog 3 | 1 | U68385 | | + | + | + | | + | | |
| (MEIS3) melanoma-associated | 1 | M12154 | ļ | 4 | ļ | <u> </u> | <u> </u> | | | |
| antigen p97 | , , | 10112154 | | | ŀ | | ļ | | | |
| (melanotransferrin) | | | <u> </u> | | | | L | <u> </u> | | |
| membrane cofactor protein (CD46, trophoblast- | 4 | X59405 | | + | + | + | | + | | |
| lymphocyte cross-reactive | | | | | | Ī | | | | |
| antigen) (MCP) membrane component, | | D24000 | | <u> </u> | | | | | | |
| chromosome 17, surface | 4 | D14696 | | + | + | + | + | + | | |
| marker 2 (ovarian | • | | | | | ļ | | | | |
| carcinoma antigen CA125) (M17S2) | | | } | | | | | | | |
| membrane metallo- | 2 | J03779 | В | | + | + | + | + | | |
| endopeptidase (neutral endopeptidase, | | | ļ | | | İ | | | • | |
| enkephalinase, CALLA, | | | | | | l | l | | | |
| CD10) (MME) | | | | <u></u> | | | | | | |
| membrane protein, palmitoylated 1 (55kD) | 2 | M64925 | | + | + | + | + | + | | |
| (MPP1) | | | | İ | | | | | | |
| meningioma expressed antigen (MGEA) | 1 | U94780 | | | | + | | | | |
| meningioma-expressed | | U73682 | + | + | | + | + | - | | - |
| antigen 11 (MEA11) | l | | | | | | | | | |
| Menkes Disease (ATP7A) putative Cu++-transporting | 1 | L06133 | | + | | | | | | |
| IP-type ATPase | | | } | | | | | | | |
| metallothionein 2A (MT2A) | 1 | V00594 | | + | + | + | + | + | | |
| metaxin 1 (MTX1) | 1 | U46920 | | + | | + | | + | | |
| methionine adenosyltransferase II. | 2 | X68836 | + | + | + | + | | + | | |
| alpha (MAT2A) | | | | | | | | | , | |
| methyl-CpG binding | 1 | Y10746 | | | | | | | | |
| domain protein 1 (MBD1) (non-exact 59%aa) | | | | | | | | | | |
| methylene tetrahydrofolate | 2 | X16396 | + | + | + | + | | + | | |
| dehydrogenase (NAD+ | | | | | | | | Ì | | |
| methenyitetrahydrofolate | | | | | | | | | | |
| cyclohydrolase (MTHFD2) | | | | | | | | | | |
| methylenetetrahydrofolate dehydrogenase (NADP+ | 1 | J04031 | | + | + | + | + | + | | |
| dependent). | | İ | | | | | | | | |
| methenyltetrahydrofolate cyclohydrolase. | | | | | | | | | | |
| formyltetrahydrofolate | | | | | | | | | | |
| synthetase (MTHFD1) | | | | | | | l | | | |
| methyltransferase, putative | 2 | AJ224442 | | | | | | | | |
| MHC antigen (HLA-B) (=L42024) | 1 | U14943 | | | | | | | | |
| MHC class 1 region | 2 | AF055066 | | ┼─┤ | | | - | \dashv | | |
| MHC class I antigen (HLA- | 1 | U70863 | | | | | | \dashv | | |
| A2) MHC class I antigen (HLA- | 1- | U19736 | | | | | | | | |
| A33) | <u>'</u> | | | | ĺ | | | | | |
| MHC class I antigen (HLA-C) | 1 | U38975 | | T | | | | | | |
| <u></u> | | <u>. </u> | | ┸ | | | | | | |

| WO 00/40/49 | | | | | | | | • | |
|--|---------------|----------|--|----------|---|---|--|---|---------|
| MHC class I antigen B*5801 (HLA-B) | 1 " | U52813 | | | | | | | |
| MHC class I antigen HLA-A (HLA-A) | 2 | AF015930 | | | | | | | |
| MHC class I antigen HLA-A | 1 | U36687 | | | | | | | |
| (HLA-A-2402 allele) MHC class I antigen HLA- | 2 | X13112 | | | | | | | |
| A11K | | 1167224 | | | | | | | |
| MHC class I antigen HLA-B (B*0801 variant) (=AF028596) | 1 | U67331 | | | | | | | |
| MHC class I antigen HLA-B (B*0801 variant) (=U88254) | 1 | U67330 | | | | | | | |
| MHC class I antigen HLA-B (B*48 allele) | 7 | AF017328 | | | | | | | |
| MHC class I antigen HLA-B (HLA-B*1502 allele) | 1 | AF014770 | | | | | | | |
| MHC class I antigen HLA-B (HLA-B*40MD) | 1 | U58643 | | | | | | | |
| MHC class I antigen HLA-B (HLA-B*4103 allele) | 1 | AF028596 | | | | | | | |
| MHC class I antigen HLA-B gene (HLA-B*4402 variant | -1 | AF035648 | | | | | | | |
| Allele) MHC class I antigen HLA-B | 1 | U52175 | | | | | | | |
| GN00110-B*3910 MHC class I antigen HLA- | 1 | D83030 | | | | | | | |
| Cw*04011 MHC class I antigen | 1 | U56434 | | | | | | | |
| R69772 HLA-A (Å*0302) MHC class I antigen | 1 | U58469 | | | | | | | |
| SHCHA (HLA-B*4403 variant) | • | 038409 | | | | | | | · |
| MHC class I histocompatibility antigen (HLA-B) (clone C21/14) | 1 | U06697 | | | | | | | |
| MHC class I HLA B71 | 2 | L07950 | | _ | | | | | |
| MHC class I HLA-A (Aw33.1) | 1 | Flp | | <u> </u> | | | | | |
| MHC class I HLA-B | 1 | U18660 | ······································ | \vdash | | | | | |
| MHC class THLA-B (HLA-B-07ZEL alleie) (=X86704) | 1 | U18661 | | | | | | | |
| MHC class THLA-B (HLA-B-08NR aliele) | 1 | U28759 | | | | | | | |
| MHC class I HLA-B*3512 | 1 | L76094 | | | - | | - | | |
| MHC class I HLA-B41 variant (=U17572) | 3 | U17572 | | | | | | | |
| MHC class I HLA-B44.2 chain | . 1 | M24038 | | | | | - | | |
| MHC class I HLA-B51- cd3.3 | 1 | L41086 | | - | | | | | |
| MHC class I HLA-C allele | 2 | Z33459 | | \vdash | | | | | |
| MHC class I HLA-Cw*0304 (=M84172; M99389) | 1 | D64150 | | | | | | | |
| MHC class I HLA-Cw*0803 | 3 | Z15144 | | | | | - | | |
| MHC class THLA-Cw6 | 1 | M28206 | | | | | | | |
| MHC class I HLA-J antigen | 1 | L56139 | | | | | | | |
| MHC class I lymphocyte antigen A2 (A2.1) variant DK1 | 1 | M19670 | | | | | | | |
| MHC class I mic-B antigen | 1 | X91625 | | | | | | | |
| MHC class I polypeptide- related sequence A (MICA) | 1 | L14848 | ····· | | | + | | | |
| MHC class I protein HLA-C heavy chain (C*0701new allele) (=AF017331) | 1 | U61274 | | | | | | | |
| MHC class II DNA Sequence (clone A37G7- 1C11) | 1 | L18885 | | | | | | | |
| / | | <u> </u> | | | | | | Ь | <u></u> |

| MHC class II DQ-alpha associated with DRw6, DQw1 protein | 1 | M16995 | + | | + | + | | + | |
|---|-----|--------------------|---|--|--------------|----------|----------|--------------|----|
| MHC class II DQ-beta | 2 . | M17564 | | + | - | + | ├ | + | |
| associated with DR2, DQw1 protein | | | | | | | | | |
| MHC class II HAL-DQ- LTR5 (DQ,w8) DNA | 1 | M33842 | | | | | | | |
| fragment, long terminal | | | | | | |] | | |
| repeat region MHC class II hia-dr alpha- | 1 | J00195 | | + | | - | \vdash | | |
| chain (=J00197:M60334:K01117 | | | | | | | l | | |
| 1;J00194;M60333;X00274) | | | | | | | | | |
| MHC class II HLA-DRB1 MHC class II HLA-DRw11- | 1 - | AF007883 M21966 | | ļ | <u> </u> | ļ | | | |
| beta-I chain (DRw11.3) | | L | | | İ | | | | |
| MHC class II lymphocyte antigen (DPw4-beta-1) | 1 | M23907 | | | | | | | |
| MHC CLASS II TRANSACTIVATOR CIITA | 1 | P33076 | | | T | | | | |
| (non-exact 57%) | | | | <u> </u> | | | | | |
| MHC HLA-E2.1 (=X87679) MHC HLA-E2.1 (alpha-2 | 1 1 | M32507 | | ļ | | | | | |
| domain) (low match) | | M32507 | | | | | L | | |
| Mi-2 autoantigen 240 kDa protein (non-exact 84%) | 1 | U08379 | | | | | | | |
| microsomal stress 70 protein ATPase core (stch) | 1 | U04735 | | | 1 | | | _ | |
| microtubule-associated | 1 | U19727 | + | + | + | + | \vdash | + | |
| protein 4 (MAP4) microtubule-associated | 1 | X73882 | | ļ | ļ | ļ | <u> </u> | | |
| protein 7 (MAP7) | | | | <u> </u> | Ĺ | | | | |
| mineralocorticoid receptor (aldosterone receptor) (MLR) | 2 | M16801 | | + | | + | | + | |
| minichromosome maintenance deficient (S. | 1 | X62153 | | + | + | + | | + | |
| cerevisiae) 3 (MCM31) | | AB011144 | ļ | <u> </u> | L. | | | | |
| maintenance deficient (S. | ' | ABUTTIAA | | + | + | + | | + | |
| cerevisiae) 3-associated protein (MCM3AP) | | | | | | | | | |
| minichromosome maintenance deficient (S. | 2 | X74795 | + | + | + | + | + | + | |
| cerevisiae) 5 (cell division | | İ | | ļ | | | | | , |
| cycle 46) (MCM5) mitochondiral cytochrome b | 1 | AF042517 | | ļ | | <u> </u> | | | |
| (CYTB) mitochondrial 16S rRNA | | | | <u> </u> | | | | | , |
| mitochondrial ATP | 11 | Z70759 X59066 | | ļ | | | | | |
| synthase (F1-ATPase) | _ | //55555 | | | | | | | |
| mitochondrial ATP | | X69907 | | ├ | | - | | | |
| synthase c subunit (P1 form) | | | | | | | | | |
| mitochondrial cytochrome b (CYTB) | 6 | AF042508 | | | | | | | |
| mitochondrial cytochrome b small subunit of complex II | 1 | AB006202 | | | | | - | | |
| mitochondrial CYTOCHROME C | 1 | P00395 | | | | | | | |
| OXIDASE POLYPEPTIDE I | | | | | | | _ | | |
| mitochondrial CYTOCHROME C | 1 | P00403 | | | | | | | |
| OXIDASE POLYPEPTIDE | | | | | | | | | |
| mitochondrial cytochrome C oxidase subunit II | 2 | P00403 | | · | | | | | |
| | | | 1 | | | | | - 1 | Į. |

| mitochondrial cytochrome oxidase subunit II (COII) (=U12692 Hsa4 | 5 | U12691 | | | | | | | | | |
|--|----|------------------|---------------|----------------|---|----------|--------------|--------------|--|---|-------------|
| mitochondrion cytochrome oxidase subunit II) | | | | | | | | | | | |
| mitochondrial DNA loop attachment sequences (clone LAS34) | 1 | X89763 | | | | | | | | | |
| mitochondrial DNA polymerase accessory subunit precursor (MtPolB) nuclear gene encoding | 1 | U94703 | | + | | | | | | | |
| mitochondrial protein, | 1 | X93334 | | - | | <u> </u> | | ├ | | | |
| complete genome mitochondrial genes for | 8 | V00710 | | ┼— | _ | | | | - | | |
| several tRNAs (Phe, Val, Leu) and 12S and 16S ribosomal RNAs. | | | | | | | | | | | |
| mitochondrial genes for tRNA (Phe) and 12S rRNA (fragment) | 3 | V00660 | | | | | | | | | |
| mitochondrial inner membrane preprotein translocase Tim17a | 1 | AF106622 | | | | | | | | | |
| mitochondrial isolate Afr7 cytochrome b(CYTB) | 1 | AF042503 | | | - | ┢ | - | | | | |
| mitochondrial loop attachment sequence (clone LAS88) | 1 | X89843 | | | | | | | | · | |
| mitochondrial NADH dehydrogenase subunit 2 (ND2) | 14 | AF014893 | | | | | | | | | |
| mitochondrial translational initiation factor 2 (MTIF2) | 1 | L34600 | | + | + | + | | + | | | |
| mitochondrion cytochrome b | 1 | U09500 | | | | | | | | | |
| mitogen inducible gene mig-2 | 1 | Z24725 | | + | + | + | | + | | | |
| mitogen inducible gene mig-2 (non-exact, 71%) | 1 | Z24725 | | | | | | | | | |
| mitogen-activated protein kinase-activated protein kinase 3 (MAPKAPK3) | 2 | U43784 | | + | + | + | | + | | | |
| MLN51 | 2 | X80199 | | + | + | + | + | + | | | |
| MLN64 (=D38255 CAB1) | 1 | X80198 | + | + | + | + | | | | | |
| moesin (MSN) | 14 | M69066 | + | + | + | + | | + | | | |
| monocytic leukaemia zinc finger protein (MOZ) | 2 | U47742 | | -+ | + | + | | + | | | |
| MOP1 () | 2 | U29165 | | | | | | | | | |
| motor protein (Hs.78504) | 2 | D21094 | + | + | + | + | | + | | | |
| mouse double minute 2, human homolog of; p53- binding protein (MDM2) | | U39736 | | | + | + | | | | | |
| M-phase phosphoprotein 6 (MPP-6) | 1 | X98263 | | + | + | + | | + | | | |
| M-phase phosphoprotein, mpp11 | 1 | X98260 | | | | | | | | | |
| MPS1 | 1 | L20314 | | | | | | | | | |
| Mr 110,000 antigen MRC OX-2, V-like region | 2 | D64154 X05324 | | + | | + | + | + | | | |
| (=M17227) mu-adaptin-related protein- | | l | . | | | | | | | | |
| 2; mu subunit of AP-4 (MU- ARP2) | | Y08387 | | | | | | | | | |
| multifunctional polypeptide similar to SAICAR synthetase and AIR carboxylase (ADE2H1) | 1 | X53793 | + | + | + | + | | + | | | |
| January (ADEZIII) | | 1 | | 1 1 | | | 1 | ı | | | |

WO 00/40749

| WO 00/40749 | | | | | | | | | |
|---|----|----------|---|---|---|---|---|---|---|
| murine leukemia viral (bmi- 1) oncogene homolog (BMI1) | 1 | L13689 | | + | | + | | + | ; |
| mutant (Daudi) beta2 - | 44 | X07621 | | | | | | | |
| mutated in colorectal cancers (MCC) | 1 | M62397 | | + | + | | | + | |
| myeloid cell leukemia sequence 1 (BCL2-related) (MCL1) | 9 | L08246 | + | + | + | + | + | | |
| myeloid cell nuclear differentiation antigeN (MNDA) | 11 | M81750 | + | | | | | + | |
| myeloid differentiation primary response gene (88) (MYD88) | 4 | U70451 | | + | + | + | | + | |
| myeloid leukemia factor 2 (MLF2) | 3 | U57342 | | + | | + | | + | |
| myeloid/lymphoid or mixed- lineage leukemia (trithorax (Drosophila) homolog); translocated to, 7 (MLLT7) | 8 | U89867 | | + | + | + | | + | |
| MYH9 (cellular myosin heavy chain) | 1 | M81105 | | | | | | | |
| myomesin (M-protein) 2 (165kD) (MYOM2) | 1 | X69089 | | | | | | | |
| myosin IE (MYO1E) | 11 | X98411 | | + | | + | | | |
| myosin light chain kinase (MLCK) | 1 | U48959 | + | | + | + | | + | |
| myosin phosphatase, target subunit 1 (MYPT1) | 2 | D87930 | | + | + | + | | + | |
| myosin regulatory light chain (=U26162) | 2 | D50372 | | | | | | | |
| myosin VIIa (low match 71) | 1 | U55208 | | | | | | | |
| myosin, heavy polypeptide 9, non-muscle (MYH9) | 3 | M81105 | + | + | + | + | | + | |
| myosin, light polypeptide, regulatory, non-sarcomeric (20kD) (MLCB) | 6 | X54304 | + | + | + | + | + | + | |
| myosin-l beta | 1 | X98507 | + | + | + | + | | + | |
| myristoylated alanine-rich protein kinase C substrate (MARCKS, 80K-L) (MACS) | 1 | D10522 | | + | + | | | | |
| myxovirus (influenza) resistance 1, homolog of murine (interferon-inducible protein p78) (MX1) | 1 | M30817 | + | + | + | + | | + | |
| myxovirus (influenza) resistance 2, homolog of murine (MX2) | 3 | M30818 | | | + | | | | |
| N-acetylgalactosaminidase, alpha- (NAGA) | 2 | M62783 | | + | + | | + | + | |
| N-acetylglucosamine receptor 1 (thyroid) (NAGR1) | 1 | L03532 | | + | + | + | | + | |
| NACP/alpha-synuclein | 2 | U46896 | | | | | | | |
| N-acylaminoacyl-peptide hydrolase (APEH) | 1 | D38441 | | + | + | | + | + | |
| N-acylsphingosine amidohydrolase (acid ceramidase) (ASAH) | 11 | U47674 | + | + | + | + | | + | |
| NAD+-specific isocifrate dehydrogenase beta subunit precursor (encoding mitochondrial protein) | 1 | U49283 | + | + | + | + | + | + | |
| NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 5 (13kD, B13) (NDUFA5) | 1 | U53468.1 | + | + | + | + | + | + | |

| NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 5 (16kD, | 1 | AF047181 | | + | + | + | + | + | |
|--|----|-----------|---|----------|---|---|-------------|--------------|----------------|
| SGDH) (NDUFB5) | | | | | | | | | |
| NADH dehydrogenase (ubiquinone) Fe-S protein 2 (49kD) (NADH-coenzyme Q eductase) (NDUFS2) | 1 | AF050640 | | + | + | + | + | + | |
| NADH dehydrogenase (ubiquinone) flavoprotein 2 (24kD) (NDUFV2) | 1 | M22538 | | | + | + | + | + | |
| NADH:ubiquinone dehydrogenase 51 kDa subunit (NDUFV1) | 2 | AF053070 | + | + | + | + | + | + | |
| NADH-CYTOCHROME B5 REDUCTASE (B5R) (50%aa) | 1 | P00387 | | | | | | | |
| NADH-ÚBIQUINONE OXIDOREDUCTASE CHAIN 1 | 1 | P03886 | | | | | | | |
| Nardilysin (N-arginine dibasic convertase) (NRD1) | 2 | U64898 | + | + | + | + | | + | |
| nascent-polypeptide- associated complex alpha polypeptide (NACA) | 5 | X80909 | | + | + | | + | + | |
| natural killer cell group 7 sequence (NKG7) | 8 | S69115 | | | | + | | + | |
| natural killer cell transcript 4 (NK4) | 19 | M32011 | + | | | | | | |
| natural killer-associated transcript 3 (NKAT3) | 1 | U30274 | + | <u> </u> | | | | t | blood only |
| natural killer-associated | 1 | AF022045 | + | <u> </u> | | - | | - | blood only |
| transcript 5 (NKAT5) natural killer-tumor | 1 | L04288 | В | | + | _ | + | + | |
| recognition sequence (NKTR) | | | | 1 | | | | | |
| N-deacetylase/N- sulfotransferase (heparan glucosaminyl) 2 (NDST2) | 2 | AF042084 | + | + | | + | | + | |
| Ndr protein kinase | 3 | Z35102 | | + | | | | | |
| Nedd-4-like ubiquitin- | 1 | U96113 | | \vdash | | | | \vdash | |
| protein ligase WWP1 nel (chicken)-like 2 | 3 | D83018 | | + | + | | | | |
| (NELL2) N-ethylmaleimide-sensitive factor attachment protein, | 1 | U39412 | | + | | | + | | |
| alpha (NAPA) N-ethylmaleimide-sensitive factor attachment protein, gamma (NAPG) | 1 | U78107 | | + | + | + | | | |
| neural precursor cell expressed, developmentally down- | 3 | X92544 | + | + | + | + | | + | high in testis |
| regulated 5 (NEDD5) neural precursor cell expressed, developmentally down- | 1 | D23662 | + | + | + | + | Ŧ | + | |
| regulated 8 (NEDD8) neuregulin 1 (NRG1) | 1 | U02330 | | + | | + | + | - | |
| neuroblastoma RAS viral (v-ras) oncogene homolog (NRAS) | 4 | AB020692 | + | + | + | + | | + | |
| Neuroblastoma RAS viral (v-ras) oncogene homolog (NRAS) (low match) | 1 | X68286 | , | | | | - | | |
| Neurofibromin 2 (bilateral acoustic neuroma) (NF2) | 1 | S73853 | | + | | | | + | |
| neuronal apoptosis | 2 | U19251 | + | + | + | | - | + | |
| inhibitory protein (NAIP) | | X00003343 | | + | + | + | <u> </u> | + | |
| neuronal cell adhesion molecule (NRCAM) | 1 | AB002341 | ľ | Į T | 7 | | | 1 ' | |

| VVO 00/40/49 | | | | <u> </u> | | - | | | |
|--|----|----------|---|----------|---|---------------|----------|---|--|
| neuropathy target esterase (NTE) | 1 | AJ004832 | | + | + | + | | + | |
| neuropeptide Y3 receptor, 5'UTR (low score) | 1 | D28433 | | | | | | | |
| neurotrophic tyrosine kinase, receptor, type 1 (NTRK1) | 14 | X03541 | + | + | + | + | + | + | |
| neutrophil cytosolic factor 4 (40kD) | 2 | U50720 | | | | | | | |
| NG31 | 1 | AF129756 | | | | | <u> </u> | | |
| NGAL (=X83006) | 1 | X99133 | | | | | | | |
| nibrin (NBS) | 1 | AF051334 | | | | | | | |
| NIK | 1 | AB014587 | | + | + | + | | + | |
| Ninjurin 1; nerve injury- induced protein-1 | 1 | U72661 | | + | + | + | | + | |
| nitrilase 1 (NIT1) (=AF069984) | 1 | AF069987 | | | | | | | |
| NKG2-D (low match) (non- exact, 58%) | 1 | X54870 | | | | | | | |
| Nmi | 1 | U32849 | | | | | | | l |
| N-myristoyltransferase 1 (NMT1) | 1 | AF043324 | | + | + | + | + | + | |
| No arches-like (zebrafish) zinc finger protein (NAR) | 1 | U79569 | | + | + | + | | + | |
| non-histone chromosome protein 2 (S. cerevisiae)- like 1 (NHP2L1) | 1 | D50420 | + | + | + | + | + | + | |
| non-muscle (fibroblast) tropomyosin | 1 | | | | | | | | |
| non-muscle alpha-actinin | 1 | U48734 | | | | | <u> </u> | | |
| non-muscle myosin alkali light chain (Hs.77385) | 3 | M22918 | + | + | + | + | + | + | High in fetal adrenal gland and BPH stroma |
| non-neuronal enolase (EC 4.2.1.11) | 1 | X16289 | | | | | | | |
| non-receptor tyrosine phosphatase 1 | 1 | M33689 | | | | | | | |
| normal keratinocyte substraction library mRNA, clone H22a | 3 | X53778 | + | + | + | * | + | + | high in many libraries |
| notch group protein (N) | 3 | M99437 | | | | | | | |
| novel protein | 1 | X99961 | | | | | | | |
| novel T-cell activation protein | 1 | X94232 | | + | + | + | | + | |
| N-ras protein NRU | 7 | A60196 | | | | | | | |
| N-sulfoglucosamine sulfohydrolase (sulfamidase) (SGSH) | 1 | U60111 | | + | | | | + | |
| nsulin induced gene 1 (INSIG1) | 7 | U96876 | + | + | + | + | + | + | |
| ntegrin, alpha 4 (antigen CD49D, alpha 4 subunit of VLA-4 receptor) (ITGA14) | 3 | L12002 | + | | | + | | | |
| nterferon, gamma-inducible protein 16 (IF116) | 1 | M63838 | + | + | + | + | | + | |
| nterleukin 1, beta (IL1RB) | 1 | M15330 | | 1 | | | | T | |
| nuclear antigen H731-like | 2 | U83908 | | + | + | + | | + | |
| nuclear antigen Sp100 (SP100) | 4 | U36501 | + | | | + | + | + | |
| Nuclear antigen Sp100 (SP100) (85%aa) | 1 | P23497 | | | | | | | |
| Nuclear antigen Sp100 (SP100) (89%aa) | 1 | P23497 | * | | | | | | · |
| nuclear autoantigenic sperm protein (histone- binding) (NASP) | 1 | M97856 | + | | + | | | | |

| nuclear corepressor KAP-1 (KAP-1) (=U95040; X97548 | -1- | U78773 | | | | | | | |
|---|---------------------------------------|----------|----------------|--|----------|--|--|-----|---------------------------------------|
| TIF1beta zinc finger | | | | | | 1 | | | |
| protein) | | | | | | | | | |
| Nuclear domain 10 protein (NDP52) | 4 | U22897 | + | + " | + | + | + | + | |
| Nuclear factor (erythroid- | 1 | S74017 | | + | + | + | + | + | |
| derived 2)-like 2 (NFE2L2) Nuclear factor of kappa | | M58603 | | + | + | - | + | + | |
| light polypeptide gene | _ | 1,100000 | | ' | , i | l | · | ļ · | |
| enhancer in B-cells 1 (p105) (NFKB1) | | · | | | | | | ľ | |
| nuclear factor of kappa | 3 | M69043 | | + | + | + | | + | |
| light polypeptide gene enhancer in B-cells | | | | | | | | | |
| inhibitor, alpha (NFKBIA) nuclear factor related to | 1 | U08191 | | + | + | + | - | + | |
| kappa B binding protein (NFRKB) | | | | | | | | | |
| nuclear mitotic apparatus protein 1 (NUMA1) | 3 | Z11583 | + | + | + | + | + | + | |
| nuclear receptor coactivator 2 (GRIP1) | 1 | X97674 | | | | | | | |
| nuclear receptor | 2 | AF010227 | + | + | + | - | | + | |
| coactivator 3 (AIB3) | - | | | | | | | | |
| nuclear receptor coactivator 4 (ELE1) | 22 | X77548 | | + | + | + | + | + | |
| nuclear receptor interacting protein 1 (NRIP1) | 1 | X84373 | | + | | + | | + | |
| nuclear respiratory factor 1 (NRF1) | 1 | U02683 | В | + | + | | | | |
| nuclear RNA helicase, | 4 | U90426 | + | + | + | + | | + | |
| DECD variant of DEAD box family (DDXL) | | | | | | | | | |
| nuclear transcription factor Y, alpha (NFYA) | 1 | X59711 | В | | | | | | |
| nuclear transcription factor, X-box binding 1 (NFX1) | 3 | U15306 | | + | ** | | + | | |
| nuclear transport factor 2 (placental protein 15) (PP15) | 1 | X07315 | + | + | + | + | | + | |
| nucleobindin (=M96824) | 1 | U31336 | | | | | | | |
| nucleobindin 1 (NUCB1) | 2 | M96824 | + | + | + | + | | + | |
| nucleolar phosphoprotein p130 (P130) | 1 | Z34289 | | + | + | | | | |
| nucleolar protein (KKE/D | 1 | Y12065 | + | + | + | + | | + | |
| repeat) (NOP56) nucleolar protein (MSP58) | | AF015308 | | | | | | | |
| nucleolar protein 1 (120kD) | | M32110 | + - | + | | - | | _ | |
| (NOL1) | | | | | | | | | |
| nucleolar protein p40 | 1 | U86602 | + | + | + | + | | + | |
| nucleolin (NCL) | 2 | M60858 | + | + | + | + | | + | |
| nucleophosmin (nucleolar phosphoprotein B23, numatrin) (NPM1) | 14 | M28699 | + | + | + | + | | + | |
| nucleophosmin-retinoic | 1 | U41742 | | | | | | | |
| acid receptor alpha fusion protein NPM-RAR long form | | | | | | | | | |
| nucleoporin (NUP358) | 2 | L41840 | | | | | \vdash | | |
| (=D42063 RanBP2 (Ran- binding protein 2)) | | | | | | | | | |
| nucleoporin 153kD (NUP153) | 1 | Z25535 | | | | | | | |
| nucleoporin 98kD (NUP98) | 1 | U41815 | | + | | - | | | |
| nucleosome assembly | 1 | D28430 | | | | | | | |
| protein nucleosome assembly | 1 | M86667 | | + | + | + | _ | + | |
| protein 1-like 1 (NAP1L1) nucleosome assembly | · · · · · · · · · · · · · · · · · · · | U77456 | + | + | <u> </u> | + | | + | · · · · · · · · · · · · · · · · · · · |
| protein 1-like 4 (NAP1L4) | | 577430 | | | | | | · | |

WO 00/40749

| Taxabaaaaaa aaaaaa aa | | | | | | | | | |
|---|--|---|---|---|-----------|---|-------------|--|---|
| nucleosome assembly protein, 5'UTR | 1 | D28430 | | | | | | | |
| olfactory receptor (OR7- 141) | 1 | U86281 | | | | | | | |
| OLFACTORY RECEPTOR- LIKE PROTEIN HGMP07E (OR17-4) (non-exact 65%) | 1 | P34982 | | | | | | | * |
| oligodendrocyte myelin glycoprotein (OMG) | 7 | L05367 | | + | | | † | | |
| O-linked N- acetylglucosamine (GlcNAc) transferase (UDP-N- acetylglucosamine:polypep tide-N-acetylglucosaminy) | 1 | U77413 | + | + | | + | + | + | |
| transferase) (OGT) oncofetal trophoblast glycoprotein 5T4 precursor | 1 | A53531 | | | | | | | |
| (non-exact 55%) Oncogene TIM (TIM) (non- exact 84%) | 1 | U02082 | | - | | - | - | | |
| ORF (Hs.77868) | 1 | M68864 | + | + | + | + | + | + | |
| ORF1; MER37; putative transposase similar to pogo element Length = 454 | 1 | U49973 | | | | | | | |
| origin recognition complex, subunit 2 (yeast homolog)- like (ORC2L) | 2 | U27459 | | | | + | | | · |
| origin recognition complex, subunit 4 (yeast homolog)- like (ORC4L) (low match) | 1 | AF022108 | | | | | | | |
| ornithine aminotransferase (gyrate atrophy) (OAT) | 2 | M23204 | | + | + | + | | | |
| ornithine decarboxylase (ODC) | 1 | M20372 | | | | | | | |
| omithine decarboxylase | 11 | D78361 | | | _ | - | | | Track in a second |
| antizyme, ORF 1 and ORF 2 | •• | D/6361 | + | + | + | + | + | + | High in pancreas, and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) | 2 | U07132 | + | + | + | + | _ | + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor | | | | | | | + | | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM- | 2 | U07132 | + | + | + | + | | + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM- 40) ovel centrosomal protein | 2 | U07132 AB002806 | + | + | + | + | | + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM- 40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast | 2 6 1 | U07132 AB002806 D28381 | + | + | + | + | | + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM- 40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast tumor protein oviductal glycoprotein 1, | 2 6 1 | U07132 AB002806 D28381 AB008515 | + | + | + | + | | + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM-40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast tumor protein oviductal glycoprotein 1, 120kD (OVGP1) oxidase (cytochrome c) | 2 6 1 1 | U07132 AB002806 D28381 AB008515 L34839 | + | + | + + | + + + | + | + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM-40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast tumor protein oviductal glycoprotein 1, 120kD (OVGP1) | 2 6 1 1 1 | U07132 AB002806 D28381 AB008515 L34839 U09550 | + | + + | + + | + + + | + | + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM-40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast tumor protein oviductal glycoprotein 1, 120kD (OVGP1) oxidase (cytochrome c) assembly 1-like (OXAIL) oxoglutarate dehydrogenase (lipoamide) | 2 6 1 1 1 1 1 1 1 | U07132 AB002806 D28381 AB008515 L34839 U09550 X80695 | + | + + | + + + + | + + + | + + | + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM-40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast tumor protein oviductal glycoprotein 1, 120kD (OVGP1) oxidase (cytochrome c) assembly 1-like (OXAIL) oxoglutarate dehydrogenase (lipoamide) (OGDH) oxysterol binding protein | 2 6 1 1 1 1 1 1 1 4 | U07132 AB002806 D28381 AB008515 L34839 U09550 X80695 D10523 | + | + | + + + + | + + + | + + + + | + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM-40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast tumor protein oviductal glycoprotein 1, 120kD (OVGP1) oxidase (cytochrome c) assembly 1-like (OXAIL) oxoglutarate dehydrogenase (lipoamide) (OGDH) oxysterol binding protein (OSBP) | 2 6 1 1 1 1 1 | U07132 AB002806 D28381 AB008515 L34839 U09550 X80695 D10523 M86917 | + | + | + + + + + | + + + + | + + + + | + + + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM-40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast tumor protein oviductal glycoprotein 1, 120kD (OVGP1) oxidase (cytochrome c) assembly 1-like (OXAIL) oxoglutarate dehydrogenase (lipoamide) (OGDH) oxysterol binding protein (OSBP) OZF (non-exact zinc finger) p21/Cdc42/Rac1-activated kinase 1 (yeast Ste20-related) (PAK1) | 2 6 1 1 1 1 4 | U07132 AB002806 D28381 AB008515 L34839 U09550 X80695 D10523 M86917 X70394 X70394 U51120 | + | + | + + + + + | + + + + | + + + + | + + + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM-40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast tumor protein oviductal glycoprotein 1, 120kD (OVGP1) oxidase (cytochrome c) assembly 1-like (OXAIL) oxoglutarate dehydrogenase (lipoamide) (OGDH) oxysterol binding protein (OSBP) OZF OZF (non-exact zinc finger) p21/Cdc42/Rac1-activated kinase 1 (yeast Ste20- related) (PAK1) P35-related protein (= S80990 ficolin) | 2 6 1 1 1 1 4 | D07132 AB002806 D28381 AB008515 L34839 U09550 X80695 D10523 M86917 X70394 X70394 U51120 D63392 | + | + + + + + + + + | + + + + + | + + + + | + + + + | + + + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM-40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast tumor protein oviductal glycoprotein 1, 120kD (OVGP1) oxidase (cytochrome c) assembly 1-like (OXAIL) oxoglutarate dehydrogenase (lipoamide) (OGDH) oxysterol binding protein (OSBP) OZF OZF (non-exact zinc finger) p21/Cdc42/Rac1-activated kinase 1 (yeast Ste20- related) (PAK1) P35-related protein (= S80990 ficolin) | 2 6 1 1 1 1 4 1 1 2 | U07132 AB002806 D28381 AB008515 L34839 U09550 X80695 D10523 M86917 X70394 X70394 U51120 | + | + + + + + + + + | + + + + + | + + + + | + + + + | + + + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM-40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast tumor protein oviductal glycoprotein 1, 120kD (OVGP1) oxidase (cytochrome c) assembly 1-like (OXAIL) oxoglutarate dehydrogenase (lipoamide) (OGDH) oxysterol binding protein (OSBP) OZF OZF (non-exact zinc finger) p21/Cdc42/Rac1-activated kinase 1 (yeast Ste20-related) (PAK1) P35-related protein (= S80990 ficolin) p40 p40phox (=U50720) | 2 6 1 1 1 1 1 1 1 2 | D07132 AB002806 D28381 AB008515 L34839 U09550 X80695 D10523 M86917 X70394 X70394 U51120 D63392 U93569 X77094 | + | + + + + + + + + | + + + + + | + + + + | + + + + | + + + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM-40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast tumor protein oviductal glycoprotein 1, 120kD (OVGP1) oxidase (cytochrome c) assembly 1-like (OXAIL) oxoglutarate dehydrogenase (lipoamide) (OGDH) oxysterol binding protein (OSBP) OZF (non-exact zinc finger) p21/Cdc42/Rac1-activated kinase 1 (yeast Ste20-related) (PAK1) P35-related protein (= S80990 ficolin) p40 p40phox (=U50720) P47 LBC oncogene | 2 6 1 1 1 1 4 1 1 2 | U07132 AB002806 D28381 AB008515 L34839 U09550 X80695 D10523 M86917 X70394 X70394 U51120 D63392 U93569 X77094 U03634 | + | + + + + + + + + | + + + + + | + + + + | + + + + | + + + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM-40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast tumor protein oviductal glycoprotein 1, 120kD (OVGP1) oxidase (cytochrome c) assembly 1-like (OXAIL) oxoglutarate dehydrogenase (lipoamide) (OGDH) oxysterol binding protein (OSBP) OZF OZF (non-exact zinc finger) p21/Cdc42/Rac1-activated kinase 1 (yeast Ste20-related) (PAK1) P35-related protein (= S80990 ficolin) p40 p40phox (=U50720) P47 LBC oncogene p53-induced protein (PIG11) | 2 6 1 1 1 1 1 1 1 2 | D07132 AB002806 D28381 AB008515 L34839 U09550 X80695 D10523 M86917 X70394 X70394 U51120 D63392 U93569 X77094 U03634 AF010315 | + | + + + + + + + + | + + + + + | + + + + | + + + + | + + + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM-40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast tumor protein oviductal glycoprotein 1, 120kD (OVGP1) oxidase (cytochrome c) assembly 1-like (OXAIL) oxoglutarate dehydrogenase (lipoamide) (OGDH) oxysterol binding protein (OSBP) OZF (non-exact zinc finger) p21/Cdc42/Rac1-activated kinase 1 (yeast Ste20-related) (PAK1) P35-related protein (= S80990 ficolin) p40 p40phox (=U50720) P47 LBC oncogene p53-induced protein | 2 6 1 1 1 1 4 1 1 2 | U07132 AB002806 D28381 AB008515 L34839 U09550 X80695 D10523 M86917 X70394 X70394 U51120 D63392 U93569 X77094 U03634 | + | + + + + + + | + + + + + | + | + + + + | + + + | and activated T cells |

| p62 nucleoporin | 1 | X58521 | | Τ - | Γ | Т | T | Τ | T |
|---|----------------------------------|--|--------------|-------------|----------|--|----------|----------------|------------------------|
| p63 mRNA for | 1 | X69910 | + | ++ | + | + | - | ++ | |
| transmembrane protein | L | | | | | | 1 | | ; |
| PAC clone DJ0701016 from 7q33-q36 (non-exact | 1 | Q07108 | | | | | | Т | |
| 54%) | 1 | | | | | | 1 | | |
| palmitoyl-protein | 10 | U44772 | | + | + | + | ╁╌ | + | |
| thioesterase (ceroid- | | | | | | 1 | | | |
| lipofuscinosis, neuronal 1, infantile: Haltia-Santavuori | | | | | | | | İ | |
| disease) (PPT) | ŀ | | | | | l | | | |
| papillary renal cell | 1 | X99720 | + | + | + | + | ╁ | + | <u> </u> |
| carcinoma (translocation- | | | | 1 | | | 1 | | 1 |
| associated) (PRCC) | | | | | | | | | |
| PAR protein | 1 | AF115850 | | + | | + | | | |
| partial EST (clone c-1gh04) | 1 | Z43627 | | | | П | | | |
| PAX3/forkhead | 1 | U02368 | | | | | | | |
| transcription factor gene fusion | | | | | | | | • | |
| paxillin (PXN) | 4 | D86862 | | + | + | + | <u> </u> | ├ ┯ | |
| PBK1 protein | 2 | AJ007398 | + | + | + | + | ┞ | | |
| PBS-EST (nz92e01.s1 | 1 | A3007396 AA732534 | | ↓ | <u> </u> | | _ | + | |
| NCI_CGAP_GCB1 clone | ' | M132034 | | | | 1 | | | 1 |
| IMAGE:1302936) (low | | 1 | | | | 1 | | | · |
| score) | | | | | | 1 | | | |
| PDZ domain protein (Drosophila inaD-like) | 1 | AJ224747 | + | | | + | | + | |
| (INALD) | • | | | 1 1 | | i i | 1 | l | |
| PEBP2aC Runt domain | 1 | Z38108 | | | | ╁ | | ├ | |
| encoding gene (=Z35728) | | | • | | İ | | 1 | | |
| peptidase D (PEPD) | 1 | J04605 | | | | | | | |
| peptidylprolyl isomerase A | 3 | Y00052 | | + | + | + | + | + | high in many libraries |
| (cyclophilin A) (PPIA) peptidylprolyl isomerase D | 2 | 144007 | | | | | _ | | |
| (cyclophilin D) (PPID) | 4 | L11667 | Т | + | + | ļ | + | + | |
| peptidylprolyl isomerase E | 1 | AF042386 | | + | + | | + | + | |
| (cyclophilin E) (PPIE) | L | | | اللل | | | | | |
| PERB11.1 (=U56942 MHC class I chain-related protein | 1 | U69630 | | | | | | | |
| A) | ſ | Į I | | 1 1 | | 1 | | | |
| | 1 | 1 | | 1 1 | | | 1 | | |
| perforin 1 (preforming | 14 | M28393 | | - | | | | | |
| protein) (PRF1) | 14 | | | | | | | | |
| protein) (PRF1) peroxisomal acyl-CoA | | M28393 X86032 | | | | | | | |
| protein) (PRF1) peroxisomal acyl-CoA thioesterase (PTE1) | 14 | X86032 | | + | - | | | - | |
| protein) (PRF1) peroxisomal acyl-CoA thioesterase (PTE1) Peroxisomal acyl- coenzyme A oxidase | 14 | | | + | + | + | + | + | |
| protein) (PRF1) peroxisomal acyl-CoA thioesterase (PTE1) Peroxisomal acyl- coenzyme A oxidase peroxisomal famesylated | 14 | X86032 | | + | + | + | + | + | |
| protein) (PRF1) peroxisomal acyl-CoA thioesterase (PTE1) Peroxisomal acyl- coenzyme A oxidase peroxisomal farmesylated protein (PXF) | 14 2 1 | X86032 X71440 X75535 | | | | | | | |
| protein) (PRF1) peroxisomal acyl-CoA thioesterase (PTE1) Peroxisomal acyl- coenzyme A oxidase peroxisomal famesylated protein (PXF) phorbol-12-myristate-13- | 14 | X86032 X71440 | B, W | | | | | | |
| protein) (PRF1) peroxisomal acyl-CoA thioesterase (PTE1) Peroxisomal acyl- coenzyme A oxidase peroxisomal tamesylated protein (PXF) phorbol-12-myristate-13- acetate-induced protein (PMAIP1) | 14 2 1 | X86032 X71440 X75535 | B, W | | | | | | |
| protein) (PRF1) peroxisomal acyl-CoA thioesterase (PTE1) Peroxisomal acyl- coenzyme A oxidase peroxisomal famesylated protein (PXF) phorbol-12-myristate-13- acetate-induced protein (PMAIP1) phosphate carrier | 14 2 1 | X86032 X71440 X75535 | B, W | | | | | | |
| protein) (PRF1) peroxisomal acyl-CoA thioesterase (PTE1) Peroxisomal acyl- coenzyme A oxidase peroxisomal famesylated protein (PXF) phorbol-12-myristate-13- acetate-induced protein (PMAIP1) phosphate carrier (mitochondrial gene?) | 14 2 1 1 1 | X86032 X71440 X75535 D90070 X77337 | <u> </u> | + | + | + | | + | |
| protein) (PRF1) peroxisomal acyl-CoA thioesterase (PTE1) Peroxisomal acyl- coenzyme A oxidase peroxisomal famesylated protein (PXF) phorbol-12-myristate-13- acetate-induced protein (PMAIP1) phosphate carrier | 14 2 1 1 1 | X86032 X71440 X75535 D90070 | B, W | | | | | | |
| protein) (PRF1) peroxisomal acyl-CoA thioesterase (PTE1) Peroxisomal acyl- coenzyme A oxidase peroxisomal farnesylated protein (PXF) phorbol-12-myristate-13- acetate-induced protein (PMAIP1) phosphate carrier (mitochondrial gene?) Phosphate carrier, mitochondrial (PHC) phosphate | 14 2 1 1 1 1 | X86032 X71440 X75535 D90070 X77337 | <u> </u> | + | + | + | | + | |
| protein) (PRF1) peroxisomal acyl-CoA thioesterase (PTE1) Peroxisomal acyl- coenzyme A oxidase peroxisomal famesylated protein (PXF) phorbol-12-myristate-13- acetate-induced protein (PMAIP1) phosphate carrier (mitochondrial gene?) Phosphate carrier, mitochondrial (PHC) phosphate cytidylytransferase 1, | 14 2 1 1 1 1 3 | X86032 X71440 X75535 D90070 X77337 X60036 | + | + | + | + | + | + | |
| protein) (PRF1) peroxisomal acyl-CoA thioesterase (PTE1) Peroxisomal acyl- coenzyme A oxidase peroxisomal famesylated protein (PXF) phorbol-12-myristate-13- acetate-induced protein (PMAIP1) phosphate carrier (mitochondrial gene?) Phosphate camer, mitochondrial (PHC) phosphate cytidylytransferase 1, choline, alpha isoform | 14 2 1 1 1 1 3 | X86032 X71440 X75535 D90070 X77337 X60036 | + | + | + | + | + | + | |
| protein) (PRF1) peroxisomal acyl-CoA thioesterase (PTE1) Peroxisomal acyl- coenzyme A oxidase peroxisomal famesylated protein (PXF) phorbol-12-myristate-13- acetate-induced protein (PMAIP1) phosphate carrier (mitochondrial gene?) Phosphate carrier, mitochondrial (PHC) phosphate cytidylytransferase 1, choline, alpha isoform (PCYT1A) PHOSPHATIDATE | 14 2 1 1 1 3 | X86032 X71440 X75535 D90070 X77337 X60036 L28957 | + | + | + | + | + | + | |
| protein) (PRF1) peroxisomal acyl-CoA thioesterase (PTE1) Peroxisomal acyl- coenzyme A oxidase peroxisomal farnesylated protein (PXF) phorbol-12-myristate-13- acetate-induced protein (PMAIP1) phosphate carrier (mitochondrial gene?) Phosphate carrier, mitochondrial (PHC) phosphate cytidylyltransferase 1, choline, alpha isoform (PCYT1A) PHOSPHATIDATE CYTIDYLYLTRANSFERAS | 14 2 1 1 1 1 3 | X86032 X71440 X75535 D90070 X77337 X60036 | + | + | + | + | + | + | |
| protein) (PRF1) peroxisomal acyl-CoA thioesterase (PTE1) Peroxisomal acyl- coenzyme A oxidase peroxisomal famesylated protein (PXF) phorbol-12-myristate-13- acetate-induced protein (PMAIP1) phosphate carrier (mitochondrial gene?) Phosphate carrier, mitochondrial (PHC) phosphate carrier (cytidylytransferase 1, choline, alpha isoform (PCYT1A) PHOSPHATIDATE CYTIDYLYLTRANSFERAS E (CDP-DIGLYCERIDE) | 14 2 1 1 1 3 1 | X86032 X71440 X75535 D90070 X77337 X60036 L28957 | + | + | + | + | + | + | |
| protein) (PRF1) peroxisomal acyl-CoA thioesterase (PTE1) Peroxisomal acyl- coenzyme A oxidase peroxisomal famesylated protein (PXF) phorbot-12-myristate-13- acetate-induced protein (PMAIP1) phosphate carrier (mitochondrial gene?) Phosphate camer, mitochondrial (PHC) phosphate camer, cytidylytransferase 1, choline, alpha isoform (PCYT1A) PHOSPHATIDATE CYTIDYLYLTRANSFERAS E (CDP-DIGLYCERIDE) phosphatidylinositol 3- | 14 2 1 1 1 3 | X86032 X71440 X75535 D90070 X77337 X60036 L28957 | + | + | + | + | + | + | |
| protein) (PRF1) peroxisomal acyl-CoA thioesterase (PTE1) Peroxisomal acyl- coenzyme A oxidase peroxisomal famesylated protein (PXF) phorbol-12-myristate-13- acetate-induced protein (PMAIP1) phosphate carrier (mitochondrial gene?) Phosphate carrier, mitochondrial (PHC) phosphate cytidylyltransferase 1, choline, alpha isoform (PCYT1A) PHOSPHATIDATE CYTIDYLYLTRANSFERAS E (CDP-DIGLYCERIDE) phosphatidylinositol 3- kinase delta catalytic subunit | 14 2 1 1 1 3 1 | X86032 X71440 X75535 D90070 X77337 X60036 L28957 | + | + | + | + | + | + | |
| protein) (PRF1) peroxisomal acyl-CoA thioesterase (PTE1) Peroxisomal acyl- coenzyme A oxidase peroxisomal farnesylated protein (PXF) phorbol-12-myristate-13- acetate-induced protein (PMAIP1) phosphate carrier (mitochondrial gene?) Phosphate carrier, mitochondrial (PHC) phosphate cytidylyltransferase 1, choline, alpha isoform (PCYT1A) PHOSPHATIDATE CYTIDYLYLTRANSFERAS E (CDP-DIGLYCERIDE) phosphatidylinositol 3- kinase delta catalytic subunit phosphatidylinositol 4- | 14 2 1 1 1 3 1 | X86032 X71440 X75535 D90070 X77337 X60036 L28957 | + | + | + | + | + | + | |
| protein) (PRF1) peroxisomal acyl-CoA thioesterase (PTE1) Peroxisomal acyl- coenzyme A oxidase peroxisomal famesylated protein (PXF) phorbol-12-myristate-13- acetate-induced protein (PMAIP1) phosphate carrier (mitochondrial gene?) Phosphate carrier, mitochondrial (PHC) phosphate carrier cytidylyltransferase 1, choline, alpha isoform (PCYT1A) PHOSPHATIDATE CYTIDYLYLTRANSFERAS E (CDP-DIGLYCERIDE) phosphatidylinositol 3- kinase delta catalytic subunit phosphatidylinositol 4- kinase, catalytic, beta | 14 2 1 1 1 3 1 | X86032 X71440 X75535 D90070 X77337 X60036 L28957 Q92903 U57843 | + | + | + + + | + | + | + | |
| protein) (PRF1) peroxisomal acyl-CoA thioesterase (PTE1) Peroxisomal acyl- coenzyme A oxidase peroxisomal famesylated protein (PXF) phorbol-12-myristate-13- acetate-induced protein (PMAIP1) phosphate carrier (mitochondrial gene?) Phosphate carrier, mitochondrial (PHC) phosphate carrier, choline, alpha isoform (PCYT1A) PHOSPHATIDATE CYTIDYLYLTRANSFERAS E (CDP-DIGLYCERIDE) phosphatidylinositol 3- kinase delta catalytic subunit phosphatidylinositol 4- kinase, catalytic, beta polypeptide (PIK4CB) | 14 2 1 1 1 3 1 1 2 3 3 | X86032 X71440 X75535 D90070 X77337 X60036 L28957 Q92903 U57843 AB005910 | + | + | + + + | + | + | + | |
| protein) (PRF1) peroxisomal acyl-CoA thioesterase (PTE1) Peroxisomal acyl- coenzyme A oxidase peroxisomal farnesylated protein (PXF) phorbol-12-myristate-13- acetate-induced protein (PMAIP1) phosphate carrier (mitochondrial gene?) Phosphate carrier, mitochondrial (PHC) phosphate cytidylyltransferase 1, choline, alpha isoform (PCYT1A) PHOSPHATIDATE CYTIDYLYLTRANSFERAS E (CDP-DIGLYCERIDE) phosphatidylinositol 3- kinase delta catalytic subunit phosphatidylinositol 4- kinase, catalytic, beta | 14 2 1 1 1 3 1 | X86032 X71440 X75535 D90070 X77337 X60036 L28957 Q92903 U57843 | + | + | + + + | + | + | + | |

| W & 00/40/49 | | | | | | | | | CI/CF | KUU/U | COU | |
|---|-------------|-------------|----------|--------------|------|----------------|--|----------|--------------|-------|-----|----------|
| phosphatidylinositol transfer protein (PI-TPbeta) | 2 | D30037 | | | | T | | | | | | ., |
| phosphatidylinositol | 2 | X98654 | В, Т | + | | 1 | | 1 | | | | |
| transfer protein, membrane-associated | · | | lymphoma | 1 | | | | ! | | | | |
| (PITPNM) | | <u>L</u> | 1 | | | | | | | | | |
| phosphatidylinositol transfer protein. | 1 | X98654 | | | | | | • | | | | |
| membrane-associated | | | l | | l | | 1 | Ì | ĺ | | | |
| (PITPNM) (non-exact 64%) | | | 1 | <u> </u> | _ | l | | | | | | |
| phosphatidylinositol-4- phosphate 5-kinase, type | 1 | U14957 | | | + | | + | | | | | |
| II, alpha (PIP5K2A) | | <u> </u> | 1 | ļ | | | 1 | 1 | | | | |
| phosphatidylinositol-4- | 1 | U85245 | | + | + | + | | + | | | | |
| phosphate 5-kinase, type II, beta (PIP5K2B) | | | 1 | | | | | | | | | |
| phosphodiesterase 7A | 1 | L12052 | B, W | + | + | - | + | | | | | |
| (PDE7A) phosphodiesterase IB | | U56976 | <u> </u> | | LY | ـــــ | <u> </u> | | ļ | | | |
| (PDES1B) | | 030970 | | 0 | IL I | | ŀ | | | | | |
| phosphoglucomutase 1 (PGM1) | 2 | M83088 | | + | + | + | | + | | | | |
| phosphogluconate dehydrogenase (PGD) | 1 | U30255 | | | + | | | | | | | |
| phosphoglycerate kinase 1 | 12 | V00572 | <u> </u> | | - | - | ├ | | | | | { |
| (PGK1) | | | | | | | | | | | | |
| phosphoglycerate mutase 1 (brain) (PGAM1) | 3 | J04173 | + | +- | + | + | + | + | • | | | |
| phosphoglycerate mutase 2 (muscle) (PGAM2) | 1 | M55673 | | + | + | | | + | <u> </u> | | | |
| phosphoinositide-3-kinase, | 1 | Z29090 | | + | + | + | ├ | | | | | |
| catalytic, alpha polypeptide (PIK3CA) | | | | | | | | | | | | |
| phosphoinositide-3-kinase, catalytic, delta polypeptide | 4 | U86453 | | + | + | + | | + | | | | |
| (PIK3CD) | | | | | | ł | 1 | | | | | ļ |
| phosphoinositide-3-kinase, catalytic, gamma | 1 | X83368 | | | | | | | | | | |
| polypeptide (PIK3CG) | | | · | | | l | | | | | | i |
| phospholipase C | 1 | X14034 | | _ | | | | \vdash | | | | _ |
| phospholipase C, delta 1 (PLCD1) | 2 | U09117 | | + | + | + | | + | | | | |
| phospholipase C, gamma 1 | 1 | M34667 | + | + | + | + | | + | | | | |
| (formerly subtype 148) (PLCG1) | | | | | | | . | | | | | |
| phospholipid scramblase | 1 | AF008445 | <u> </u> | | | \vdash | | | | | | \dashv |
| phosphoribosyl | 1 . | D61391 | | + | + | - | _ | + | | | | |
| pyrophosphate synthetase- associated protein 1 | | | | | | | | | | | | |
| (PRPSAP1) | | l | | | | | | | | | | |
| phosphoribosylglycinamide formyltransferase. | 3 | X54199 | | + | + | + | + | + | | | | |
| phosphoribosylglycinamide | | |] | | | | | | | | | |
| synthetase, phosphoribosylaminoimida | | | | | | | | | , | | | |
| zole synthetase (GART) | | | ł | | | | | | | | | 1 |
| phosphorylase kinase, | 3 | D38616 | | + | + | + | + | + | | | | \dashv |
| alpha 2 (liver), glycogen storage disease IX | | İ | | | | | | | | | | |
| (PHKA2) | | | | | | | | | | | | - |
| phosphorylase, glycogen; brain (PYGB) | 1 | U47025 | + | + | + | | | + | | | | |
| phosphorylase, glycogen; brain (PYGB) (low match, | 1 | U47025 | | | | | | | | | | |
| non-exact, 75%) phosphorylase, glycogen; | | Y15233 | | + | + | + | | + | | | | |
| liver (Hers disease, | ' | 110200 |] | 🔻 | 7 | • | | 7 | | | | |
| lycogen storage disease type VI) (PYGL) | | 1 | 1 | | | | | | | | | |
| phosphorylation regulatory | 2 | | | | | | | | | | | |
| protein HP-10 phosphotidylinositol | | Danas | | | | | | | | | | |
| transfer protein (PITPN) | 1 | D30036 | + | + | + | + | | + | | | | |
| | | | | | | | | | _ | | | |

| | | | | | | | | • • | JI/CAU | ,, 0000. | • |
|--|-----|--------|-------------|--|--------------|--|----------|----------|----------|----------------|-----|
| pigment epithelium-denved factor (PEDF) | 7 | U29953 | Ŧ | + | + | + | + | TŦ | | | .,- |
| pim-1 oncogene (PIM1) | 1 | M24779 | + | + | + | | | + | <u> </u> | | |
| pinin, desmosome associated protein (PNN) | 1 | U77718 | | В | , mor | nocyt | ė, T | lymp | homa | | |
| placenta (Diff33) | 5 | U49188 | | + | + | + | | T + | | | |
| placenta (Diff33) (non- exact, 69%) | 1 | U49188 | | | | | | | | | |
| placenta (Diff48) | 18 | U49187 | + | +- | | ┼─ | + | +- | - | | |
| placenta (Diff48) (low match) | 1 | U49187 | | 1 | | | \vdash | | | | |
| placenta(Diff48) (low match) | 1 | U49187 | | 1 | | | | | - | | |
| plasminogen activator, urokinase receptor (PLAUR) | 1 | X74039 | | + | | + | | + | | | |
| platelet factor 4 (PF4) | 1 | M25897 | | + | + | + | - | + | | | |
| platelet/endothelial cell | 8 | M37780 | | + | + | + | + | + | | | |
| adhesion molecule (CD31 ntigen) (PECAM1) | | | | | ļ | | : | | | | |
| platelet-activating factor acetylhydrolase 2 (40kD) (PAFAH2) | 4 | U89386 | | + | + | + | | | | | |
| platelet-activating factor acetylhydrolase, isoform lb, alpha subunit (45kD) (PAFAH1B1) | 1 | U72342 | + | + | + | + | + | + | | | |
| platelet-activating factor | 1 | D10202 | | + | | | _ | + | | · | |
| receptor (PTAFR) pleckstrin (PLEK) | 10 | X07743 | | _ | + | + | _ | <u> </u> | | | |
| pleckstrin (PLEK) (low | 1 | X07743 | | | | <u> </u> | | + | | | |
| match) | | | | | | | | | | | |
| pleckstrin homology, Sec7 and coiled/coil domains 1(cytohesin 1) (PSCD1) | 4 | M85169 | + | + | | + | | + | | . . | |
| pleckstrin homology, Sec7 and coiled/coil domains, binding protein (PSCDBP) | 4 | L06633 | + | | | + | | | | • | · : |
| pM5 protein | 1 | X57398 | + | + | + | + | | + | | | |
| PMP69 | 2 | Y14322 | | | | 一 | _ | | | | |
| poly (ADP-ribose) polymerase (NAD (+) ADP- ribosyltransferase) (=X16674) | 1 | X56140 | , | | | | | | | | |
| poly(A) polymerase (PAP) | 1 | X76770 | + | + | + | + | | + | | | |
| poly(A)-binding protein-like 1 (PABPL1) | 19 | Y00345 | + | + | + | + | + | + | | | |
| poly(rC)-binding protein 1 (PCBP1) | 3 | X78137 | + | + | + | + | + | + | | | |
| polyadenylate binding protein | 1 - | U75686 | | | | | | | | | |
| polycystic kidney disease 1 (autosomal dominant) (PKD1) | 5 | U24498 | | | | | | | | | |
| polymerase (DNA directed), beta (POLB) | 1 | D29013 | | + | | | + | + | | | |
| polymerase (DNA directed), gamma (POLG) | 6 | D84103 | | | | | | | | | |
| polymerase (RNA) II (DNA directed) polypeptide A (220kD) (POLR2A) | 1 | X63564 | + | + | + | + | + | + | | | |
| polymyositis/scleroderma autoantigen 2 (100kD) (PMSCL2) | 1 | L01457 | + | + | + | + | + | + | | | |
| polypyrimidine tract binding protein (heterogeneous nuclear ribonucleoprotein I) (PTB) | r - | X65372 | + | + | + | + | + | + | | | |

PCT/CA00/00005

WO 00/40749

| positive regulator of programmed cell death ICH-1L (Ich-1) | 3 | U13021 | | | + | | | | |
|---|--------------|----------|--------|--|---|---|---|--|------------------|
| postmeiotic segregation increased 2-like 12 (PMS2L12) | 1 | M16514 | + | + | + | + | | + | |
| postmeiotic segregation increased 2-like 8 (PMS2L8) | 1 | U38964 | + | + | + | + | | + | |
| potassium inwardly- rectifying channel, subfamily J, member 15 (KCNJ15) | 1 | D87291 | | | | + | | + | |
| potassium voltage-gated channel, KQT-like subfamily, member 1 (KCNQ1) | 1 | AF051426 | | + | + | + | | + | |
| POU domain, class 2, associating factor 1 (POU2AF1) | 1 | Z49194 | | | | + | | | |
| POU domain, class 2, transcription factor 1 (POU2F1) | 2 | X13403 | | + | | + | | | |
| PPAR binding protein (PPARBP) | 1 | Y13467 | + | + | + | + | | + | |
| PPAR gamma2 | | D83233 | | 1 | | 1 | | <u> </u> | <u> </u> |
| pre-B-cell colony- enhancing factor (PBEF) | 8 | U02020 | | | | | | | |
| prefoldin 1 (PFDN1) | 1 | Y17392 | + | + | + | + | + | + | |
| prefoldin 5 (PRFLD5) | 3 | D89667 | В | + | + | 1 | + | | |
| prefoldin subunit 3 (=U96759 von Hippel- Lindau binding protein (VBP-1)) | 1 | Y17394 | | | | | | | |
| pregnancy-associated plasma protein A (PAPPA) | 1 | U28727 | | + | | + | | | high in placenta |
| pre-mRNA splicing factor SF3a (60kD), similar to S. cerevisiae PRP9 (spliceosome-associated protein 61) (SF3A60) | 1 | U08815 | + | + | + | + | | + | |
| pre-mRNA splicing factor SF3a (60kD), similar to S. cerevisiae PRP9 (spliceosome-associated protein 61) (SF3A60) (low score) | 1 | U08815 | | | | | | | |
| pre-mRNA splicing factor SRp20, 5'UTR | 2 | D28423 | | | | Г | | | |
| preprotein translocase (TIM17) | 3 | X97544 | + | + | + | + | | + | |
| prion protein | 1 | X82545 | | | | | | | |
| prion protein (p27-30) (Creutzfeld-Jakob disease, Gerstmann-Strausler- Scheinker syndrome, fatal familial insomnia) (PRNP) | 1 | M13899 | | + | + | + | | + | |
| pristanoyl-CoA oxidase (low match) | - 1 | Y11411 | | | | | | | |
| pristanoyl-CoA oxidase (low score) | 1 | Y11411 | ****** | | | | | | |
| procollagen-lysine, 2- oxoglutarate 5- dioxygenase (lysine hydroxylase, Ehlers-Danlos syndrome type VI) (PLOD) | 1 | M98252 | | + | + | + | | + | · |
| procollagen-proline, 2- oxoglutarate 4- dioxygenase (proline 4- hydroxylase), alpha polypeptide 1 (P4HA1) | | M24486 | + | + | + | + | + | + | |

| W O 00/40/49 | | | | | | | | 10 | 1/CA00/00003 | |
|---|----|--------|---|---------------|---|---|---|----------|--------------|----------|
| procollagen-proline, 2- oxoglutarate 4- dioxygenase (proline 4- hydroxylase), beta polypeptide (protein disulfide isomerase; thyroid hormone binding protein p55) (P4HB) | | X05130 | + | + | + | + | + | + | · | |
| profilin 1 (PFN1) | 1 | J03191 | + | + | + | + | + | + | | I |
| progesterone receptor- associated p48 protein (P48) | 2 | U28918 | | + | | | | | | |
| prohibitin (PHB) | 1 | S85655 | | + | + | + | + | + | | \neg |
| proliferating cell nuclear antigen (PCNA) | 3 | J04718 | + | + | + | + | | + | | |
| proliferation-associated gene A (natural iller- enhancing factor A) (PAGA) | 4 | L19184 | + | + | + | + | + | + | | |
| proline-rich protein BstNI subfamily 2 (PRB2) (non- exact, 43%aa) | 1 | S62936 | | | | | | | | |
| proline-serine-threonine phosphatase interacting protein 1 (PSTPIP1) | 1 | U94778 | | | | | | | | |
| prolyl endopeptidase (PREP) | 2 | X74496 | | + | | + | | + | - | \Box |
| prolylcarboxypeptidase (angiotensinase C) (PRCP) | 5 | L13977 | | + | + | + | + | + | | ٦ |
| promyelocytic leukemia (PML) | 1 | M80185 | + | + | + | + | | + | | \neg |
| properdin P factor. | 4 | X57748 | + | † | | | | | | ᅱ |
| complement (PFC) pro-platelet basic protein | | M54995 | | - | + | + | | + | | \dashv |
| (includes platelet basic protein, beta-thromboglobulin, connective tissue-activating peptide III, neutrophil-activating peptide-2) (PPBP) | | | | | | | | | | |
| pro-platelet basic protein (includes platelet basic protein, beta- thromboglobulin, connective tissue- activating peptide III, neutrophil-activating peptide-2) (PPBP) | 7 | M54995 | + | | + | | + | | | |
| proprotein convertase subtilisin/kexin type 7 (PCSK7) | 4 | U40623 | | | | | | | | |
| prosaposin (variant Gaucher disease and variant metachromatic leukodystrophy) (PSAP) | 89 | D00422 | + | + | + | + | + | + | | |
| prostaglandin- endoperoxide synthase 1 (prostaglandin G/H synthase and cyclooxygenase) (PTGS1) | 1 | U63846 | В | + | | | + | + | | |
| prostaglandin- endoperoxide synthase 2 (prostaglandin G/H synthase and cyclooxygenase) (PTGS2) | 2 | L15326 | | | | | | | | |
| prostagiandin- endoperoxide synthase-1 (=L08404; U84208) (all promoters) | 7 | D64068 | | | | | | | | |
| prostate carcinoma tumor | 2 | L78132 | | | | | | \vdash | | \dashv |
| antigen (pcta-1) | | | L | | | | | | |] |

| M93056 | WO 00/40749 | | | | | | | | PC | T/CA00/00005 |
|--|---|----|----------|---|-------|-----|-----|-----|-----|------------------------|
| Processors Institutor Processors Pro | | 17 | KU5515 | | 1 + 1 | + 1 | + 1 | + 1 | + [| high in many libraries |
| professe infibitor 2 (anti- leastase), monocyte/neutrophil (ELANF2) (low match) professome (prosome, macropain) 285 subunit, ATPase, 3 (PSMC3) professome (prosome, macropain) 285 subunit, ATPase, 3 (PSMC4) professome (prosome, macropain) 285 subunit, ATPase, 3 (PSMC4) professome (prosome, macropain) 285 subunit, ATPase, 3 (PSMC5) professome (prosome, macropain) 285 subunit, ATPase, 3 (PSMC5) professome (prosome, macropain) 285 subunit, ATPase, 4 (PSMC5) professome (prosome, macropain) 285 subunit, non-ATPase, 1 (PSMD11) professome (prosome, macropain) 285 subunit, non-ATPase, 2 (PSMD5) professome (prosome, macropain) 285 subunit, non-ATPase, 5 (PSMD5) professome (prosome, macropain) 285 subunit, non-ATPase, 5 (PSMD5) professome (prosome, macropain) 285 subunit, non-ATPase, 5 (PSMD5) professome (prosome, macropain) 285 subunit, non-ATPase, 5 (PSMD5) professome (prosome, macropain) 285 subunit, non-ATPase, 5 (PSMD5) professome (prosome, macropain) 285 subunit, non-ATPase, 7 (Mov34 homolog) (PMSD7) professome (prosome, macropain) 385 subunit, non-ATPase, 7 (Mov34 homolog) (PMSD7) professome (prosome, macropain) 380 subunit, alpha type, 3 (PSMA7) professome (prosome, macropain) subunit, alpha type, 3 (PSMA7) (low match) type, 7 (PSMA7) (low match) professome (prosome, macropain) subunit, beta type, 6 (PSMB0) professome (prosome, macropain) subunit, beta type, 8 (PSMB0) professome (prosome, macropain) subunit, beta type, 9 (PSMB0) professome (prosome, macropain) subunit, beta type, 9 (PSMB0) professome (prosome, macropain) subunit, beta type, 9 (PSMB0) professome (prosome, macropain) subunit, beta type, 9 (PSMB0) | eiastase), alpha-1- | 17 | NOZZ1Z | | | | | | | |
| (ELANI-12) (low match) professore (prosome, macropain) 28S subunit, ATPase, 1 (PSMC1) professore (prosome, macropain) 28S subunit, ATPase, 1 (PSMC4) professore (prosome, macropain) 28S subunit, ATPase, 2 (PSMC4) professore (prosome, macropain) 28S subunit, ATPase, 4 (PSMC4) professore (prosome, macropain) 28S subunit, ATPase, 4 (PSMC4) professore (prosome, macropain) 28S subunit, ATPase, 6 (PMSC6) professore (prosome, macropain) 28S subunit, ATPase, 6 (PMSC6) professore (prosome, macropain) 28S subunit, ATPase, 6 (PMSC6) professore (prosome, macropain) 28S subunit, atPase, 6 (PMSC6) professore (prosome, macropain) 28S subunit, non-ATPase, 2 (PSMD2) professore (prosome, macropain) 28S subunit, non-ATPase, 2 (PSMD2) professore (prosome, macropain) 28S subunit, non-ATPase, 6 (PSMD6) professore (prosome, macropain) 28S subunit, non-ATPase, 1 (PSMD12) professore (prosome, macropain) 28S subunit, non-ATPase, 1 (PSMD12) professore (prosome, macropain) 28S subunit, non-ATPase, 1 (PSMD12) professore (prosome, macropain) advistor subunit (PSME1) professore (prosome, macropain) subunit, siphs (pps, 3 (PSMD6)) professore (prosome, macropain) subunit, siphs (pps, 3 (PSMA)) professore (prosome, macropain) subunit, siphs (pps, 4 (PSMA)) professore (prosome, macropain) subunit, beta (pps, 4 (PSMA)) to macropain) subunit, beta (pps, 6 (PSMB1)) professore (prosome, macropain) subunit, beta (pps, 6 (PSMB1)) professore (prosome, macropain) subunit, beta (pps, 6 (PSMB1)) professore (prosome, macropain) subunit, beta (pps, 6 (PSMB1)) professore (prosome, macropain) subunit, beta (pps, 6 (PSMB1)) professore (prosome, macropain) subunit, beta (pps, 6 (PSMB1)) professore (prosome, macropain) subunit, beta (pps, 6 (PSMB1)) professore (prosome, macropain) subunit, beta (pps, 6 (PSMB1)) professore (prosome, macropain) subunit, beta (pps, 6 (PSMB1)) professore (prosome, macropain) subunit, beta (pps, 6 (PSMB1)) professore (prosome, macropain) subunit, beta (pps, 6 (PSMB1)) | protease inhibitor 2 (anti- | 1 | M93056 | | | | + | | + | |
| Macropain) 28S subunit, ATPase, (PSMC1) 1 M34079 + <td>(ELANH2) (low match)</td> <td>3</td> <td>L02426</td> <td>В</td> <td>+</td> <td>+</td> <td>-</td> <td></td> <td>+</td> <td></td> | (ELANH2) (low match) | 3 | L02426 | В | + | + | - | | + | |
| AFO01212 AFOO1212 | macropain) 26S subunit, ATPase, 1 (PSMC1) | | | | | | | | | |
| ### ### ### ### ### ### ### ### ### ## | macropain) 26S subunit, | 1 | M34079 | + | + | * | | | • | |
| professome (prosome, macropain) 268 subunit, ATPase, 5 (PSMC5) professome (prosome, macropain) 268 subunit, ATPase, 6 (PMSC6) professome (prosome, macropain) 268 subunit, non-ATPase, 11 (PSMD11) professome (prosome, macropain) 268 subunit, non-ATPase, 2 (PSMD2) professome (prosome, macropain) 268 subunit, non-ATPase, 2 (PSMD5) professome (prosome, macropain) 268 subunit, non-ATPase, 7 (Mov34 homolog) (PMSD7) professome (prosome, macropain) 268 subunit, non-ATPase, 7 (Mov34 homolog) (PMSD7) professome (prosome, macropain) 268 subunit, non-ATPase, 12 (PMSD12) professome (prosome, macropain) 268 subunit, non-ATPase, 12 (PMSD12) professome (prosome, macropain) 268 subunit, non-ATPase, 12 (PMSD12) professome (prosome, macropain) subunit, alpha type, 3 (PSMA3) professome (prosome, macropain) subunit, alpha type, 3 (PSMA5) professome (prosome, macropain) subunit, alpha type, 5 (PSMS) professome (prosome, macropain) subunit, alpha type, 7 (PSMA7) (low match) professome (prosome, macropain) subunit, beta type, 7 (PSMB10) professome (prosome, macropain) subunit, beta type, 7 (PSMB10) professome (prosome, macropain) subunit, beta type, 6 (Ramba) subunit, beta type, 6 (Ramba) subunit, beta type, 6 (Ramba) subunit, beta type, 6 (Ramba) subunit, beta type, 6 (Ramba) subunit, beta type, 6 (Ramba) subunit, beta type, 6 (Ramba) subunit, beta type, 6 (Ramba) subunit, beta type, 6 (Ramba) subunit, beta type, 6 (Ramba) subunit, beta type, 6 (Ramba) subunit, beta type, 6 (Ramba) subunit, beta type, 6 (Ramba) subunit, beta type, 8 (Ramba) subunit, beta type, 8 (Ramba) subunit, beta type, 8 (Ramba) subunit, beta type, 8 (Ramba) subunit, beta type, 8 (Ramba) subunit, beta type, 8 (Ramba) subunit, beta type, 8 (Ramba) subunit, beta type, 8 (Ramba) subunit, beta type, 8 (Ramba) subunit, beta type, 8 (Ramba) subunit, beta type, 8 (Ramba) subunit, beta type, 9 (Ramba) subunit, beta type, 8 (Ramba) subunit, beta type, 9 (Ramba) subunit, beta type, 8 (Ramba) subunit, beta type, 8 (Ramba) subunit, beta type, 8 (Ramba) subunit, beta type, 8 (| macropain) 26S subunit. | 2 | AF020736 | | | | | | | |
| D76275 | proteasome (prosome, macropain) 26S subunit. | 5 | L38810 | + | + | + | + | + | + | |
| AF001212 | proteasome (prosome, macropain) 26S subunit, | 2 | D78275 | + | + | + | + | | + | |
| D78151 | proteasome (prosome, macropain) 26S subunit, | 1 | AF001212 | | + | | | + | | |
| proteasome (prosome, macropain) 26S subunit, non-ATPase, 5 (PSMD5) proteasome (prosome, macropain) 26S subunit, non-ATPase, 7 (Mov34 homolog) (PMSD7) proteasome (prosome, macropain) 36S subunit, on-ATPase, 7 (Mov34 homolog) (PMSD7) proteasome (prosome, macropain) activator subunit 1 (PA28 alpha) (PSME1) proteasome (prosome, macropain) subunit, alpha type, 3 (PSMA3) proteasome (prosome, macropain) subunit, alpha type, 7 (PSMA7) (pow match) proteasome (prosome, macropain) subunit, alpha type, 7 (PSMA7) (pow match) proteasome (prosome, macropain) subunit, alpha type, 7 (PSMA7) (low match) proteasome (prosome, macropain) subunit, alpha type, 7 (PSMA7) (low match) proteasome (prosome, macropain) subunit, beta type, 1 (PSMB10) proteasome (prosome, macropain) subunit, beta type, 6 (PSMB10) proteasome (prosome, macropain) subunit, beta type, 6 (PMSB6) proteasome (prosome, macropain) subunit, beta type, 6 (PMSB6) proteasome (prosome, macropain) subunit, beta type, 6 (PMSB6) proteasome (prosome, macropain) subunit, beta type, 6 (PMSB6) proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional protease 7) (PSMB8) proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional protease 7) (PSMB8) proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional protease 7) (PSMB8) proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional protease 7) (PSMB9) proteasome (prosome, macropain) subunit, beta type, 9 (large multifunctional protease 7) (PSMB9) proteasome (prosome, macropain) subunit, beta type, 9 (large multifunctional protease 7) (PSMB9) proteasome (prosome, macropain) subunit, beta type, 9 (large multifunctional protease 7) (PSMB9) proteasome (prosome, macropain) subunit, beta type, 9 (large multifunctional protease 7) (PSMB9) proteasome (prosome, macropain) subunit, beta type, 9 (large multifunctional protease 7) (PSMB9) proteasome (prosome, macropain) subunit, beta type, 9 (large multifunctional protease 7) (PSMB9) proteasome (prosome, macropain) su | proteasome (prosome, macropain) 26S subunit, | 2 | D78151 | | | | | | + | |
| profeasome (prosome, macropain) 26S subunit, non-ATPase, 7 (Mov34 homolog) (PMSDT) profeasome (prosome, macropain) 26S subunit, on-ATPase, 8, 12 (PMSD12) profeasome (prosome, macropain) activator subunit 1 (PA28 alpha) ((PSME1) profeasome (prosome, macropain) subunit, alpha type, 3 (PSMA3) profeasome (prosome, macropain) subunit, alpha type, 3 (PSMA5) profeasome (prosome, macropain) subunit, alpha type, 5 (PSMA5) profeasome (prosome, macropain) subunit, alpha type, 7 (PSMA5) profeasome (prosome, macropain) subunit, alpha type, 7 (PSMA7) profeasome (prosome, macropain) subunit, alpha type, 7 (PSMA7) (low match) profeasome (prosome, macropain) subunit, beta type, 10 (PSMB10) profeasome (prosome, macropain) subunit, beta type, 10 (PSMB10) profeasome (prosome, macropain) subunit, beta type, 8 (PSMSB) profeasome (prosome, macropain) subunit, beta type, 8 (PSMSB) profeasome (prosome, macropain) subunit, beta type, 8 (PSMSB) profeasome (prosome, macropain) subunit, beta type, 8 (Brasome, profeasome, profeasome (prosome, macropain) subunit, beta type, 8 (Brasome, profeasome, profeasome (prosome, macropain) subunit, beta type, 8 (Brasome, profeasome, profeasome (prosome, macropain) subunit, beta type, 8 (Brasome, profeasome, profeasome (prosome, macropain) subunit, beta type, 8 (Brasome, profeasome, profeasome (prosome, macropain) subunit, beta type, 8 (Brasome, profeasome, profeasome, profeasome (prosome, macropain) subunit, beta type, 9 (Brasome, profea | proteasome (prosome, macropain) 26S subunit, | 1 | 579862 | T | + | + | | + | | |
| proteasome (prosome, macropain) 26S subunit, on-ATPase, 12 (PMSD12) proteasome (prosome, macropain) activator subunit 1 (PA28 alpha) (PSME1) proteasome (prosome, macropain) subunit, alpha type, 3 (PSMA3) proteasome (prosome, macropain) subunit, alpha type, 3 (PSMA5) proteasome (prosome, macropain) subunit, alpha type, 5 (PSMA5) proteasome (prosome, macropain) subunit, alpha type, 7 (PSMA7) proteasome (prosome, macropain) subunit, alpha type, 7 (PSMA7) flow match) proteasome (prosome, macropain) subunit, alpha type, 7 (PSMB1) proteasome (prosome, macropain) subunit, beta type, 10 (PSMB1) proteasome (prosome, macropain) subunit, beta type, 10 (PSMB1) proteasome (prosome, macropain) subunit, beta type, 10 (PSMB10) proteasome (prosome, macropain) subunit, beta type, 10 (PSMB10) proteasome (prosome, macropain) subunit, beta type, 6 (PMSB6) proteasome (prosome, macropain) subunit, beta type, 6 (PMSB6) proteasome (prosome, macropain) subunit, beta type, 6 (PMSB6) proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional protease 7) (PSMB8) proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional protease 7) (PSMB8) proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional protease 7) (PSMB8) proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional protease 7) (PSMB8) proteasome (prosome, macropain) subunit, beta type, 9 (large | proteasome (prosome, macropain) 26S subunit, non-ATPase, 7 (Mov34 | 1 | D50063 | | + | + | + | | + | high in many libraries |
| proteasome (prosome, macropain) activator subunit 1 (PA28 alpha) (PSME1) proteasome (prosome, macropain) subunit, alpha type, 3 (PSMA3) proteasome (prosome, macropain) subunit, alpha type, 5 (PSMA5) proteasome (prosome, macropain) subunit, alpha type, 5 (PSMA5) proteasome (prosome, macropain) subunit, alpha type, 7 (PSMA7) proteasome (prosome, macropain) subunit, alpha type, 7 (PSMA7) (low match) proteasome (prosome, macropain) subunit, beta type, 1 (PSMB1) proteasome (prosome, macropain) subunit, beta type, 1 (PSMB10) proteasome (prosome, macropain) subunit, beta type, 10 (PSMB10) proteasome (prosome, macropain) subunit, beta type, 6 (PMSB6) proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional protease 7) (PSMB8) proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional protease 7) (PSMB8) proteasome (prosome, macropain) subunit, beta type, 9 (large | proteasome (prosome, macropain) 26S subunit, | 1 | AB003103 | | + | + | + | | + | |
| proteasome (prosome, macropain) subunit, alpha type, 3 (PSMA3) proteasome (prosome, macropain) subunit, alpha type, 5 (PSMA5) proteasome (prosome, macropain) subunit, alpha type, 7 (PSMA7) proteasome (prosome, macropain) subunit, alpha type, 7 (PSMA7) proteasome (prosome, macropain) subunit, alpha type, 7 (PSMA7) (low match) proteasome (prosome, macropain) subunit, beta type, 1 (PSMB1) proteasome (prosome, macropain) subunit, beta type, 10 (PSMB10) proteasome (prosome, macropain) subunit, beta type, 6 (PMSB6) proteasome (prosome, macropain) subunit, beta type, 6 (PMSB6) proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional protease 7) ((PSMB8) proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional protease 7) ((PSMB8) proteasome (prosome, macropain) subunit, beta type, 9 (large | proteasome (prosome, macropain) activator subunit 1 (PA28 alpha) | 3 | L07633 | + | | | | | | |
| proteasome (prosome, macropain) subunit, alpha type, 5 (PSMA5) proteasome (prosome, macropain) subunit, alpha type, 7 (PSMA7) proteasome (prosome, macropain) subunit, alpha type, 7 (PSMA7) (low match) proteasome (prosome, macropain) subunit, beta type, 1 (PSMB1) proteasome (prosome, macropain) subunit, beta type, 10 (PSMB10) proteasome (prosome, macropain) subunit, beta type, 10 (PSMB10) proteasome (prosome, macropain) subunit, beta type, 6 (PMSB6) proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional protease 7) (PSMB8) proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional protease 7) (PSMB8) proteasome (prosome, macropain) subunit, beta type, 9 (large | proteasome (prosome, macropain) subunit, alpha | 2 | D00762 | | | | | | | |
| proteasome (prosome, macropain) subunit, alpha type, 7 (PSMA7) proteasome (prosome, macropain) subunit, alpha type, 7 (PSMA7) (low match) proteasome (prosome, macropain) subunit, beta type, 1 (PSMB1) proteasome (prosome, macropain) subunit, beta type, 10 (PSMB10) proteasome (prosome, macropain) subunit, beta type, 10 (PSMB10) proteasome (prosome, macropain) subunit, beta type, 6 (PMSB6) proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional protease 7) (PSMB8) proteasome (prosome, macropain) subunit, beta type, 9 (large macropain) subunit, beta type, 9 (large macropain) subunit, beta type, 9 (large | proteasome (prosome, macropain) subunit, alpha | 3 | X61970 | + | + | + | - | | | |
| proteasome (prosome, macropain) subunit, alpha type, 7 (PSMA7) (low match) proteasome (prosome, macropain) subunit, beta type, 1 (PSMB1) proteasome (prosome, macropain) subunit, beta type, 10 (PSMB10) proteasome (prosome, macropain) subunit, beta type, 6 (PMSB6) proteasome (prosome, macropain) subunit, beta type, 6 (PMSB6) proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional protease 7) (PSMB8) proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional protease 7) (PSMB8) proteasome (prosome, macropain) subunit, beta type, 9 (large | proteasome (prosome, macropain) subunit, alpha | 3 | AF054185 | | + | + | + | + | + | |
| proteasome (prosome, macropain) subunit, beta type, 1 (PSMB1) proteasome (prosome, macropain) subunit, beta type, 10 (PSMB10) proteasome (prosome, macropain) subunit, beta type, 6 (PMSB6) proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional protease 7) (PSMB8) proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional protease 7) (PSMB8) proteasome (prosome, macropain) subunit, beta type, 9 (large | proteasome (prosome, macropain) subunit, alpha type, 7 (PSMA7) (low | 1 | AF022815 | | | | | | | |
| proteasome (prosome, macropain) subunit, beta type, 10 (PSMB10) proteasome (prosome, macropain) subunit, beta type, 6 (PMSB6) proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional protease 7) (PSMB8) proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional protease 7) (PSMB8) proteasome (prosome, macropain) subunit, beta type, 9 (large | proteasome (prosome, macropain) subunit, beta | 1 | D00761 | + | + | + | + | + | + | |
| proteasome (prosome, macropain) subunit, beta type, 6 (PMSB6) proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional protease 7) (PSMB8) proteasome (prosome, macropain) subunit, beta type, 9 (large multifunctional protease 7) | proteasome (prosome, macropain) subunit, beta | 1 | X71874 | + | + | | + | + | | |
| proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional protease 7) (PSMB8) proteasome (prosome, macropain) subunit, beta type, 9 (large | proteasome (prosome, macropain) subunit, beta | 1 | D29012 | | + | + | + | | | |
| proteasome (prosome, 3 Z14977 + + + + + + + + + + + + + + + + + + | proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional protease 7) | 1 | U17497 | + | + | + | + | | | · |
| multifunctional protease 2) (PSMB9) | proteasome (prosome, macropain) subunit, beta type, 9 (large multifunctional protease 2) | 3 | Z14977 | + | | | + | | + | |

| professome (prosome, macropain) subunit, beta ype, 7 (PSMB7) | WO 00/40/49 | | | | | | | | rc | . I/CA00/00003 |
|--|--|----|----------|-----|----------|---|----------|---|----|------------------------|
| Internative protein for belangual colorables September Septe | macropain) subunit, beta | 1 | D38048 | + | + | + | + | + | + | • |
| protein A alternatively spliced form 2 (A-2) protein activator of the interferon-induced protein (RHS (PACT) protein decivator of the interferon-induced protein (RHS (PACT) protein disulfide isomerase-related protein (PS) protein disulfide isomerase-related protein (PS) protein disulfide isomerase-related protein (PS) protein formation (PS) protein formation (PS) protein formation (PS) protein formation (PS) protein formation (PS) protein formation (PS) protein formation (PS) protein formation (PS) protein finase A anchoring protein kinase A anchoring protein kinase A anchoring protein kinase A anchoring protein kinase C substrate (PRKCSH) protein kinase C substrate (PRKCSH) protein kinase C, delta (PRKCSH) protein kinase C, delta (PRKCSH) protein kinase C, that (PRKCSH) protein kinase C, that (PRKCSH) protein kinase C like 1 (PRKCSH) protein kinase C like 1 (PRKCSH) protein kinase C like 1 (PRKCSH) protein kinase C like 1 (PRKCSH) protein kinase C like 1 (PRKCSH) protein kinase C like 1 (PRKCSH) protein kinase C like 1 (PRKCSH) protein kinase, AMP-activated, gamma 1 non-catalytic subunit (PRKAGT) protein kinase, AMP-activated, gamma 1 non-catalytic subunit (PRKAGT) protein kinase, DNA-activated (PRKCH) protein kinase, AMP-activated (PRKCSH) protein kinase, AMP-activated (PRKCSH) protein kinase, NINGer-activated (PRKCH) protein kinase, NINGer-activated (PRKCH) protein kinase, NINGer-activated (PRKCH) protein kinase, NINGer-activated (PRKCH) protein kinase, NINGer-activated (PRKCH) protein kinase, NINGer-activated (PRKCH) protein kinase, NINGer-activated (PRKCH) protein kinase, NINGer-activated (PRKCH) protein kinase, NINGer-activated (PRKCH) protein kinase, NINGer-activated (PRKCH) protein kinase, NINGer-activated (PRKCH) protein kinase (PRKCH) protein kinase (PRKCH) protein | protective protein for beta- galactosidase | 3 | M22960 | + | + | + | + | + | + | |
| AF072860 | protein A alternatively | 1 | U47925 | | + | | | | | |
| protein disulfide isomerase- related protein (PS) protein geranylgeranyltransferase type I, beta subunit (PGGT18) protein homologous to chicken 8 complex protein, guanine nucleotide binding (H12.3) protein kinase A anchoring protein kinase A anchoring protein kinase C substrate 80K-H (PRKCSH) protein kinase C, beta 1 (PRKCSH) protein kinase C, beta 1 (PRKCSH) protein kinase C, beta 1 (PRKCSH) protein kinase C, delta (PRKCSH) protein kinase C, tal (PRKCSH) protein kinase C, tal (PRKCSH) protein kinase C, tal (PRKCSH) protein kinase C, tal (PRKCSH) protein kinase C, tal (PRKCSH) protein kinase C, tal (PRKCSH) protein kinase C, tal (PRKCSH) protein kinase C, tal (PRKCSH) protein kinase C, tal (PRKCSH) protein kinase C, tal (PRKCSH) protein kinase C, tal (PRKCSH) protein kinase C, tal (PRKCSH) protein kinase C, tal (PRKCSH) protein kinase, AMP- activated, gamma 1 non- catalytic subunit (PRKACSI) protein kinase, CAMP- dependent, regulatory, type I, alpha (tissue specific exinguisher 1 (PRKACH) protein kinase, DMA- cativated 1 (MAP kinase 1 poly, p41) (PRKMT) protein kinase, mitogen- activated 1 (MAP kinase 1 poly, p41) (PRKMT) protein kinase 3 (MAP kinase kinase 3) (PRKMI) protein kinase 3 (MAP kinase kinase 3) (PRKMI) protein kinase 3 (MAP kinase kinase 3) (PRKMI) protein phosphatase 1 to the complex protein phosphatase 1 to the complex protein phosphatase 2 to the complex protein phosphatase 2 to the complex protein phosphatase 2 to the complex protein phosphatase 2 to the complex protein phosphatase 2 to the complex protein phosphatase 2 to the complex protein phosphatase 2 to the complex protein phosphatase 2 to the complex protein phosphatase 2 to the complex protein phosphatase 2 to the complex protein phosphatase 2 to the complex protein phosphatase 2 to the complex protein phosphatase 2 to the complex protein phosphatase 2 to the complex protein phosphatase 2 to the complex phosphatase 2 to the complex phosphatase 2 to the complex phosphatase 2 to the complex phosphatase 2 to the complex phosphatase 2 to the | protein activator of the interferon-induced protein | 1 | AF072860 | | + | + | + | | + | high in testis |
| Decision Supering 1 | protein disulfide isomerase- | 2 | D49489 | + | + | + | + | + | + | |
| protein homologous to chicken B complex protein, guanine nucleotide binding (H12.3) protein kinase A anchoring protein protein kinase A anchoring protein protein kinase C substrate 80K-H (PRKCSH) protein kinase C. beta 1 (PRKCSH) protein kinase C. beta 1 (PRKCSH) protein kinase C, delta (PRKCSH) protein kinase C, delta (PRKCSH) protein kinase C, delta (PRKCSH) protein kinase C, cata (PRKCSH) protein kinase C, cata (PRKCSH) protein kinase C, cata (PRKCSH) protein kinase C, delta (PRKCSH) protein kinase C, tata 1 (PRKCSH) protein kinase C, ata 1 (PRKCSH) protein kinase C, ata 1 (PRKCSH) protein kinase C, ata 1 (PRKCSH) protein kinase C, ata 1 (PRKCSH) protein kinase C, ata 1 (PRKCSH) protein kinase, AMP-activated, against a tata 1 (PRKCSH) protein kinase, C, ata 1 (PRKCSH) protein kinase, C, ata 1 (PRKCSH) protein kinase, C, ata 1 (PRKCSH) protein kinase, C, ata 1 (PRKCSH) protein kinase, C, ata 1 (PRKCSH) protein kinase, C, ata 1 (PRKCSH) protein kinase, C, ata 1 (PRKCSH) protein kinase, C, ata 1 (PRKCSH) protein kinase, C, ata 1 (PRKCSH) protein kinase, C, ata 1 (PRKCSH) protein kinase, milogen-activated 6 (extracellular signal-regulated kinase) protein kinase, milogen-activated 6 (extracellular signal-regulated kinase) protein kinase (PRKCSH) protein kinase | protein geranylgeranyltransferase type I, beta subunit | 1 | L25441 | + | + | + | | | | |
| protein kinase C substrate B0K-H (PRKCSH) protein kinase C, beta 1 (PRKCSH) protein kinase C, beta 1 (PRKCSH) protein kinase C, beta 1 (PRKCSH) protein kinase C, delta (PRKCH) protein kinase C, delta (PRKCH) protein kinase C, teta (PRKCH) protein kinase C, teta (PRKCH) protein kinase C, teta (PRKCH) protein kinase C, tike 1 (PRKCH) protein kinase C, tike 1 (PRKCH) protein kinase C, tike 1 (PRKCH) protein kinase C, tike 1 (PRKCH) protein kinase, cAMP- activated, gamma 1 non- catalytic subunit (PRKAG1) protein kinase, cAMP- dependent, regulatory, type I, alpha (tissue specific extinguisher 1) (PRKAR1A) protein kinase, DNA- activated, catalytic polypeptide (PRKCC) protein kinase, mitogen- activated 1 (MAP kinase 1; p40, p41) (PRKM1) protein kinase, mitogen- activated d (extracellular signal-regulated kinase, p37) (PRKM6) protein kinase, mitogen- activated d (extracellular signal-regulated kinase, p37) (PRKM6) protein kinase, mitogen- activated kinase 3 (MAP kinase kinase 3) (MAP kinase kinase 3) (MAP kinase kinase 3) (MAP kinase kinase 3) (PPCA) protein phosphatase 1, regulatory subunit 10 (PPPRTO) protein phosphatase 1, regulatory subunit 7 (PPPNRTO) protein phosphatase 2 1 X12658 + + + + + + + ** ** ** ** ** ** ** ** ** ** ** ** * | chicken B complex protein, guanine nucleotide binding | 20 | M24194 | + | + | + | + | + | + | high in many libraries |
| BOK-H (PRKCSH) protein kinase C, beta 1 (PRKCB1) protein kinase C, delta (PRKCB1) protein kinase C, delta (PRKCB1) protein kinase C, delta (PRKCB1) protein kinase C, eta (PRKCH) protein kinase C, eta (PRKCH) protein kinase C, like 1 (PRKCM) Protein kinase C-like 1 (PRKCB1) protein kinase, AMP- activated, gamma 1 non- catalytic subunit (PRKAG1) protein kinase, AMP- dependent, regulatory, type l, alpha (tissue specific extinguisher 1) (PRKAR1A) protein kinase, DNA- activated, catalytic polypepilide (PRKDC) protein kinase, mitogen- activated 6 (extracellular signal-regulated kinase, p97) (PRKM6) protein kinase, mitogen- activated 6 (extracellular signal-regulated kinase, p97) (PRKM6) protein kinase, mitogen- activated 6 (extracellular signal-regulated kinase) protein kinase, mitogen- activated 6 (extracellular signal-regulated kinase) protein kinase, mitogen- activated kinase 3 (MAR kinase kinase 3) (PRKM6) protein phosphatase 1, regulatory subunit 10 (PPPRCD) protein phosphatase 1, regulatory subunit 7 (PPPRCD) protein phosphatase 2 1 X12656 + + + + + + + ** ** ** ** ** ** ** ** ** ** ** ** * | protein | | | | L | | | | | |
| PRKCB1 | 80K-H (PRKCSH) | _ | | | | | | | | |
| PRKCD Protein kinase C, eta 1 M55284 + + + + + + + + + | (PRKCB1) | | | | | | | | | |
| CPRKCH protein kinase C, mu (PRKCM) (non-exact 78%) | (PRKCD) | | _ | + | + | | _ | | | |
| (PRKCM) (non-exact 78%) Protein kinase C-like 1 | (PRKCH) | | | | <u> </u> | + | | | + | |
| (PRKCL1) protein kinase, AMP- activated, gamma 1 non- catalytic subunit (PRKAG1) protein kinase, cAMP- dependent, regulatory, type I, alpha (tissue specific extinguisher 1) (PRKAR1A) protein kinase, DNA- activated, catalytic polypeptide (PRKDC) protein kinase, mitogen- activated 1 (MAP kinase 1; p40, p41) (PRKM1) protein kinase, mitogen- activated 6 (extracellular signal-regulated kinase, p97) (PRKM6) protein kinase, mitogen- activated kinase 3 (MAP kinase kinase 3) (PRKM6) protein phosphatase 1, catalytic subunit, alpha isoform (PPP1CA) protein phosphatase 1, regulatory subunit 10 (PPPR10) protein phosphatase 1, regulatory subunit 7 (PPP1R7) protein phosphatase 2 (formerly 2A), catalytic subunit, beta isoform (PPP2CB) | (PRKCM) (non-exact 78%) | · | | | | | <u> </u> | | | |
| activated, gamma 1 non-catalytic subunit (PRKAG1) protein kinase, cAMP- dependent, regulatory, type I, alpha (issue specific extinguisher 1) (PRKAR1A) protein kinase, DNA- activated, catalytic polypeptide (PRKDC) protein kinase, miltogen- activated 1 (MAP kinase 1; p40, p41) (PRKM1) protein kinase, miltogen- activated 6 (extracellular signal-regulated kinase, p97) (PRKM6) protein kinase 3 (MAP kinase 3 (M | (PRKCL1) | | | | | | <u> </u> | | _ | |
| dependent, regulatory, type I, alpha (tissue specific extinguisher 1) (PRKAR1A) protein kinase, DNA- activated, catalytic polypeptide (PRKDC) protein kinase, mitogen- activated 1 (MAP kinase 1; p40, p41) (PRKM1) protein kinase, mitogen- activated 6 (extracellular signal-regulated kinase, p97) (PRKM6) protein kinase, mitogen- activated, kinase 3 (MAP kinase kinase, p97) (PRKM6) protein protein phosphatase 1, 5 M63960 + + + + + + + + + + + + + + + + + + + | activated, gamma 1 non- catalytic subunit (PRKAG1) | | | | | + | | | | |
| Drotein kinase, DNA- | dependent, regulatory, type I, alpha (tissue specific | 4 | M18468 | | + | + | + | + | + | |
| Drotein kinase, mitogen-activated 1 (MAP kinase 1; p40, p41) (PRKM1) Protein kinase, mitogen-activated 6 (extracellular signal-regulated kinase, p97) (PRKM6) Protein kinase, mitogen-activated, kinase 3 (MAP kinase kinase 3) (PRKMK3) Protein phosphatase 1, catalytic subunit, alpha isoform (PPP1CA) Protein phosphatase 1, regulatory subunit 10 (PPR10) Protein phosphatase 1, regulatory subunit 7 (PP1R7) Protein phosphatase 2 (formerly 2A), catalytic subunit, beta isoform (PPP2CB) Protein phosphatase 2 (formerly 2A), catalytic subunit, beta isoform (PPP2CB) Protein phosphatase 2 Protein phosphatase 2 Protein phosphatase 3 Protein phosphatase 4 Protein phosphatase 5 Protein phosphatase 6 Protein phosphatase 7 Protein phosphatase 8 Protein phosphatase 9 Prote | protein kinase, DNA- activated, catalytic | 1 | U47077 | | + | + | | * | + | |
| protein kinase, mitogenactivated 6 (extracellular signal-regulated kinase, p97) (PRKM6) protein kinase, mitogenactivated, kinase 3 (MAP kinase kinase 3) ((PRKMK3) protein phosphatase 1, catalytic subunit, alpha isoform (PPP1CA) protein phosphatase 1, regulatory subunit 10 (PPPR10) protein phosphatase 1, regulatory subunit 7 (PPP1R7) protein phosphatase 2 (formerly 2A), catalytic subunit, beta isoform (PPPCCB) | protein kinase, mitogen- activated 1 (MAP kinase 1; | 1 | Z11695 | В | + | | | + | | |
| activated, kinase 3 (MAP kinase kinase 3) (PRKMK3) (PRKMK3) protein phosphatase 1, catalytic subunit, alpha isoform (PPP1CA) protein phosphatase 1, regulatory subunit 10 (PPPR10) protein phosphatase 1, regulatory subunit 7 (PPP1R7) protein phosphatase 2 (formerly 2A), catalytic subunit, beta isoform (PPP2CB) | protein kinase, mitogen- activated 6 (extracellular signal-regulated kinase, | 1 | L77964 | | + | | + | + | + | |
| catalytic subunit, alpha isoform (PPP1CA) protein phosphatase 1, regulatory subunit 10 (PPPR10) protein phosphatase 1, 2 Z50749 + + + + + + + + + + + + + + + + + + + | activated, kinase 3 (MAP kinase kinase 3) (PRKMK3) | | | + " | + | + | + | + | | |
| regulatory subunit 10 (PPPR10) protein phosphatase 1, 2 Z50749 + + + + + + + + + + + + + + + + + + + | catalytic subunit, alpha isoform (PPP1CA) | | | + | + | + | + | + | + | · |
| regulatory subunit 7 (PPP1R7) protein phosphatase 2 1 X12656 + + + + + + + + + + + + + + + + + + | regulatory subunit 10 (PPPR10) | 3 | Y13247 | | + | + | + | | + | |
| (formerly 2A), catalytic subunit, beta isoform (PPP2CB) | protein phosphatase 1, regulatory subunit 7 (PPP1R7) | | | | | | | | | |
| | (formerly 2A), catalytic subunit, beta isoform (PPP2CB) | | | + | + | + | + | | | |
| (formerly 2A), regulatory subunit B" (PR 72), alpha isoform and (PR 130), beta isoform (PPP2R3) | subunit B" (PR 72), alpha isoform and (PR 130), beta | 1 | L07590 | | | + | + | | + | |

80

| WO 00/40/49 | | | | | | | | 1 | .1/CA00/00003 |
|--|----|----------|---|---|---|-----|---|---|----------------|
| protein phosphatase 2, regulatory subunit B (B56), alpha isoform (PPP2R5A) | 2 | L42373 | + | + | + | + | | + | 5 |
| protein phosphatase 2, regulatory subunit B (B56), delta isoform (PPP2R5D) | 3 | D78360 | | + | + | + | | + | |
| protein phosphatase 2, regulatory subunit B (B56), gamma isoform (PPP2R5C) | 1 | D26445 | + | + | + | + | | + | |
| protein phosphatase 2A regulatory subunit alpha- isotype (alpha-PR65) | 5 | J02902 | + | + | + | + | | + | |
| protein phosphatase 4 (formerly X), catalytic subunit (PPP4C) | 2 | AF097996 | + | + | + | + | | + | |
| protein tyrosine kinase 2 beta (PTK2B) | 4 | L49207 | | + | | + | | + | |
| protein tyrosine phosphatase epsilon | 1 | X54134 | | | | | | | |
| protein tyrosine phosphatase type IVA, member 2 (PTP4A2) | 2 | L48723 | + | + | + | + | | + | |
| protein tyrosine phosphatase, non-receptor type 1 (PTPN1) | 1 | M31724 | + | + | + | + | | | |
| protein tyrosine phosphatase, non-receptor type 12 (PTPN12) | 1 | M93425 | | + | + | + | | + | high in testis |
| protein tyrosine phosphatase, non-receptor type 12 (PTPN12) (non- exact, 70%) | 1 | M93425 | | | | | | | · |
| protein tyrosine phosphatase, non-receptor type 2 (PTPN2) | 2 | M25393 | | + | + | + | | + | |
| protein tyrosine phosphatase, non-receptor type 4 (megakaryocyte) (PTPN4) | 1 | M68941 | | | + | + | | + | |
| protein tyrosine phosphatase, non-receptor type 6 (PTPN6) | 7 | M74903 | + | + | + | + | | + | |
| protein tyrosine phosphatase, non-receptor type 7 (PTPN7) | 1 | D11327 | + | | | + | | + | |
| protein tyrosine phosphatase, receptor type, alpha polypeptide (PTPRA) | | M34668 | + | + | + | + | | + | |
| protein tyrosine phosphatase, receptor type, c polypeptide (PTPRC) | 44 | Y00638 | + | + | | + | | + | |
| protein tyrosine phosphatase, receptor type, M (PTPRM) | 1 | X58288 | | + | + | + : | | + | |
| protein tyrosine phosphatase, receptor type, N polypeptide 2 (PTPRN2) | 2 | U81561 | | + | | + | | + | |
| protein with polyglutamine repeat (ERPROT213-21) | 1 | U94836 | + | + | + | + | | + | |
| protein-kinase, interferon- inducible double stranded RNA dependent inhibitor (PRKRI) | 1 | U28424 | | + | + | + | + | + | |
| protein-L-isoaspartate (D- aspartate) O- methyltransferase (PCMT1) | 4 | D13892 | | + | + | | | | |
| proteoglycan 1, secretory granule (PRG1) | 7 | J03223 | | + | | + | | + | |
| prothymosin, alpha (gene sequence 28) (PTMA) | 12 | M14483 | + | + | + | + | + | + | |
| | | | | | | | | | |

| prp28, U5 snRNP 100 kd | 7 | AF026402 | + | + | + | + | Γ | + | , |
|--|----|----------|-------------|----|-------------|--|----|-----|--|
| protein (U5-100K) PRP4/STK/WD splicing | | AF001687 | | ++ | + | + | ├- | + | |
| factor (HPRP4P) | | | | | | <u> </u> | | ļ ' | |
| PTK7 protein tyrosine kinase 7 (PTK7) | 1 | U40271 | | + | + | + | | + | |
| punnergic receptor P2X, ligand-gated ion channel, 4 (P2RX4) | 3 | AF000234 | | + | + | + | | + | |
| punnergic receptor P2X, ligand-gated ion channel, 7 (P2RX7) | 1 | Y12851 | + | | | | | | macrophage only |
| puromycin-sensitive aminopeptidase (PSA) | 1 | Y07701 | | + | + | | | + | |
| putative ATP(GTP)-binding protein | 2 | AJ010842 | | + | | | | + | |
| putative brain nuclearly- targeted protein (KIAA0765) | 1 | AB018308 | + | + | + | + | | + | |
| putative chemokine receptor; GTP-binding protein (HM74) | 1 | D10923 | + | | - | | | | |
| putative dienoyl-CoA isomerase (ECH1) | 1 | AF030249 | | + | · · · · · · | \vdash | _ | | |
| putative G-binding protein | 1 | AF065393 | | + | | | - | - | |
| Putative human HLA class II associated protein ! (PHAP1) | 1 | U73477 | В | + | | | + | | |
| Putative L-type neutral amino acid transporter (KIAA0436) | 1 | AB007896 | , | | - | | | | |
| putative mitochondrial space protein 32.1 | 1 | AF050198 | | | | | | | |
| PUTATIVE MUCIN CORE PROTEIN PRECURSOR 24 (MULTI- | 1 | Q04900 | | | | | | | |
| GLÝCOSYLATED CORE PROTEIN 24) (MGC-24) (MUC-24) | | | | | | | | | · |
| putative nucleic acid binding protein | 2 | X76302 | + | + | + | + | | + | |
| putative outer mitochondrial membrane 34 kDa translocase Htom34 | 1 | U58970 | | + | + | + | | + | |
| putative p150 (non-exact 88%) | 1 | U93568 | | | | | | | |
| putative translation initiation factor (SUI1) | 1 | L26247 | + | + | + | + | + | + | High in moderately differentiated colon adenocarcinoma |
| putative tumor suppressor protein (123F2) | 1 | AF061836 | | + | + | + | | + | |
| pyrroline 5-carboxylate reductase | 1 | M77836 | + | + | + | + | | + | |
| pyruvate dehydrogenase (lipoamide) alpha 1 (PDHA1) | 1 | D90084 | | + | + | + | + | + | |
| pyruvate dehydrogenase (lipoamide) beta (PDHB) | 2 | J03576 | + | + | + | + | | + | |
| Pyruvate dehydrogenase complex, lipoyl-containing component X; E3-binding protein (PDX1) | 3 | Y13145 | | + | + | | | | |
| pyruvate kinase, muscle (PKM2) | 11 | M23725 | | | | | + | | |
| RAB, member of RAS oncogene family-like (RABL) | 1 | U18420 | | + | + | + | | + | |
| RAB1, member RAS oncogene family (RAB1) | 3 | M28209 | | + | + | + | | + | |
| RAB11A, member RAS | 2 | X56740 | + | + | + | + | | + | high in spleen |
| oncogene family (RAB11A) | | | | | | | | | |

| RAB11B, member RAS oncogene family (Rab11B) | 1 | D45418 | | + | | | | + | |
|---|--------------|----------|---|--|---|----------|--|----|---------------|
| RAB27A, member RAS oncogene family (RAB27A) | 3 | U38654 | | + | | + | | + | |
| RAB5B, member RAS | | X54871 | | ++ | + | + | ├ | + | |
| oncogene family (RAB5B) | | | | | | | <u>L</u> | | |
| RAB6, member RAS oncogene family (RAB6) | 1 | M28212 | | + | | | | + | |
| RAB7, member RAS oncogene family (RAB7) | 1 | X93499 | + | + | + | + | | + | |
| RAB7, member RAS oncogene family-like 1 | 2 | D84488 | | + | + | + | | + | |
| (RAB7L1) RAB9, member RAS | 1 | U44103 | | 1 | | <u> </u> | <u> </u> | 1_ | |
| oncogene family (RAB9) RAD50 (S. cerevisiae) | 2 | | | 1 | | <u> </u> | | | |
| homolog (RAD50) | | U63139 | | + | + | + | | | |
| RAD51 (S. cerevisiae) homolog C (RAD51C) | 1 | AF029669 | | + | + | + | | + | |
| Radin blood group (RD) | 2 | L03411 | | + | + | + | | + | |
| RAE1 (RNA export 1, S.pombe) homolog (RAE1) | 3 | U84720 | + | + | + | + | | + | |
| ralA-binding protein (RLIP76) | 2 | L42542 | + | + | + | + | | | |
| RAN binding protein 2-like 1 (RANBP2L1) | 2 | AF012086 | | | | | | | |
| Ran GTPase activating protein 1 (RANGAP1) | 3 | X82260 | + | + | + | + | | + | <u> </u> |
| RAN, member RAS oncogene family (RAN) | 1 | M31469 | | † † | | 一 | | - | |
| (low match) RanBP2 (Ran-binding | -1 | D42063 | | 1 | | | | | |
| protein 2) (=U19248; L41840 sapiens | ' | D42063 | | | | | | | |
| nucleoporin (NUP358)) | | | | | | <u></u> | L | L | |
| ransforming growth factor, beta receptor II (70-80kD) (TGFBR2) | 4 | D50683 | + | + | + | + | | + | |
| RAP1A, member of RAS oncogene family (RAP1A) | 10 | M22995 | + | + | + | Ŧ | + | + | |
| RAR-related orphan receptor C (RORC) | 1 | U16997 | | | | | | + | |
| RAS guanyl releasing protein 2 (calcium and DAG-regulated) | 1 | Y12336 | + | + | | | | | |
| ras homolog gene family, member A (ARHA) | 12 | X05026 | + | + | + | + | + | + | high in ovary |
| ras homolog gene family, | 1 | X61587 | + | + | + | + | | | |
| member G (rho G) (ARHG) ras homolog gene family, | 2 | Z35227 | + | +- | + | | | + | |
| member H (ARHH) ras inhibitor (RIN1) | 2 | M37191 | | + | | L., | | - | |
| Ras-GTPase activating | 2 | AF053535 | + | + | + | + | | + | |
| protein SH3 domain- binding protein 2 (KIAA0660) | | | | | | | | | |
| Ras-GTPase-activating protein SH3-domain- binding protein (G3BP) | 3 | U32519 | + | + | + | + | | + | |
| ras-related C3 botulinum | 11 | M29871 | | ┼┤ | + | | | + | |
| toxin substrate 2 (rho family, small GTP binding protein Rac2) (RAC2) | | | | | | | | | |
| RAS-RELATED PROTEIN RAP-1B (GTP-BINDING PROTEIN SMG P21B) | 1 | P09526 | | | | | | | |
| RBQ-1 | 1 | X85133 | | + | + | + | | | |
| rearranged T cell receptor beta variable region (TCRB) (=X58810) | 1 | L06891 | | | | | | | |
| regulator of Fas-induced | 1 | AF057557 | В | | - | | + | | |
| apoptosis (TOSO) | | L | | 11 | | l | | | |

| WO 00/40/49 | | | | | | | | | | | _ |
|---|---------------|----------|------|------------------|---|---|---|----------|------|----|---------------|
| regulator of G protein signalling 6 (RGS6) | 1 | AF073920 | | + | | | | | | ., | |
| regulator of G-protein signalling 14 (RGS14) | 2 | AF037195 | + | + | + | + | | | | | |
| regulator of G-protein signalling 2, 24kD (RGS2) | 6 | L13391 | + | + | + | + | | + | | | |
| regulator of G-protein signalling 5 (RGS5) (49% | 1 | O15539 | | | | | | | | | |
| regulatory factor X, 4 (influences HLA class II | 1 | M69297 | | | + | + | | | | | 1 |
| expression) (RFX4) regulatory factor X, 5 (influences HLA class II | 2 | X85786 | | + | + | | | + | | | 1 |
| expression (RFX5) | 1 | M63488 | + | + | + | + | | + | | | $\frac{1}{2}$ |
| (RPA1) replication protein A3 | -1 | L07493 | | - | | | | \dashv | | | ┪ |
| (14kD) (RPA3) (low match) reproduction 8 (D8S2298E) | 1 | D83767 | | + | + | + | | | | | \dashv |
| l ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' | 2 | U94585 | + | + | + | + | | + | | · | 4 |
| requiem, apoptosis response zinc finger gene (REQ) | | | · | | | | | | | | |
| requiem, apoptosis response zinc finger gene (REQ) (=AF001433) (low match) | 1 | U94585 | | | | | | | | | |
| restin (Reed-Steinberg cell- expressed intermediate filament-associated | 1 | M97501 | В, Т | + | + | | | | | | |
| protein) (RSN) retinoblastoma 1 (including | 3 | L11910 | + | + | + | + | | | | | 1 |
| osteosarcoma) (RB1) retinoblastoma binding protein 2 homolog 1 | 1 | AF087481 | | | | | | | | | |
| (RBBP2H1) | 1 | S66427 | + | + | | | | | | | 1 |
| retinoblastoma-binding | 5 | S66431 | + | + | + | + | | + | | • | 1 |
| retinoblastoma-binding protein 4 (RBBP4) | 1 | X71810 | | + | + | + | | + | | | ٦ |
| retinoblastoma-binding protein 4 (RBBP4) | 1 | X74262 | | + | + | + | | + | | | 1 |
| retinoblastoma-binding protein 7 (RBBP7) | 1 | U35143 | | | | | | | | | ٦ |
| retinoblastoma-like 2 (p130) (RBL2) | 1 | X76061 | | + | + | + | | + | | | ٦ |
| retinoic acid receptor responder (tazarotene induced) 3 (RARRES3) | 1 | AF060228 | | + | | + | + | + | - | | |
| retinoic acid receptor, alpha (RARA) | 1 | X06538 | + | + | | + | | | | | |
| retinoic acid responsive (NN8-4AG) | 1 | U50383 | | + | | + | | + | | | |
| retinoid X receptor beta (RXR-beta) | 2 | X66424 | | + | + | Ť | | + | | | |
| REV3 (yeast homolog)-like, catalytic subunit of DNA polymerase zeta (REV3L) | 1 | AF035537 | | | | | | | | | |
| Rho GDP dissociation inhibitor (GDI) beta (ARHGDIB) | 23 | L07916 | + | + | + | + | + | + | | | |
| Rho GTPase activating protein 4 (ARHGAP4) | 2 | X78817 | + | + | L | | | | | | |
| Rho GTPase activating protein 4 (ARHGAP4) (low match) | 1 | P98171 | | | | | | | | | |
| Rho-associated, coiled-coil containing protein kinase 2 (ROCK2) | 1 | AB014519 | | | | | | | | | |
| ribonuclease 6 precursor (RNASE6PL) | 2 | U85625 | + | + | + | + | + | + | | | |
| | · | | | | | | | | | | _ |

| ribonuclease 6 precursor | | | | | | | | | |
|---|----|--------|-----|--|-------------|----------|---|----------|--------------------------------------|
| (RNASE6PL) (low match) | 1 | U85625 | | | | | | | ;, |
| ribonuclease, RNase A family, 2 (liver, eosinophil- derived neurotoxin) | 1 | X55988 | | | | | + | | |
| (RNASE2) | | | | _l | | | | | |
| nbonucleáse/angiogenin inhibitor (RNH) | 3 | M36717 | + | + | + | + | | + | |
| nbonucleoside diphosphate reductase M1 subunit | 1 | X65708 | | | | | | | |
| ribonucleotide reductase M2 polypeptide (non-exact 91%) | 1 | P31350 | | | | | | | |
| ribophorin I (RPN1) | 1 | Y00281 | + | + | + | + | | + | |
| ribophorin II (RPN2) | 1 | Y00282 | + | + | + | + | + | + | |
| ribosomal 18S rRNA | 3 | M10098 | | 1 | <u> </u> | <u> </u> | | | |
| ribosomal 28S RNA | 1 | M11167 | | | | ! | | <u> </u> | |
| ribosomal phosphoprotein P0, 5'UTR (low match) | 1 | D28418 | | | | | - | | |
| Ribosomal protein | 1 | | | | | | | 1 | |
| nbosomal protein L10 (RPL10) | 30 | L25899 | + | + | + | + | + | + | high in many libraries |
| RIBOSOMAL PROTEIN L10A (CSA-19) | 2 | P53025 | | | | | | | |
| ribosomal protein L11 (RPL11) | 4 | X79234 | + | + | + | + | + | + | Alveolar rhabdomyosarcoma |
| (RPL19) | 2 | L06505 | + | + | + | + | + | + | • |
| ribosomal protein L13 (PRL13) | 1 | P26373 | + | + | + | + | + | + | high in many libraries |
| ribosomal protein L14 (RPL14) | 4 | D87735 | + | + | + | + | + | + | high in many libraries |
| ribosomal protein L17 (RPL17) | 4 | X53777 | + | | | | | | blood only |
| ribosomal protein L18 (RPL18) | 10 | L11566 | + | + | + | + | | + | |
| ribosomal protein L18a (RPL18A) | 5 | L05093 | | + | + | + | + | + | High in fetal adrenal gland and skin |
| ribosomal protein L18a homologue | 2 | X80821 | | | | + | | | |
| ribosomal protein L19 (RPL19) | 15 | X63527 | + | + | + | + | + | + | |
| ribosomal protein L21 (RPL21) | 6 | U14967 | + | + | + | + | + | + | · |
| ribosomal protein L22 (RPL22) | 3 | D17652 | + | + | + | + | | + | |
| ribosomal protein L23 (RPL23) | 2 | X55954 | + | + | + | + | + | + | high in many libraries |
| пbosomal protein L23a (RPL23A) | 5 | U37230 | + | + | + | + | + | + | high in many libraries |
| ribosomal protein L26 (RPL26) | 8 | X69392 | + | + | + | + | + | + | |
| ribosomal protein L27 (RPL27) | 6 | L05094 | + | + | + | + | | + | |
| ribosomal protein L27a (RPL27A) | 10 | U14968 | + | + | + | + | + | + | |
| ribosomal protein L28 (RPL28) | 6 | U14969 | + | + | + | + | | + | |
| ribosomal protein L29 (RPL29) | 6 | U10248 | + | + | + | + | + | + | |
| ribosomal protein L3 (RPL3) | 81 | | + | + | + | + | + | + | high in many libraries |
| ribosomal protein L3 homologue | 81 | X06323 | | | | | | | |
| ribosomal protein L30 (RPL30) | 6 | X79238 | + . | + | + | + | + | + | high in lymphoma |
| ribosomal protein L30 (RPL30) (low score) | 1 | X79238 | | | | | | | |
| ribosomal protein L31 (RPL31) | 10 | X15940 | + | + | + | + | + | + | High in alveolar rhabdomyosarcoma |

| 342 + 7440 941 941 661 + 699 + 876 + 660 + 6844 + 966 + | + + + + + + + | + + + + + + | + + + + + + + + | + + + + + + | + + + + | high in barstead prostate high in many libraries |
|---|--|--|--|-------------|---------|--|
| 941 941 661 + 699 + 876 + 660 + 6844 + 966 + | + + + + + | + + + + + | + + + + | + + | + + + | prostate high in many libraries |
| 941 681 + 699 + 876 + 660 + 6844 + 966 + | + + + + | + + + + + + | + + + | + + | + | prostate high in many libraries |
| 681 + 699 + 876 + 660 + 6844 + 966 + | + + + + | + + + + | + + + | + | + | prostate high in many libraries |
| 699 + 876 + 660 + 6844 + 966 + | + + + + | + + + + | + + + | + | + | prostate high in many libraries |
| 876 + 660 + 6844 + 966 + | + + + | + | + | + | | |
| 660 + 6844 + 966 + | + | + | + | | + | |
| 966 + 966 | + | + | <u> </u> | + | 1 | high in many libraries |
| 966 + 966 | | | ++ | 1 | + | high in many libraries |
| 966 | + | + | ' | + | + | high in many libraries |
| | | 1 | + | + | + | High in alveolar rhabdomyosarcoma |
| 391 + | | + | | ╁ | ┼── | maddoniyodaroonia |
| 391 + | | | | | | |
| I | + | + | + | + | + | high in many libraries |
| 967 + | + | + | + | + | + | high in conorm |
| 072 + | + | + | + | + | + | High in uterus, and seminoma |
| 407 + | + | + | + | + | + | high in ovary |
| 953 | + | + | + | + | + | |
| 972 + | _ + | + | + | + | + | high in many libraries |
| 617 + | + | + | + | + | + | high in many libraries |
| 7152 | | | | | | |
| 505 + | + | + | + | + | + | high in many libraries |
| 124 | + | + | + | + | + | |
| 934 + | + | T + | + | + | + | |
| 405 + | + | + | + | + | + | |
| 854 + | + | + | + | + | + | High in prostate invasive tumor |
| 932 + | + | + | + | + | + | high in many libraries |
| 150 | | 1 | | | | |
| 757 + | + | + | + | + | + | high in many libraries |
| 206 + | + | + | + | + | + | high in many libraries |
| 880 | | | | | | |
| 498 + | + | + | + | + | + | high in many libraries |
| 483 + | + | + | + | + | + | high in CD34+/CD38- hematopoietic cells and skin tumor |
| 530 | + | + | + | | + | |
| 520 + | + | + | + | + | + | high in uterus |
| 716 + | + | + | + | + | + | high in barstead |
| | | | | | [| prostate |
| 554 | + | + | + | + | + | |
| 1 8 2 8 17 2 8 4 4 5 5 | 24 934 + 9354 + 932 + 937 + 938 + 930 + 930 + 931 + 931 + 932 + 933 + 934 + 935 + 955 + 957 + 957 + 958 + 95 | 24 + + + + + + + + + + + + + + + + + + + | 24 | 24 | 24 | 24 |

| ribosomal protein S28 | | T. Trende | | | | | | | |
|--|-------------|-------------|----------------|--|--------------|--------------|------------|--------------|------------------------|
| (RPS28) | 3 | U58682 | + | + | + | + | | + | · |
| ribosomal protein S29 (RPS29) | 2 | U14973 | + | + | + | + | + | + | |
| nbosomal protein S3 (RPS3) | 9 | X55715 | + | + | + | + | + | + | high in many libraries |
| ribosomal protein S3 (RPS3) (low match) | 1 | U14990 | | | | | T | | |
| ribosomal protein S3A (RPS3A) | 21 | Z83334 | | + | + | + | + | + | high in many libraries |
| ribosomal protein S3A (RPS3A) (low score) | 1 | M77234 | | + | + | | \vdash | \vdash | |
| ribosomal protein S4, X- linked (RPS4X) | 9 | M58458 | + | + | + | + | | + | high in ovary and |
| ribosomal protein S4, Y- linked (RPS4Y) | 2 | M58459 | + | + | + | + | + | + | Synovial sarcoma |
| ribosomal protein S5 (RPS5) | 4 | U14970 | + | + | +- | + | + | + | high in lymphoma |
| RIBOSOMAL PROTEIN S6 | 1 1 | P10660 | | + | | ┿ | ⊢ – | | ļ |
| (PHOSPHOPROTEIN NP33) | | | | ļ | | | | İ | |
| ribosomal protein S6 (RPS6) | 22 | M20020 | + | + | + | + | + | + | |
| ribosomal protein S6 (RPS6) (non-exact 86%) | 1 | M77232 | | | | | | | |
| ribosomal protein S6 | 3 | L07597 | + | + | + | + | | + | |
| kinase, 90kD, polypeptide 1 (RPS6KA1) ribosomal protein S6 | | | | | | | | | |
| kinase, 90kD, polypeptide 2 (RPS6KA2) | 1 | X85106 | | | | | | | |
| nbosomal protein S7 (RPS7) | 4 | Z25749 | | + | + | + | + | + | |
| ribosomal protein S8 (RPS8) | 6 | X67247 | | + | + | + | + | + | |
| ribosomal protein S9 (RPS9) | 8 | U14971 | | | | | | | colon tumor |
| ribosomal protein, large, PO (RPLP0) | | M17885 | 1 | | + | П | | + | · |
| ribosomal protein, large, P1 (RPLP1) | 12 | M17886 | Т | + | + | | + | | |
| ribosomal RNA 18S (≃M10098; K03432) (=polyadenylating sequence) | 11 | X03205 | | | | | | | · |
| ribosomal RNA 28S | 2 | M11167 | | | \vdash | | | | <u> </u> |
| ribosomal RNA, 16S | 1 | U25123 | - | +- | | | | | |
| ring finger protein (non- exact 58%) | 1 | AJ001019 | <u> </u> | | | | | | |
| ring finger protein 3 (RNF3) | 1 | AJ001019 | | | - | \vdash | | | |
| ring finger protein 4 (RNF4) | 3 | AB000468 | | + | + | + | | + | |
| ring zinc-tinger protein (ZNF127-Xp) | 3 | U41315 | | + | + | + | - | + | |
| RNA (guanine-7-) methyltransferase (RNMT) | 1 | AB007858 | | + | + | + | | + | |
| RNA binding motif protein 5 (RBM5) | 4 | U23946 | + | + | + | + | | + | |
| RNA binding motif, single stranded interacting protein 2 (RBMS2) | 1 | D28483 | | + | - | + | | + | |
| RNA helicase (putative), (Myc-regulated DEAD box protein) (MRD8) | 1 | X98743 | + | + | + | + | | + | |
| RNA helicase-related protein | 1 | AF083255 | | + | + | + | + | + | |
| RNA pol II largest subunit | 2 | X74872 | | | | | \dashv | | |
| RNA polymerase I subunit (RPA40) | 1 | AF008442 | | + | + | - | \dashv | + | |
| RTVP-1 protein | 2 | X91911 | + | + | + | + | - | + | |
| | | | <u> </u> | | | | | | |

| WO 00/40/49 | | | | | | | | | |
|---|---------------------------------------|-----------|---|-----|--------------|--|--------------|--|--|
| S100 calcium-binding | 2 | M81457 | | | + | | + | + | · · · · · · · · · · · · · · · · · · · |
| protein A10 (annexin II | | | | | | | | 1 | |
| ligand, calpactin I, light | | | | | | | | l | |
| polypeptide (p11)) (S100A10) | | 1 | | | | | | | |
| S100 calcium-binding | 7 | X80201 | | ++ | + | + | | + | |
| protein A11 (calgizzarin) | • | //00201 | | | | | | ŀ | |
| (S100A11) | | | |]] | | | | | |
| S100 calcium-binding | 3 | M80563 | В | | + | | + | | |
| protein A4 (calcium protein, | | | | | | i I | | | 1 |
| calvasculin, metastasin, | | | | | | | | | <u> </u> |
| murine placental | | | | | | 1 | | l | j |
| homolog)(S100A4) | | M21005 | | | + | + | | + | high in bone marrow |
| S100 calcium-binding protein A8 (calgranulin A) | , , , , , , , , , , , , , , , , , , , | 1005 | | | • | * | | * | migri in bone manow |
| (S100A8) | | | | | | | | | 1 |
| S100 calcium-binding | 14 | X06233 | | + | + | + | | | high in invasive |
| protein A9 (calgranulin B) | | | | | | | | 1 | larynx squamous cell |
| (S100A9) | | 1 1 | | | | | | | carcinoma |
| S164 gene | 1 | AF109907 | | | | | | | |
| S-adenosylmethionine | 3 | M88003 | + | + | + | + | <u> </u> | + | |
| decarboxylase 1 (AMD1) | _ | | | | | ł | ŀ | | i l |
| SB classII | 5 | M27487 | + | + | + | + | | + | |
| histocompatibility antigen | | | | | | | | | ŧ |
| alpha-chain | | | | | | <u> </u> | | | |
| SC35-interacting protein 1 | 5 | AF030234 | + | + | + | + | + | + | |
| (SRRP129) | | 117775 | + | + | + | + | | + | <u> </u> |
| scaffold attachment factor B (SAFB) | 1 | U72355 | * | + | _ | + | | _ | |
| scaffold attachment factor | | U72355 | | + | | - | - | - | |
| B (SAFB) (non-exact 78%) | ' | 072333 | | | | | | | |
| scRNA molecule. | 1 | L13713 | | + | | | \vdash | _ | |
| transcribed from Alu repeat | · | | | | | 1 | 1 | 1 | |
| SEC14 (S. cerevisiae)-like | 4 | D67029 | | + | + | + | | + | |
| (SEC14L) | | | | | | | | | |
| SEC23-like protein B | 2 | X97065 | + | + | + | + | | + | |
| (SEC23B) | | | | | | | | | |
| SEC63 (SEC63) | 1 | AF100141 | | + | + | l | | + | |
| secreted protein, acidic, | 7 | M25746 | | + | + | + | + | + | high in bone marrow |
| cysteine-rich (osteonectin) | | į | | | | 1 | | l | stroma |
| (SPARC) | | | | 1 | | Ļ., | | Щ | |
| secretory carrier | 1 | AF038966 | | + | | + | İ | l | ļ : |
| membrane protein 1 (SCAMP1) | | | | | ŀ | 1 | | 1 | |
| secretory carrier | 1 | AF005038 | + | + | + | + | + | + | |
| membrane protein 2 | l ' | Ai 000000 | • | 1 | ' | ` | ' | ` | |
| (SCAMP2) | | | | | | | | 1 | |
| secretory carrier | 1 | AF005039 | | | | 1 | | | |
| membrane protein 3 | | | | | ì | | | 1 | 1 |
| (SCAMP3) | | | | | | | | | |
| secretory granule | 1 | M33649 | | 1 | 1 | 1 | 1 | | |
| proteoglycan core (clones | | | | | | | [| l | |
| lambda-PG[6,7,8]) | 43 | X17519 | + | | ļ | + | ├ | + | |
| selectin L (lymphocyte adhesion molecule 1) | 43 | V1/218 | • | 1 | | * | | ~ | |
| (SELL) | | <u> </u> | | 1 | ł |] | 1 | ļ | 1 |
| selectin P ligand (SELPLG) | 13 | U02297 | + | + | | | | | |
| sema domain. | 2 | U60800 | | +- | | + | <u> </u> | + | ļ |
| immunoglobulin domain | _ | 000000 | | 1 |] | " | [| " | |
| (lg), transmembrane | | 1 | | 1 | | 1 | | 1 | |
| domain (TM) and short | | | | | ŀ | | 1 | | |
| cvtoplasmic domain. | | | | 1 | l | 1 | 1 | 1 | 1 |
| (semaphorin) 4D | | | | 1 | 1 | | l | 1 | |
| (SEMA4D) | | <u> </u> | | 1 | <u> </u> | <u> </u> | ļ | <u> </u> | |
| Ser/Arg-related nuclear | 4 | AF048977 | | + | + | + | + | + | 1 |
| matrix protein (plenty of prolines 101-like) | 1 | 1 | | i | | 1 | | | |
| (SRM160) | | | | | | 1 | 1 | 1 | |
| serine palmitoyltransferase | 1 | Y08685 | | ++ | + | + | | + | |
| subunit I (SPTI) | ' ' | .55555 | | 1 | · . | ` | 1 | 1 | |
| serine palmitoyltransferase, | 1 | AB011098 | + | + | + | + | 1 | + | |
| subunit II (LCB2) | • | | | | L_{-} | L | | L | |
| <u> </u> | | | | | | | | | |

| | | , _{RREF} | | | | | , | | | | |
|---|-----|---------------------|------|---|---|---|---|----|---------------------------------------|---|--|
| serine protease | 1 | J02907 | | | | | | | | | |
| serine protease inhibitor, Kunitz type, 2 (SPINT2) | 1 | U78095 | + | + | + | + | | + | | | |
| serine/threonine kinase 10 (STK10) | 1 | AB015718 | + | + | + | + | | + | | | |
| serine/threonine kinase 19 (STK19) | 1 | L26260 | +, | + | + | + | | | | | |
| serine/threonine kinase 4 (STK4) | 1 | U18297 | | + | | | | + | | | |
| serine/threonine protein kinase KKIALRE (KKIALRE) | 1 | X66358 | | + | + | + | | + | | | |
| serine/threonine protein- kinase (NIK) | 1 | Y10256 | | + | + | + | | | | | |
| SERINE/THREONINE- PROTEIN KINASE RECEPTOR R3 PRECURSOR (SKR3) | 1 | P37023 | | | | | | | | | |
| serologically defined colon cancer antigen 16 (NY-CO- 16) | 2 | AF039694 | | | | | | | | | |
| serologically defined colon cancer antigen 33 (SDCCAG33) | 1 | AF039698 | В, Т | + | + | | + | | | | |
| serologically defined colon cancer antigen 33 (SDCCAG33) (low score) | 1 | AF039698 | | | | | | | | | |
| serologically defined colon cancer antigen 33 (SDCCAG33) (low score) | 1 | AF039698 | | | | | | į. | | | |
| serum deprivation response (phosphatidylserine-binding protein) (SDPR) (=S67386) | 1 | AF085481.1 | | | | | | | | | |
| serum/glucocorticoid regulated kinase (SGK) | 2 | Y10032 | + | + | + | + | | + | | | |
| SET domain, bifurcated 1 (SETDB1) | 2 | D31891 | + | + | + | | | + | | | |
| SH2 domain protein 1A, Duncan's disease lymphoproliferative syndrome) (SH2D1A) | 1 | AF073019 | , | | | | | + | | | |
| SH3 binding protein (SAB) | 2 | AB005047 | + | + | + | + | | + | | - | |
| SH3 domain protein 1B (SH3D1B) | 4 | U61167 | + | | | + | | + | | | |
| SH3BGR PROTEIN (=21- GLUTAMIC ACID-RICH PROTEIN;21-GARP) (non- exact 82%aa) | 1 | P55822 | | | | | | | | | |
| SH3-binding domain glutamic acid-rich protein like (SH3BGRL) | 1 - | AF042081 | + | + | + | + | | + | | | |
| SH3-domain GRB2-like 1 (SH3GL1) | 1 | U65999 | + | + | + | + | | + | | | |
| SHC (Src homology 2 domain-containing) transforming protein 1 (SHC1) | 2 | X68148 | | + | + | + | | + | | | |
| siah binding protein 1 (SiahBP1) | 2 | U51586 | | + | + | + | | + | | | |
| siah binding protein 1 (SiahBP1) (non-exact, 69%) | T | U51586 | | | | | | | | | |
| Sialomucin CD164 (CD164) | 9 | D14043 | | | | | · | | | | |
| sialophorin (gpL115, leukosialin, CD43) (SNP) | 2 | J04536 | | | | | | | | | |
| sialyltransferase (STHM) | 1 | U14550 | | | + | + | | + | | | |
| sialyltransferase 1 (beta- galactoside alpha-2,6- sialytransferase) (SIAT1) | 2 | X17247 | + | + | + | + | + | + | · · · · · · · · · · · · · · · · · · · | | |

| galactosidase alpha-2.3-sialytransferase) (SIATAA) sialytransferase) (SIATAA) sialytransferase) (SIATAA) sialytransferase) (SIATAA) sialytransferase) (SIATAA) sialytransferase) (SIATAA) signal repotitions particle 1 | | | | | | | | | | | _ |
|---|--|---|----------|---|---|---|---------|---|---|-------------|---|
| Siapyltransferase 8 (alpha-2, 8- polysiapyltransferase) D (SIAT8D) | galactosidase alpha-2,3- | 1 | AF059321 | В | + | + | | + | + | 5 | |
| Signal peptidase 25kDa 1 | sialyltransferase 8 (alpha- 2, 8-polysialytransferase) D | 1 | L41680 | | + | | | | | | |
| signal recognition particle 14kD (hemologous Alu RNA-binding protein) (SRP14) signal recognition particle 5kD (SRP94) signal recognition particle 5kD (SRP94) signal recognition particle 5kD (SRP94) signal recognition particle 5kD (SRP3) signal recognition particle 7kD (SRP3) signal recognition particle 7kD (SRP3) signal recognition particle 7kD (SRP3) signal recognition particle 7kD (SRP3) signal recognition particle 7kD (SRP3) signal recognition particle 7kD (SRP3) signal sequence receptor (adapta (translocon-associated protein labla) (SSR1) signal sequence receptor, 2kD (SRP3) signal stransducer and activator of transcription (STAT5A) signal transducer and activator of transcription 2 signal stransducer and activator of transcription 2 signal transducer and activator of transcription 3 signal transducer and activator of transcription 3 signal transducer and activator of transcription 3 signal transducer and activator of transcription 5 A (STAT5A) signal transducer and activator of transcription 5 A (STAT5A) signal transducer and activator of transcription 5 A (STAT5A) signal transducing adaptor molecule (SR3 domain and ITAM motif) 1 (STAM) silencing mediator of retinoid and thyroid homone action (SMR1) similar to beta-transducin superfamily proteins (SAZD) similar to S. cerevisiae SSM (TEB4) ST protein 1 AJ010059.1 Siggier syndrome antigen 1 AB011169 + + + + + + + + + + + + + + + + + + + | signal peptidase 25kDa | 1 | L38950 | | 1 | | | | | | 1 |
| signal recognition particle \$40 (SRPS4) \$10 (SRP9) \$10 (SRP9) \$10 (SRP9) \$10 (SRP9) \$10 (SRP9) \$10 (SRP9) \$10 (SRP9) \$10 (SRP9) \$10 (SRP9) \$10 (SRP9) \$10 (SRP9) \$10 (SRP8) \$10 (SRP9) \$10 (SRP8) \$10 (SRP9) \$10 (SRP8) \$10 | signal recognition particle 14kD (homologous Alu RNA-binding protein) | 1 | X73459 | + | + | + | + | + | + | | |
| SkD (SRP9) Signal recognition particle receptor (docking protein') SRPR Signal regulatory protein, beta, 1 (SIRP-BETA-1) 5 | signal recognition particle 54kD (SRP54) | 1 | U51920 | | | | | | | | |
| receptor (docking protein') SRPR Signal regulatory protein, beta, 1 (SIRP-BETA-1) Signal sequence receptor, alpha (translocon- associated protein alpha) (SSR1) Signal sequence receptor, beta (translocon- associated protein beta) (SSR2) Signal transducer and activator of transcription (STATSA) Signal transducer and activator of transcription 2, 113KD (STAT2) Signal transducer and activator of transcription 3 (acute-phase response factor) (STAT3) Signal transducer and activator of transcription 5A (STATSA) Signal transducer and activator of transcription 3 (acute-phase response factor) (STAT3) Signal transducer and activator of transcription 5A (STATSA) Signal transducer and activator of transcription 5A (STATSA) Signal transducer and activator of transcription 5A (STATSA) Signal transducer and activator of transcription 5A (STATSA) Signal transducer and activator of transcription 5A (STATSA) Signal transducing adaptor molecule (SH3 domain and ITAM motif) 1 (STAM) Silencing mediator of retinoid and thyroid hornone action (SMRT) Similar to beta-transducin superfamily proteins (SAZD) Similar to Scerevisiae SSM4 (TEB4) Similar to yeast pre-mRNA 1 M62800 A1 (SZK0, INDEX SARO) (SSA1) Siggren syndrome antigen A1 (SZK0, INDEX SARO) (SSA1) Siggren syndrome antigen A1 (SZK0, INDEX SARO) (SSA1) Siggren syndrome antigen A1 (SSK0, INDEX SARO) (SSA1) INDEX SARON (SSA1) INDEX SARON INDEX SAR | 9kD (SRP9) | 2 | | | + | + | + | + | + | | |
| beta, 1 (SIRP-BETA-1) signal sequence receptor, alpha (translocon-associated protein alpha) (SSR1) signal sequence receptor, beta (translocon-associated protein beta) (SSR2) signal sequence receptor, beta (translocon-associated protein beta) (SSR2) signal transducer and activator of transcription (STATSA) signal transducer and activator of transcription 3 (squate-phase response factor) (STAT3) signal transducer and activator of transcription 3 (squate-phase response factor) (STAT3) signal transducer and activator of transcription 5A (STAT5A) signal transducer and activator of transcription 5A (STAT5A) signal transducer and activator of transcription 5A (STAT5A) signal transducing adaptor molecule (SH3 domain and ITAM motip) 1 (STAM) silencing mediator of retinoid and thyroid hormone action (SMRT) similar to beta-transducin superfamily proteins (SAZD) similar to S. cerevisiae SSM4 (TEB4) similar to S. cerevisiae SSM4 (TEB4) similar to yeast pre-mRNA 1 AB011169 + + + + + + + + + + + + + + + + + + + | receptor ('docking protein') SRPR | 5 | X06272 | | | | | | | | |
| alpha (translocon-associated protein alpha) (SSR1) signal sequence receptor, beta (translocon-associated protein beta) (SSR2) signal transducer and activator of transcription (STAT5A) signal transducer and activator of transcription 2, 113KD (STAT2) signal transducer and activator of transcription 3 (acute-phase response factor) (STAT3) signal transducer and activator of transcription 3 (squte-phase response factor) (STAT3) signal transducing adaptor molecule (SH3A) signal transducing adaptor molecule (SH3A) signal transducing adaptor molecule (SH3A) signal transducing adaptor molecule (SH3A) signal transducing adaptor molecule (SH3A) signal transducing adaptor molecule (SH3A) signal transducing adaptor molecule (SH3A) signal transducing adaptor molecule (SH3A) signal transducing adaptor molecule (SH3A) signal transducing adaptor 1 U37146 retinoid and thyroid hormone action (SMRT) similar to beta-transducin superfamily proteins (SAZD) similar to S. cerevisiae SSM4 (TEB4) similar to yeast pre-mRNA 1 AF025031 + + + + + + + + + + + + + + + + + + + | beta, 1 (SIRP-BETA-1) | | Y10376 | | + | | | | | | |
| beta (translocon-associated protein beta) (SSR2) signal transducer and activator of transcription (STAT5A) signal transducer and activator of transcription 2, 113KD (STAT2) signal transducer and activator of transcription 3 (acute-phase response factor) (STAT3) signal transducer and activator of transcription 3 (acute-phase response factor) (STAT3) signal transducer and activator of transcription 5A (STAT5A) signal transducing adaptor molecule (SH3 domain and ITAM motif) 1 (STAM) silencing mediator of retinoid and thyroid hormone action (SMRT) similar to beta-transducin superfamily proteins (SAZD) similar to S. cerevisiae SSM4 (TEB4) similar to Veast pre-mRNA splicing factors, Prp1/Zer1 and Prp6 SIT protein 1 AJ010059.1 Signal store, SA/Ro) (SSA1) Signal store, SA/Ro) (SSA1) Signal store, SA/Ro) (SSA1) Signal store, SA/Ro) (SSA1) Signal store, SA/Ro) (SSA1) Signal store, SA/Ro) (SSA1) M62800 A1 (SSAC) | alpha (translocon- associated protein alpha) (SSR1) | | | | | | | | | | |
| activator of transcription (STAT5A) signal transducer and activator of transcription 2, 113KD (STAT2) signal transducer and activator of transcription 3 (acute-phase response factor) (STAT3) signal transducer and activator of transcription 54 (STAT5A) signal transducer and activator of transcription 5A (STAT5A) signal transducer and activator of transcription 5A (STAT5A) signal transducing adaptor molecule (SH3 domain and ITAM motif) 1 (STAM) silencing mediator of retinoid and thyroid hormone action (SMRT) similar to beta-transducin similar to beta-transducin similar to S. cerevisiae (SAZD) similar to yeast pre-mRNA splicing factors, Prp1/Zer1 and Prp6 SIT protein 1 AJ010059.1 Slogren syndrome antigen A1 (SZKD, ribonucleoprotein autoantigen SS-A/Ro) (SSA1) Slogren syndrome antigen A1 (SZKD, | beta (translocon- associated protein beta) (SSR2) | _ | | | | | | | | | |
| activator of transcription 2, 113KD (STAT2) signal transducer and activator of transcription 3 (acute-phase response factor) (STAT3) signal transducer and activator of transcription 5A (STAT5A) signal transducing adaptor molecule (SH3 domain and ITAM molif) 1 (STAM) silencing mediator of retinoid and thyroid hormone action (SMRT) similar to beta-transducin superfamily proteins (SAZD) similar to S. cerevisiae SSM4 (TEB4) similar to yeast pre-mRNA splicing factors, Prp1/Zer1 and Prp6 SIT protein 1 AJ010059.1 SIT protein 1 AJ010059.1 SIT protein 1 AJ010059.1 SIGgren syndrome antigen A1 (52kD, ribonucleoprotein autoantigen SS-A/Ro) (SSA1) Siggren syndrome antigen A1 (52kD, Siggren syndrome antigen A2 (52kD, Siggren syndrome antigen A2 (52kD, Siggren syndrome antigen A3 (52kD, Siggren syndrome antigen A3 (52kD, Siggren syndrome antigen A3 (52kD, Sig | activator of transcription (STAT5A) | · | | + | + | + | + | + | | | |
| activator of transcription 3 (acute-phase response factor) (STAT3) signal transducer and activator of transcription 5A (STAT5A) signal transducing adaptor molecule (SH3 domain and ITAM motif) 1 (STAM) silencing mediator of retinoid and thyroid hormone action (SMRT) similar to beta-transducin similar to beta-transducin similar to S. cerevisiae (SAZD) similar to S. cerevisiae SSM4 (TEB4) similar to yeast pre-mRNA splicing factors, Prp1/Zer1 and Prp6 SIT protein 1 AJ010059.1 Sjogren syndrome antigen A1 (52kD, indonucleoprotein autoantigen SS-A/Ro) (SSA1) Sjogren syndrome antigen A1 (52kD, indonucleoprotein autoantigen SS-A/Ro) (SSAS) Sjogren syndrome antigen A1 (52kD, indonucleoprotein autoantigen SS-A/Ro) (SSA1) Sjogren syndrome antigen A1 (52kD, indonucleoprotein autoantigen SS-A/Ro) (SSA1) | activator of transcription 2, 113KD (STAT2) | | | | | | | | + | | |
| signal transducer and activator of transcription 5A (STAT5A) signal transducing adaptor molecule (SH3 domain and ITAM motif) 1 (STAM) silencing mediator of retinoid and thyroid hormone action (SMRT) similar to beta-transducin superfamily proteins (SAZD) similar to S. cerevisiae SM4 (TEB4) similar to yeast pre-mRNA splicing factors, Prp1/Zer1 and Prp6 SIT protein 1 AJ010059.1 Sjogren syndrome antigen A1 (S2kD, ISJOgren synd | activator of transcription 3 (acute-phase response | 3 | L29277 | | | | | | | - | |
| molecule (SH3 domain and ITAM motif) 1 (STAM) silencing mediator of retinoid and thyroid hormone action (SMRT) similar to beta-transducin 1 U02609 + + + + + + superfamily proteins (SAZD) similar to S. cerevisiae 1 AB011169 + + + + + + SSM4 (TEB4) similar to yeast pre-mRNA 1 AF026031 + + + + + + splicing factors, Prp1/Zer1 and Prp6 SIT protein 1 AJ010059.1 SJogren syndrome antigen A1 (S2kD, ribonucleoprotein autoantigen SS-A/Ro) (SSA1) SJogren syndrome antigen 1 M62800 A1 (S2kD, SJogren syndrome antigen SJogren syndrome antigen I M62800 A1 (S2kD, SJogren syndrome antigen I M62800 A1 (S2kD, SJogren syndrome antigen I M62800 A1 (S2kD, SJogren syndrome antigen I M62800 A1 (S2kD, SJogren syndrome antigen I M62800 A1 (S2kD, SJogren syndrome antigen I M62800 A1 (S2kD, SJogren syndrome antigen I M62800 A1 (S2kD, SJOgren syndrome antigen I M62800 A1 (S2kD | signal transducer and activator of transcription 5A (STAT5A) | | | + | + | + | + | + | + | | |
| retinoid and thyroid hormone action (SMRT) similar to beta-transducin 1 U02609 + + + + + + + + + + + + + + + + + + + | molecule (SH3 domain and ITAM motif) 1 (STAM) | | | | | | | | | | |
| superfamily proteins (SAZD) similar to S. cerevisiae 1 AB011169 + + + + + + SSM4 (TEB4) similar to yeast pre-mRNA 1 AF026031 + + + + + + + + + + + + + + + + + + + | retinoid and thyroid hormone action (SMRT) | | | | | | | | | | |
| SSM4 (TEB4) similar to yeast pre-mRNA splicing factors, Prp1/Zer1 and Prp6 SIT protein 1 AJ010059.1 Slogren syndrome antigen 2 M62800 + A1 (52kD, ribonucleoprotein autoantigen SS-A/Ro) (SSA1) Slogren syndrome antigen 1 M62800 A1 (52kD, | superfamily proteins (SAZD) | | | + | | | | | | | |
| splicing factors, Prp1/Zer1 and Prp6 SIT protein 1 AJ010059.1 Sjogren syndrome antigen 2 M62800 + | SSM4 (TEB4) | | | | | | <u></u> | | | | |
| Sjogren syndrome antigen 2 M62800 + A1 (52kD, ribonucleoprotein autoantigen SS-A/Ro) (SSA1) | splicing factors, Prp1/Zer1 and Prp6 | , | · _ | + | + | + | + | | + | | |
| A1 (52kD, ribonucleoprotein autoantigen SS-A/Ro) (SSA1) Siggren syndrome antigen 1 M62800 A1 (52kD, | F | | I : | | | | | | | | |
| Sjogren syndrome antigen 1 M62800 A1 (52kD, | A1 (52kD, ribonucleoprotein autoantigen SS-A/Ro) | 2 | M62800 | | | | | + | | | |
| ribonucleoprotein autoantigen SS-A/Ro) (SSA1) (non-exact 63%) (match to zinc finger) | Sjogren syndrome antigen A1 (52kD, ribonucleoprotein autoantigen SS-A/Ro) (SSA1) (non-exact 63%) | 1 | M62800 | | | | | | | · | |
| SKAP55 homologue 1 AJ004886 + + + + + + (SKAP-HOM) | SKAP55 homologue | 1 | AJ004886 | | + | + | + | | + | | |
| 3kb1 (S. pombe) homolog 2 AF015913 + <td< td=""><td>skb1 (S. pombe) homolog</td><td>2</td><td>AF015913</td><td>+</td><td>+</td><td>+</td><td>+</td><td></td><td>+</td><td></td><td></td></td<> | skb1 (S. pombe) homolog | 2 | AF015913 | + | + | + | + | | + | | |

| lekeletel musele abundant | | | | | | | | | |
|--|-------------|----------|----------------|---|---|---|----|----------|---|
| skeletal muscle abundant protein | 1 | X87613 | + | + | + | + | | + | , |
| SMA3 (SMA3) | 1 | X83300 | + | + | | + | | + | - |
| small acidic protein | 3 | U51678 | + | + | + | + | ٠. | + | |
| small EDRK-rich factor 2 (SERF2) | 2 | Y10351 | + | + | + | + | + | + | high in fetal lung |
| small inducible cytokine A5 (RANTES) (SCYA5) | 2 | M21121 | + | + | + | + | + | + | high in many libraries |
| small inducible cytokine subfamily C, member 2 (SCYC2) | 1 | D63789 | | | | | | | |
| small nuclear ribonucleoprotein polypeptide B" (SNRPB2) | 2 | M15841 | | + | + | + | | + | |
| small nuclear ribonucleoprotein polypeptide N (SNRPN) | 4 | J04815 | + | + | + | + | + | + | |
| small nuclear ribonucleoprotein polypeptides B and B1 (SNRPB) | 2 | J04564 | + | + | + | + | | + | |
| small nuclear RNA activating complex, polypeptide 5, 19kD (SNAPC5) | 1 | AF093593 | + | + | + | + | | + | |
| smallest subunit of ubiquinol-cytochrome c reductase | 1 | D55636 | + | + | + | + | + | + | high in fetal lung |
| SMC (mouse) homolog, X chromosome (SMCX) | 1 | L25270 | + | + | + | + | | + | |
| SMT3B protein (2) | 2 | X99585 | + | + | + | + | + | + | <u> </u> |
| SNARE protein (YKT6) | | U95735 | | | | | - | <u> </u> | |
| (low match) SNC19 | | | | | | | | | |
| 1 1 | 1 | U20428 | | | | | | | |
| SNC73 protein (SNC73) | 2 | J00220 | + | + | | + | + | + | high in many libraries |
| solute carrier family 1 (neutral amino acid transporter), member 5 (SLC1A5) | 2 | U53347 | | + | | + | | + | |
| Solute carrier family 11 (proton-coupled divalent metal ion transporters), member 1 (SLC11A1) | 7 | D50403 | + | | | | | | |
| solute camer family 17 (sodium phosphate), member 3 (SLC17A3) | 1 | U90545 | | | | + | | | |
| solute carrier family 19 (folate transporter), member 1 (SLC19A1) | 1 | U17566 | B, lymphoma | + | | | + | | |
| solute carrier family 2 (facilitated glucose transporter), member 1 (SLC2A1) | 1 | K03195 | + | + | + | + | + | + | |
| solute carrier family 23 (nucleobase transporters), member 2 (SLC23A2) | 3 | D87075 | | + | + | + | | + | |
| solute carrier family 25 (mitochondrial carrier, oxoglutarate carrier), member 11 (SLC25A11) | 1 | AF070548 | B, T | + | + | | + | + | |
| solute carrier family 31 (copper transporters), member 2 (SLC31A2) | 3 | U83461 | | + | | + | | | |
| solute carrier family 4, anion exchanger, member 2 (erythrocyte membrane protein band 3-like 1) (SLC4A2) | 1 | X62137 | | + | + | | | + | |
| solute carrier family 4, sodium bicarbonate cotransporter, member 8 (SLC4A8) | 1 | AB018282 | | + | | | | | |

WO 00/40749

| W 0 00/40/49 | | | | | | | | r | C1/CA00/00005 |
|---|--|--------------|---------------------------------------|--|----------|----------|--|----------|---------------------------------------|
| solute carner family 7 (cationic amino acid | 2 | M80244 | T, W | + | + | | + | Γ | : |
| transporter, v+ system) | | | | | | | 1 | ĺ | |
| member 5 (SLC7A5) | | | 1 | l | | İ | 1 | ļ | |
| solute carrier family 7 | 3 | D87432 | + | + | + | | | + | |
| transporter, y+ system), | |] | 1 | | 1 | i | | | İ |
| member 6 (SLC7A6) | | | 1 | | | 1 | | · | |
| solute carrier family 7 | 1 | D87432 | | | | 1 | | 1 | |
| (cationic amino acid transporter, y+ system). | | İ | | | 1 | 1 | | | |
| member 6 (SLC7A6) (non- | l | | İ | ļ | | | Ī | | |
| exact 77%) | | l | <u> </u> | 1 | ļ | 1 | | | i : |
| solute carrier family 9 (sodium/hydrogen | 1 | AF030409 | | + | + | + | | + | |
| exchanger), isoform 6 | | | 1 | 1 | | l | 1 | | |
| (SLC9A6) | | | ļ | [| | 1 | | | |
| somatic cytochrome c (HCS) | 2 | M22877 | | | | | | | |
| SON DNA binding protein | 2 | X63753 | | + | + | + | _ | + | |
| (SON) | | /// | | 1 | ' | ' | | l ' | |
| son of sevenless | 1 | L13858 | + | + | | + | | | |
| (Drosophila) homolog 1 (SOS1) | Í | | | | | 1 | | | |
| sorcin (SRI) | 1 | M32886 | | | | | | | |
| sortilin 1 (SORT1) | 2 | X98248 | | ++ | | + | | + | |
| sortilin-related receptor. | 6 | Y08110 | | + | | ├— | - | | |
| L(DLR class) A repeats- containing (SORL1) | | | | | | 1 | | | |
| sorting nexin 1 (SNX1) | 3 | U53225 | + | ++ | | + | | + | |
| sorting nexin 2 (SNX2) | 2 | AF043453 | · · · · · · | ļ <u>'</u> | <u> </u> | <u> </u> | | | |
| sorting nexin 6 (SNX6) | 1 | AF121856.1 | | <u> </u> | | | | | |
| (=U83194.1 TRAF4- | ' | A1 12 1030.1 | | | | ! | | | |
| associated factor 2) | | | | | | | | | |
| Sp3 transcription factor (SP3) | 1 | X68560 | + | + | + | + | | + | |
| Sp3 transcription factor (SP3) | 4 | M97191 | + | + | + | + | | + | · · |
| special AT-rich sequence | 1 | M97287 | | + | | | \vdash | | |
| binding protein 1 (binds to | | | | | | | | ĺ | |
| nuclear matrix/scaffold- associating DNA's) | | | | | | | | | |
| (SATB1) | | | | | | | ı | | |
| speckle-type POZ protein | 4 \ | AJ000644 | · · · · · · · · · · · · · · · · · · · | | | | - | | |
| (SPOP) speckle-type POZ protein | 1 | AJ000644 | | | | | | | · · · · · · · · · · · · · · · · · · · |
| (SPOP) (non-exact) | | | | | | | | | , [|
| spectrin SH3 domain binding protein 1 | 6 | U87166 | + | + | + | + | | | |
| (SSH3BP1) | | | | | | l | | i | |
| Spectrin, alpha, non- | 2 | J05243 | | + | + | | _ | + | |
| erythrocytic 1 (alpha-fodrin) (SPTAN1) | | | • | 1 | | | - 1 | - 1 | |
| spermidine/spermine N1- | 11 | M55580 | | | | \dashv | \dashv | | |
| acetyltransferase (SAT) | | | | | | | | | |
| spermidine/spermine N1- acetyltransferase (SAT) | 1 | U40369 | | <u> </u> | | | | | |
| (non-exact, 84%) | | | | | | | ļ | ŀ | |
| spermine synthase (SMS) | 1 | AD001528 | + | + | + | + | \neg | + | |
| SPF31 (SPF31) | 1 | AF083190 | + | + | + | 7 | | + | |
| sphingomyelin phosphodiesterase 1, acid | The state of the s | X52679 | | + | + | | + | | |
| lysosomal (acid | ĺ | } | | | | - 1 | | Ì | |
| sphingomyelinase) | | [| | | | | | | |
| (SMPD1) SPINDLIN HOMOLOG | | Connec | | | | | | | |
| (PROTEIN DXF34) | 1 | Q99865 | | | | | | | |
| spinocerebellar ataxia 1 | 3 | X79204 | В | +- | \dashv | | + | \dashv | |
| (olivopontocerebellar ataxia 1, autosomal dominant, | | Ì | | | - 1 | | | | |
| ataxin 1) (SCA1) | | | | | | | | - 1 | |
| | | | | | | | | | |

| WO 00/40/49 | | | | | | | | • • | C1/CA00/00003 |
|--|-----|----------|----|---|---|---|---|-----|---------------|
| spinocerebellar ataxia 2 (olivopontocerebellar ataxia 2, autosomal dominant, ataxin 2) (SCA2) | 1 | U70323 | В | | | | + | | · |
| spinocerebellar ataxia 7 (olivopontocerebellar atrophy with retinal degeneration) (SCA7) | 2 | AJ000517 | | + | | | | | |
| spliceosome associated protein (SAP 145) | 3 | U41371 | | + | + | + | + | + | |
| splicing factor (CC1.3) (CC1.3) | 2 | L10910 | + | + | + | + | + | + | |
| splicing factor SRp40-1 (SRp40) | 7 | U30826 | + | + | + | + | + | + | |
| splicing factor, arginine/serine-rich 11 (SFRS11) | 3 | M74002 | В | + | + | | + | + | |
| splicing factor, arginine/serine-rich 7 (35kD) (SFRS7) | 4 | L41887 | | + | + | + | | + | |
| Src-like adapter protein (non-exact, 76%aa) | 1 | U30473 | | | | | | | |
| Src-like-adapter (SLA) | 6 | D89077 | | + | + | + | | + | |
| Src-like-adapter (SLA) (low match) | 1 - | D89077 | | | | | | | |
| Src-like-adapter (SLA) (low score) | 1 | U44403 | | | | | | | |
| stannin (SNN) | 2 | AF030196 | + | + | + | + | | + | |
| STAT induced STAT inhibitor 3 (SSI-3) | 1 | AB004904 | | | | + | | | |
| STE20-like kinase 3 (MST-3) | 2 | AF024636 | + | + | + | + | | + | |
| step II splicing factor SLU7 (SLU7) | 1 | AF101074 | | + | | + | + | + | |
| steroid sulfatase | 1 | M17591 | | | | | | | |
| steroid sulfatase (microsomal), arylsulfatase C, isozyme S (STS) | 1 | J04964 | | + | + | + | | | · |
| sterol carrier protein 2 (SCP2) | 1 | M55421 | | + | + | + | + | + | |
| sterol O-acyltransferase (acyl-Coenzyme A: cholesterol acyltransferase) 1 (SOAT1) | 7 | AF059202 | | | | | + | | |
| stimulated trans-acting factor (50 kDa) (STAF50) | 6 | X82200 | + | + | | + | | | |
| Striatin, calmodulin-binding protein (STRN) (low match, 71%aa) | 1 | U17989 | | | | | | | |
| Stromal antigen 2 (STAG2) | 2 | Z75331 | | 1 | + | + | + | + | |
| stromal interaction molecule 1 (STIM1) | 3 | U52426 | + | + | + | + | | + | |
| structure specific recognition protein 1 (SSRP1) | 1 | M86737 | | + | + | + | | + | |
| succinate dehydrogenase complex, subunit A, flavoprotein (Fp) (SDHA) | 5 | L21936 | | | + | | | | |
| succinate dehydrogenase complex, subunit B, iron sulfur (Ip) (SDHB) | 1 | U17248 | + | + | + | + | | + | |
| succinate dehydrogenase complex, subunit C, integral membrane protein, 15kD (SDHC) | | U57877 | + | + | + | + | | + | |
| succinate dehydrogenase complex, subunit D, Integral membrane protein (SDHD) | 3 | AB006202 | | + | + | | + | | |
| succinate-CoA ligase, GDP-forming, beta subunit (SUCLG2) | 1 | AF058954 | | + | + | + | + | + | |
| | | | 12 | | | | | | |

| WO 00/40749 | | | | | | | | PC | 71/CA00/00005 |
|---|----|----------|----------|---|---|---|---|----|---------------|
| succinyl CoA synthetase | 1 | Z68204 | | | | | | | 1. |
| sudD (suppressor of bimD6, Aspergillus | 2 | AF013591 | | + | | | + | + | |
| nidulans) homolog (SUDD) sulfotransterase family 1A, phenol-preferring, member 1 (SULT1A1) | 1 | L19999 | <u> </u> | + | | | + | + | |
| sulfotransferase family 1A, phenol-preferring, member 3 (SULT1A3) (non-exact 67%) | 1 | U37686 | | | | | | | |
| superoxide dismutase 1, soluble (amyotrophic lateral sclerosis 1 (adult)) (SOD1) | 4 | X02317 | | + | + | | + | + | |
| superoxide dismutase 2, mitochondrial (SOD2) | 5 | Y00985 | | + | + | + | + | + | |
| supervillin (SVIL) | 2 | AF051851 | | 1 | + | + | | + | |
| suppression of tumorigenicity 5 (ST5) | 2 | U15131 | | + | | + | | + | |
| suppression of tumorigenicity 5 (ST5) (non-exact 82%) | 1 | U15779 | | | | | | | |
| suppressor of K+ transport defect 1 (SKD1) | 1 | AF038960 | | | + | + | | | |
| suppressor of Ty (S.cerevisiae) 3 homolog (SUPT3H) | 1 | AF064804 | + | + | + | + | | + | |
| suppressor of Ty (S.cerevisiae) 4 homolog 1 (SUPT4H1) | 2 | U38817 | + | + | + | + | | + | |
| suppressor of Ty (S.cerevisiae) 5 homolog (SUPT5H) | 2 | U56402 | | + | | | | + | · |
| suppressor of Ty (S.cerevisiae) 6 homolog (SUPT6H) | 2 | U46691 | + | + | + | + | + | + | |
| suppressor of variegation 3-9 (Drosophila) homolog 1 (SUV39H1) | 1 | AF019968 | | + | + | + | | | |
| survival of motor neuron 1, telomeric (SMN1) | 1 | U18423 | | | | | | | |
| SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a, member 1 (SMARCA1) (non-exact, 75%) | | M88163 | | | + | + | | + | |
| SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a, member 2 (SMARCA2) | 2 | D26155 | | + | | | | | |
| SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a, member 4 (SMARCA4) | 1 | D26156 | + | + | + | + | + | + | |
| SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily c, member 2 (SMARCC2) | 4 | U66616 | + | + | + | + | + | + | |
| SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily e, member 1 (SMARCE1) | 2 | AF035262 | B, W | + | + | | + | + | |
| synaptobrevin-like 1 (SYBL1) | 1 | X95803 | | + | + | | | + | |
| synaptosomal-associated protein, 23kD (SNAP23) | 2 | AJ011915 | | + | + | + | | + | |
| syndecan binding protein (syntenin) (SDCBP) | 15 | AF006636 | + | + | + | + | | + | |

| cofactor-1 Idiated 3A (X-linked): andocardial fibrelastosis 2: Barth syndrome) (TAZ) + + + + + + + + + + + + + + + + + + + | WO 00/40749 | | | | | | | | P | CT/CA00/00005 |
|--|---|-------|-----------|---|----------|--------------|--|--|--------------|---------------|
| Syntaxin 16 | translocated to X | 2 | X79201 | | + | | | | | |
| Syntaxin 6 (STX6) | | 1 . | AF038897 | | + | | ┼- | | | |
| Syntaxin 6 (STX6) | syntaxin 3A (STX3A) | 2 | U32315 | | + | | ┝┿ | | + | |
| SYMIAXIN BINDING PROTEIN CUNC-18C SYMIAWIN-18C I AP008937 SYMIAWIN-18C I AP008938 SYMIAWIN-18C I | L | _ | 1 | | + | <u> </u> | ļ | | <u> </u> | |
| PROTEIN 3 (UNC-18 HOMOLOG 3) (UNC-18C) syntaxin-18C | , . | | | | | ļ | <u> </u> | <u> </u> | ├ | ļ <u></u> . |
| SYT Interacting protein (SIP) AP080561 + + + + + + + + | PROTEIN 3 (UNC-18 HOMOLOG 3) (UNC-18C) | | | | | _ | | : | | |
| (SIP) Toell activation, increased late expression (TACTILE) Toell receiptor V alpha gene segment V-alpha-7 (clone ICRR1) Toell receiptor V alpha gene segment V-alpha-7 (clone ICRR1) Toell receiptor V alpha gene segment V-alpha-27 (clone ICRR1) Toell receiptor V alpha gene segment V-alpha-27 (clone ICRR1) Toell receiptor-associating cofactor-1 (affazzin (cardiomyopathy, dilated 3A (X-dinked); endocardial fibroelastosis endocardial fibroelastosis (clone) (clon | , - | 1 | AF008937 | | | | | | | |
| late expression (TACTILE) | (SIP) | - | AF080561 | | + | + | + | | + | |
| gene segment V-alpha-7 (clone IGRA11) I cell receptor V alpha 27 T3 receptor-associating cofactor-1 Iafazzin (cardiomyopathy, dialeted 3A (X-Inked); and cardial fibroelastosis 2; Barth syndrome) (TAZ) TAFIIT00 protein (non-exact 53%) Iankyrase, IRF1: 1 | late expression (TACTILE) | | M88282 | | | | + | | | |
| Toell receptor V alpha 1 | gene segment V-alpha-7 | 2 | X58744 | | | | | | | |
| gene segment V-aipha-w27 T3 receptor-associating | | 1 | Y58740 | | + | | . | ļ | ┢ | |
| T3 receptor-associating cofactor. I lafazzin (cardiomyopathy, dilated 3A (X-linked); endocardial fibroelastosis 2: Barth syndrome) (TAZ) TAPITIO0 protein (non-exact 53%) [Inhyrase, TIR+1; interacting ankyrin-related ADP-ribose polymerase (TNKS) [TAPI, TAP2, LMP2, LMP7] [Inhyrase, TIR+1; interacting ankyrin-related ADP-ribose polymerase (TNKS) [TAPI, TAP2, LMP2, LMP7] [Inhyrase, TIR+1; interacting ankyrin-related ADP-ribose polymerase (TNKS) [TAPI, TAP2, LMP2, LMP7] [Inhyrase, TAR1, TAP2, LMP2, LMP7] [Inhyrase, TAR1, TAP2, LMP2, LMP7] [Inhyrase, TAR2, TAR2, TAR3, T | Igene segment V-alpha-w27 | ' | 7,50740 | | 1 | | 1 | | } | |
| Initialization Carolimorpopathy | T3 receptor-associating | 5 | 583390 | + | + | + | + | + | + | |
| dilated 3A (X-linked); and condital fibroelastosis 2; Barth syndrome) (TAZ) TAPITIOD protein (non-exact 53%) is an interacting ankyrin-related ADP-ribose polymerase (TNKS) TAPIT, TAP2, LMP2, LMP7 and DOB TAR DNA-binding protein 6 U23731 + + + + + + + + + + + + + + + + + + + | | | ļ | | | | | | | |
| TAPITIOD protein (non-exact 53%) Iankyrase, IRF1- 1 | dilated 3A (X-linked); endocardial fibroelastosis | 1 | X92763 | + | + | | + | | + | |
| Iankyrase, TRF1. | TAFII100 protein (non- | 1 | U80191 | | | | | | _ | · . — — — — |
| interacting ankyrin-related ADP-ribose polymerase (TNKS) TAP1, TAP2, LMP7 and DOB TAR DNA-binding protein- 43 Tat interactive protein (80kD) (TIP90) TARA box binding protein (TBP)-associated factor, RNA polymerase II, C1, 130kD (TAF2C) TATA box binding protein (TBP)-associated factor, RNA polymerase II, C5, 130kD (TAF2C) TATA box binding protein (TBP)-associated factor, RNA polymerase II, C7, 130kD (TAF2C) TATA box binding protein (TBP)-associated factor, RNA polymerase II, C7, 130kD (TAF2C) TATA box binding protein TATA box binding protein TATA box binding protein TBP)-associated factor, RNA polymerase II, I, 28kD TATA box binding protein TAPA-bax C1 (TAF2C) TATA box binding protein TAPA-bax C1 (TAF2C) TATA box binding protein TAPA-bax C1 (TAF2C) TATA box C1 (TAF2C) TATA box Dinding protein TAPA-bax C1 (TAF2C) TATA box Dinding protein TAPA-bax C1 (TAF2C) TAPA-bax C1 (TAF2C) TAPA box Dinding protein TAPA-bax C1 (TAF2C) TAPA box Dinding protein TAPA-bax C1 (TAF2C) TAPA box Dinding protein TAPA-bax C1 (TAF2C) TAPA box Dinding protein TAPA-bax C1 (TAF2C) TAPA box Dinding protein TAPA-bax C1 (TAF2C) TAPA box Dinding protein TAPA-bax C1 (TAF2C) TAPA box Dinding protein TAPA-bax | | 4 | NEADOCE O | | <u></u> | | <u> </u> | | <u> </u> | |
| and DOB TAR DNA-binding protein- 43 Tat interactive protein (60kD) (TiP60) TATA box binding protein (TBP)-associated factor, RNA polymerase II, C1, 130kD (TAF2C1) (non- exact, 55%) TATA box binding protein (TBP)-associated factor, RNA polymerase II, F, 55kD (TAF2F) TATA box binding protein (TBP)-associated factor, RNA polymerase II, G, 32kD (TAF2G) TATA box binding protein (TBP)-associated factor, RNA polymerase II, G, 32kD (TAF2G) TATA box binding protein (TBP)-associated factor, RNA polymerase II, 1, 28kD (TAF2G) TATA box binding protein (TBP)-associated factor, RNA polymerase II, 1, 28kD (TAF2G) TATA box binding protein (TBP)-associated factor, RNA polymerase II, 1, 28kD (TAF2G) Tax1 (human 1-cell leukemia virus type I) binding protein 1 (TAX1BP1) Tax2 (human 1-cell leukemia virus type I) binding protein 1 (TAX1BP1) T-box 2 (TBX2) (non-exact 1 T-box 2 (TBX2) (non-exact 1 T-box 2 (TBX2) (non-exact 1 T-box 2 (TBX2) (non-exact 1 T-box 2 (TBX2) (non-exact 1 T-box 2 (TBX2) (non-exact 1 T-box 2 (TBX2) (non-exact 1 T-cell death-associated 1 gene 8 (TDAG8) T-cell leukemia/lymphoma 1 X82240 T-cell receptor (delta D2- T-cell receptor (delta D2- T-cell receptor (delta D2- T-cell receptor (delta D2- T-cell receptor (delta D2- T-cell receptor (delta D2- | interacting ankyrin-related ADP-ribose polymerase | 1 | AF082556 | | + | + | + | | + | |
| 13 Interactive protein (60kD) (TIP60) 2 U40989 + + + + + + + + + + + + + + + + + + | TAP1, TAP2, LMP2, LMP7 | 1 | X66401 | | | | | | | |
| (60kD) (TIP60) TATA box binding protein (TBP)-associated factor, RNA polymerase II, C1, 130kD (TAF2C1) (non- exact, 55%) TATA box binding protein (TBP)-associated factor, RNA polymerase II, F, 55kD (TAF2C) TATA box binding protein (TBP)-associated factor, RNA polymerase II, G, 32kD (TAF2G) TATA box binding protein (TBP)-associated factor, RNA polymerase II, G, 32kD (TAF2G) TATA box binding protein (TBP)-associated factor, RNA polymerase II, i, 28kD (TAF2I) Tax1 (human 1-ceil | | 6 | U23731 | + | + | + | + | | + | |
| (TBP)-associated factor, RNA polymerase II, C1, 130kD (TAF2C1) (non-exact, 55%) TATA box binding protein (TBP)-associated factor, RNA polymerase II, F, 55kD (TAF2F) TATA box binding protein 2 U21858 + + + + + + (TBP)-associated factor, RNA polymerase II, G, 32kD (TAF2G) TATA box binding protein 1 D63705 + + + + + + (TBP)-associated factor, RNA polymerase II, I, 28kD (TAF2I) TATA box binding protein 1 U33821 + + + + + + + (TBP)-associated factor, RNA polymerase II, I, 28kD (TAF2I) Tax1 (human T-cell 1 U33821 + + + + + + + + + + + + + + + + + + + | (60kD) (TIP60) | 2 | U40989 | + | + | + | + | | + | |
| (TBP)-associated factor, RNA polymerase II, F, 55kD (TAF2F) TATA box binding protein (TBP)-associated factor, RNA polymerase II, G, 32kD (TAF2G) TATA box binding protein (TBP)-associated factor, RNA polymerase II, I, 28kD (TAF2G) TATA box binding protein 1 D63705 + + + + + + + + + + + + + + + + + + + | (TBP)-associated factor, RNA polymerase II, C1, 130kD (TAF2C1) (non- exact, 55%) | 1 | O00268 | | | | | | | |
| (TBP)-associated factor, RNA polymerase II, G, 32kD (TAF2G) TATA box binding protein (TBP)-associated factor, RNA polymerase II, I, 28kD (TAF2I) Tax1 (human T-cell 1 U33821 + + + + + + + + + + + + + + + + + + + | (TBP)-associated factor, RNA polymerase II, F, 55kD (TAF2F) | , | | | + | + | + | + | + | |
| (TBP)-associated factor, RNA polymerase II, I, 28kD (TAF2I) Tax1 (human T-cell leukemia virus type I) binding protein 1 (TAX1BP1) T-box 2 (TBX2) (non-exact 1 U28049 + + + + + + + + + + + + + + + + + + + | (TBP)-associated factor, RNA polymerase II, G, 32kD (TAF2G) | 2 | | | | | | + | | |
| leukemia virus type I) binding protein 1 (TAX1BP1) T-box 2 (TBX2) (non-exact 1 U28049 + + + + + 77%) TBP-associated factor 172 1 AJ001017 + + + + + + + + + + + + + + + + + + + | (TBP)-associated factor, RNA polymerase II, I, 28kD (TAF2I) | 1 | | + | + - | + | + | | + | |
| T-box 2 (TBX2) (non-exact 1 | leukemia virus type I) binding protein 1 | 1 | U33821 | | + | + | + | + | + | |
| (TAF-172) T-cell death-associated 1 U95218 + gene 8 (TDAG8) T-cell leukemia/lymphoma 1 X82240 + 1A (TCL1A) T-cell leukemia/lymphoma 1 X82240 T-cell leukemia/lymphoma 1 X82240 T-cell receptor (delta D2- 1 M22197 T- | T-box 2 (TBX2) (non-exact 77%) | 1 | U28049 | | | + | + | | + | |
| gene 8 (TDAG8) T-cell leukemia/lymphoma 1 X82240 + | TBP-associated factor 172 (TAF-172) | 1 | AJ001017 | | + | | + | | + | |
| 1A (TCL1A) 1-cell leukemia/lymphoma | gene 8 (TDAG8) | | | | | | + | | | |
| 1A (TCL1A) (low match) T-cell receptor (delta D2- | 1A (TCL1A) | | | + | | | | | | |
| | 1A (TCL1A) (low match) | | | | | | | | | |
| | T-cell receptor (delta D2- J1-region) (clone K3B) | 1 | M22197 | | | | | | | |

WO 00/40749

| | | | | | | | | _ | |
|--|--|----------|---------------|--|----------|----------|---|----|---------------------|
| T-cell receptor (V beta 5.1, J beta 1.5, C beta 1) (low | 1 | M97705 | | | | | | | |
| match) T-cell receptor alpha delta (=M94081) | 2 | AE000662 | | + | - | + | | ├- | - |
| T-cell receptor alpha | | B39625 | | | - | ┿ | ┼ | ╄ | |
| enhancer-binding protein, | , | | | | | 1 | ļ | • | |
| short form (=X58636 Mouse LEF1 lymphoid | 1 | | | | 1 | | 1 | | |
| enhancer binding factor 1 | | | | 1 | | 1 | 1 | 1 | |
| (=D16503)) T-cell receptor delta gene | | Hadron | | ļ | <u> </u> | ↓_ | | | |
| D2-J1-region, clone K3B | 1 | M22197 | | | <u> </u> | | | | |
| T-cell receptor germline beta chain gene V-region (V) V-beta-MT1-1 | 1 | M11955 | | | | | | | |
| T-cell receptor germline | | M14159 | + | + | - | - | - | ├ | only in blood |
| beta-chain gene J2.1 exon | | | | | | | | | only in blood |
| T-cell receptor germline delta-chain D-J region | 2 | M22152 | | | | | | | |
| T-cell receptor interacting molecule (TRIM) protein | 2 | AJ224878 | | | | | | + | |
| T-cell receptor rearranged delta-chain, V-region (V- | 1 | M21784 | | | | | | | |
| delta 3-J) T-cell receptor, alpha | 3 | AE000660 | + | + | + | + | - | + | |
| (V,D,J,C) (TCRA) T-cell receptor, beta cluster | 3 | L34740 | + | + | + | + | + | + | high in pancreas |
| (TCRB) T-cell receptor, delta | 2 | X73617 | | | + | + | | + | |
| (V,D,J,C) (TCRD) T-cell, immune regulator 1 | 3 | U45285 | · | ļ | | Ĺ | | Ŀ | |
| (TCIRG1) | | | | | | | | | only found in tumor |
| TCF-1 mRNA for T cell factor 1 | 1 | X59870 | | | | | | | |
| TCF-1 mRNA for T cell factor 1 (splice form B) (low match) | 1 | X59870 | | | | | | | |
| T-COMPLEX PROTEIN 1, | 1 | Q99832 | | | | \vdash | | _ | |
| ETA SUBUNIT (TCP-1- ETA) (CCT-ETA) (HIV-1 | | | | | 1 | 1 | | | |
| NEF INTERACTING PROTEIN) | | | | | | | | | |
| T-COMPLEX PROTEIN 1. | 1 | P50990 | | | | _ | | | |
| THETA SUBUNIT (TCP-1- THETA) (CCT-THETA) | | | | | | | | | |
| (KIAA0002) | 1 | S94421 | | ļ | | <u> </u> | | | |
| receptor(eta-exon) | • | 00+121 | | 1 | İ | | | | |
| TCR V Beta 13.2 | 1 | X75419 | | | | | | | |
| TERA | 1 | AC004472 | | | | | | | |
| testis enhanced gene transcript (TEGT) | 33 | X75861 | + | + | + | + | + | + | |
| tetracycline transporter-like protein (TETRAN) | 2 | L11669 | | + | + | + | | + | |
| tetratricopeptide repeat domain 1 (TTC1) | 1 | U46570 | + | + | + | + | | + | |
| tetratricopeptide repeat domain 2 (TTC2) | 1 | U46571 | | + | | + | | + | |
| tetratricopeptide repeat domain 3 (TTC3) | 1 | D84296 | + | + | + | + | | + | |
| TGFB1-induced anti- | 7 | D86970 | + | + | + | + | | + | |
| apoptotic factor 1 (TIAF1) thioredoxin reductase 1 | 3 | S79851 | - <u></u> | + | + | + | | + | |
| (TXNRD1) THIOREDOXIN- | | P30048 | | | | | | | |
| DEPENDENT PEROXIDE REDUCTASE | • | | | | | | | | |
| PRECURSOR, | | | | | | | | | |
| mitochondrial (ANTI- OXIDANT PROTEIN 1) | | | | | | | | | |
| (AOP-1) | | | | | | | | | |
| | | | | | | | | | |

| VV O 00/40/49 | | | | | | | | | C1/CA00/00005 |
|--|-----|------------|---|---|---|---|---|---|--|
| threonyl-tRNA synthetase (TARS) | 1 | M63180 | | + | + | + | | + | l. |
| thrombin inhibitor | 1 | Z22658 | | | | T | | | |
| thrombospondin 1 (THBS1) | 2 | X04665 | | + | + | + | + | + | · · · · · · · · · · · · · · · · · · · |
| thromboxane A synthase 1 (platelet, cytochrome P450, subfamily V) (TBXAZ1) | 1 | M80647 | | + | | + | + | + | |
| thymidine kinase 2, mitochondrial (TK2) | 2 | X76104 | | + | + | | + | | |
| thymidylate kinase (CDC8) | 1 | L16991 | | + | + | + | | + | |
| thymine-DNA glycosylase (TDG) | 2 | U51166 | + | + | + | + | | + | |
| Thymosin, beta 10 (TMSB10) | 2 | M20259 | + | + | + | + | + | + | |
| thymosin, beta 4, X chromosome (TMSB4X) | 29 | M17733 | | + | + | + | | + | |
| thyroid autoantigen 70kD (Ku antigen) (G22P1) | 7 | J04611 | | | | | | | |
| thyroid hormone receptor coactivating protein (SMAP) | 1 | AF016270 | | + | | + | | + | |
| thyroid hormone receptor interactor 7 (TRIP7) | 2 | L40357 | | + | + | + | | + | |
| thyroid hormone receptor interactor 8r (TRIP8) | 4 | L40411 | | + | | | | | |
| thyroid hormone receptor- associated protein, 230 kDa subunit (TRAP230) | | D83783 | | | | | | | |
| thyroid receptor interacting protein 15 (TRIP15) | 2 | L40388 | + | + | + | + | | | |
| TI-227H | 1 | D50525 | | | | | | | |
| TIA1 cytotoxic granule- associated RNA-binding protein (TIA1) | 1 | M77142 | | + | + | + | | + | |
| tissue inhibitor of metalloproteinase 1 (erythroid potentiating activity, collagenase inhibitor) (TIMP1) | 1 " | X02598 | + | + | + | + | + | + | |
| tissue inhibitor of metalloproteinase 2 (TIMP2) | 7 | M32304 | + | + | + | + | | + | high in placenta |
| tissue specific transplantation antigen P35B (TSTA3) | 1 | U58766 | + | + | + | + | | + | |
| titin (TTN) | 1 | X64697 | + | + | + | + | | + | high in muscle |
| TNF receptor-associated factor 2 (TRAF2) | 1 | U12597 | | + | + | + | | + | |
| TNF receptor-associated factor 3 (TRAF3) | 1 | AF110908.1 | | + | | | | | |
| TNF receptor-associated factor 6 (TRAF6) (low match) | 1 | U78798 | | | | | | | |
| toll-like receptor 1 (TLR1) | 1 | U88540 | | | | + | | | |
| toll-like receptor 2 (TLR2) | 1 | U88878 | + | + | | + | | + | |
| toll-like receptor 4 (TLR4) | 1 | U88880 | | + | | | + | | |
| toll-like receptor 5 (TILR5) | 1 | AF051151 | | + | | + | | | |
| topoisomerase (DNA) I (TOP1) | 1 | J03250 | | + | + | + | | | |
| topoisomerase (DNA) II beta (180kD) (TOP2B) | 2 | X68060 | + | + | + | + | | + | |
| topoisomerase (DNA) III beta (TOP3B) | 3 | D87012 | + | | | | | | |
| TR3beta | 1 | D85245 | | + | | | | | |
| TRAF family member- associated NF-kB activator (TANK) | 3 | U63830 | + | + | + | + | + | + | |
| TRANSALDOLASE | 11 | P37837 | | | | | | | |
| transaldolase 1 (TALDO1) | 4 | L19437 | | + | + | + | + | + | |

97

| transaldolase-related protein | 1 | AF010398 | | Ţ | | Τ | | | | ř. |
|---|----|----------|-------------|---|---|---|---|---|---|-----------------------|
| transcobalamin II (TCII) | 1- | AF047576 | | + | | | - | | | ··- |
| transcription elongation factor B (SIII), polypeptide 1-like (TCEB1L) | 2 | Z47087 | + | + | + | + | | + | | |
| transcription elongation factor B (SIII), polypeptide 3 (110kD, elongin A) (TCEB3) | 1 | L47345 | + | + | + | + | + | + | | |
| transcription factor 12 (HTF4, helix-loop-helix transcription factors 4) (TCF12) | 1 | M83233 | + | + | + | + | | + | | |
| transcription factor 17 (TCF17) | 2 | D89928 | | + | | + | | | - | |
| transcription factor 4 (TCR4) | 2 | X52079 | | + | + | + | | + | | |
| transcription factor 6-like 1 (mitochondrial transcription factor 1-like) (TCF6L1) | 2 | M62810 | + | + | + | + | | | | |
| transcription factor 7-like 2 (T-cell specific, HMG-box) (TCF7L2) | 1 | Y11306 | | + | + | + | | + | | |
| transcription factor binding to IGHM enhancer 3 (TFE3 | 1 | X96717 | + | + | + | + | | + | | |
| transcription factor IL-4 Stat | 7 | AF067575 | + | + | + | + | + | + | | |
| transcription factor IL-4 Stat (low match) | 1 | U16031 | | | | | | | | |
| transcription factor ISGF-3 (=M97936) | 4 | M97935 | | | | | | | | |
| transcription factor REST | 7 | A56138 | | 1 | | | | | | |
| transcription factor TFIID | 1 | Z22828 | ·· | 1 | | | | | | |
| transcriptional adaptor 2 (ADA2, yeast, homolog)- like (TADA2L) | 1 | AF064094 | | | | | | | | |
| transcriptional intermediary factor 1 (TIF1) (non-exact 72%) | 1 | AF009353 | | | | | | | - | |
| transducin (beta)-like 1 (TBL1) | 1 | Y12781 | + | + | + | + | | + | | |
| transducin-like enhancer of split 3, homolog of Drosophila E(sp1) (TLE3) | 1 | M99438 | + | + | | | | | | |
| Transformation/transcription domain-associated protein (TRRAP) | 1 | AF076974 | + | + | + | + | | + | - | - ,,,, |
| transformation-sensitive, similar to Saccharomyces cerevisiae STI1 (STI1L) | 2 | M86752 | | + | + | + | | + | | |
| transforming growth factor beta-activated kinase 1 (TAK1) (non-exact 78%) | 7 | AB009356 | | | | | | | | |
| transforming growth factor beta-stimulated protein TSC-22 (TSC22) | 3 | AJ222700 | + | + | + | + | | + | | |
| transforming growth factor, beta receptor III (betaglycan, 300kD) (TGFBR3) | 1 | L07594 | | + | + | + | | + | | |
| transforming growth factor, beta-induced, 68kD (TGFBI) | 2 | 4507466 | + | + | + | + | + | + | | |
| TRANSFORMING GROWTH FACTOR-BETA INDUCED PROTEIN IG-H3 PRECURSOR (BETA IG- H3) | 2 | Q15582 | | | | | | | | |
| transforming, acidic coiled- coil containing protein 1 (TACC1) (non-exact 70%) | 1 | AF049910 | | | | | | | | |

| | | | _ | | | | | | |
|--|----|----------|---------------------------------------|--------------|---|--|---|----------|------------------------|
| transgelin 2 (TAGLN2) | 14 | D21261 | + | + | + | + | + | + | ٠, |
| transgelin 2 (TAGLN2) (non-exact) | 1 | D21261 | · · · · · · · · · · · · · · · · · · · | | | | | | |
| trans-Golgi network protein (46, 48, 51kD isoforms) (TGN51) | 2 | AF029316 | | + | | + | | | |
| transient receptor potential channel 1 (TRPC1) | 1 | X89066 | | + | + | + | | + | |
| transketolase (Wernicke- Korsakoff syndrome) (TKT) | 7 | L12711 | | + | + | + | | + | |
| translation factor sui1 homolog (GC20) | 1 | AF064607 | | + | + | + | + | + | |
| translin (TSN) | 3 | X78627 | + | + | + | + | | + | |
| translin-associated factor X (TSNAX) | 1 | X95073 | | + | + | + | | + | |
| transmembrane glycoprotein (A33) | 1 | U79725 | | | | | | | |
| transmembrane protein (63kD), endoplasmic reticulum/Golgi intermediate compartment (P63) | 1 | X69910 | + | + | + | + | | + | |
| transmembrane protein 1 (TMEM2) | 1 | AB001523 | | + | | + | | + | |
| TRANSMEMBRANE PROTEIN SEX PRECURSOR (non-exact 65%) | 1 | P51805 | | | | | | | |
| transmembrane trafficking protein (TMP21) | 2 | X97442 | + | + | + | + | + | + | |
| transporter 1, ABC (ATP binding cassette) (TAP1) | 3 | L21208 | + | + | + | + | | + | |
| Treacher Collins- Franceschetti syndrome 1 (TCOF1) | 2 | U40847 | + | + | + | + | | + | high in many libraries |
| triosephosphate isomerase 1 (TPI1) | 2 | X69723 | + | + | + | + | + | + | |
| tropomyosin | 2 | X04201 | | + | + | + | | + | |
| tropomyosin 4 (TPM4) | 2 | X05276 | + | + | + | + | | + | |
| TRPM-2 protein | 2 | M63376 | | | | | t | \vdash | |
| tryptase I precursor (non- exact 64%)(=P20231) | 1 | A35863 | | 1 | | | | | |
| tryptophan rich basic protein (WRB) | 1 | Y12478 | | | | | | | |
| tryptophanyl-tRNA synthetase (WARS) | 1 | X59892 | + | + | + | + | + | + | |
| Ts translation elongation factor, mitochondrial (TSFM) | 1 | L37936 | + | + | | + | | + | |
| ttopoisomerase (DNA) II beta (180kD) | 1 | Z15115 | | + | + | | | + | |
| Tu translation elongation factor, mitochondrial (TUFM) | 4 | L38995 | | | | | | | |
| fuberous sclerosis 1 (TSC1) | 1 | AF013168 | | + | + | + | | + | |
| tuberous scierosis 2 (TSC2) | 1 | X75621 | | + | + | + | | + | |
| tubulin, alpha 1 (testis specific) (TUBA1) | 1 | X06956 | | + | | | + | | |
| tubulin, alpha, ubiquitous (K-ALPHA-1) | 11 | K00558 | + | + | + | + | + | + | high in many libraries |
| tubulin, alpha, ubiquitous (K-ALPHA-1) (low match) | 1 | K00558 | | | | | | | |
| tubulin-specific chaperone c (TBCC) | 1 | U61234 | | + | + | + | | + | |
| tumor necrosis factor (ligand) superfamily, member 10 (TNFSF10) | 7 | U37518 | | + | + | + | | + | |

| WU 00/40/49 | | | | | | | | | C1/CA00/00003 |
|--|--------------------|----------|---------------------------------------|---|----------|---|---|---|------------------------|
| tumor necrosis factor (ligand) superfamily, | 1- | AF046888 | + | + | | + | | + | |
| member 13 (TNFSF13) tumor necrosis factor (ligand) superfamily. | 1 | AF036581 | | | | _ | | | |
| member 14 (TNFSF14) | | D38122 | + | - | | | | | Found only in library |
| (ligand) superfamily, member 6 (TNFSF6) | | | | | | | | | 386: T-cell lymphoma |
| tumor necrosis factor (ligand) superfamily, member 8 (TNFSF8) | 1 | L09753 | B only | | | | | | |
| tumor necrosis factor alpha-inducible cellular protein containing leucine | 7 | AF061034 | | + | + | + | | + | |
| zipper domains (FIP2) Tumor necrosis factor | 2 | M63928 | | + | | | + | | |
| receptor superfamily member 7 (TNFRSF7) | | | <u></u> | | <u> </u> | | | | |
| tumor necrosis factor receptor superfamily, member 10b (TNFRSF10B) | 1 | AF016266 | | + | + | + | + | + | |
| tumor necrosis factor receptor superfamily, member 10c, decoy without an intracellular domain (TNFRSF10C) | 3 | AF012629 | | | | | + | | |
| tumor necrosis factor receptor superfamily, member 10d, decoy with truncated death domain (TNFRSF10D) | 1 | AF023849 | | | | | | | found only in prostate |
| (non-exact 84%) tumor necrosis factor receptor superfamily, member 12 (translocating chain-association membrane protein) | 7 | U94508 | + | + | + | + | | + | |
| (TNFRSF12) tumor necrosis factor receptor superfamily, member 14 (herpesvirus entry mediator) (TNFRSF14) | T | U70321 | + | + | + | + | | + | · |
| tumor necrosis factor receptor superfamily, member 1B (TNFRSF1B) | 5 | U52165 | + | + | + | + | | + | |
| tumor necrosis factor receptor superfamily, member 6 (TNFRSF6) | 1 · · · | X63717 | B, W | | | | | + | |
| tumor necrosis factor receptor superfamily, member 7 (TNFRSF7) | 1 | M63928 | + | + | | | | | |
| tumor necrosis factor, alpha-induced protein 2 (TNFAIP2) | 8 | M92357 | | + | + | | + | | |
| tumor necrosis factor, alpha-induced protein 3 (TNFAIP3) | 2 | M59465 | | | | | | | |
| tumor protein 53-binding protein, 1 (TP53BP1) | 1 | AF078776 | | + | + | + | | + | |
| tumor protein p53 (Li- Fraumeni syndrome) (TP53) | 1 | M14695 | + | + | | | | + | |
| Tumor protein p53-binding protein (TP53BPL) | 1 | U82939 | + | | | + | | + | |
| tumor protein, translationally-controlled 1 (TPT1) | 35 | X16064 | | | | | | | |
| tumor protein, translationally-controlled 1 (TPT1) (low score) | 1 | X16064 | · · · · · · · · · · · · · · · · · · · | | | | | | |
| tumor rejection antigen (gp96) 1 (TRA1) | 9 | X15187 | +. | + | + | + | + | + | |

| | | | _ | | | | | | · |
|--|---|----------|---|--|--|--------------|----------|---|---------------------|
| lumorous imaginal discs (Drosophila) homolog (TID1) | 2 | AF081749 | | + | | | | | |
| TXK tyrosine kinase (TXK) | 2 | L27071 | | - | | \vdash | ┼~ | + | |
| type II integral membrane protein (NKG2-E) | 1 | AJ001685 | | | | | + | 1 | found only in fetal |
| TYRO protein tyrosine kinase binding protein (TYROBP) | 3 | AF019562 | | | + | | | | inverspicen |
| tyrosine 3- monooxygenase/tryptopha in 5-monooxygenase activation protein, beta polypeptide (YWHAB) | 1 | X57346 | + | + | + | + | | + | high in ecnorm |
| tyrosine 3- monooxygenase/tryptopha n 5-monooxygenase | 1 | M86400 | | | | | | | |
| activation protein, zeta polypeptide (YWHAZ) | | | | | | | | | |
| tyrosine 3- monooxygenase/tryptopha n 5-monooxygenase activation protein, zeta polypeptide (YWHAZ) | 1 | M86400 | | | | | | | |
| Tyrosine kinase 2 (TYK2) | 3 | X54637 | | + | + | + | | + | |
| TYROSINE-PROTEIN KINASE ZAP-70 (70 KD ZETA-ASSOCIATED PROTEIN) (SYK-RELATED TYROSINE KINASE) | 2 | P43403 | | | | | | | |
| tyrosyFtRNA synthetase (YARS) | 7 | U89436 | + | + | + | + | | + | |
| Ù1 smáll nuclear RNA | 1 | M14387 | | 1 | | | | 1 | |
| U19H snoRNA (=M63485 R.norvegicus matrin 3) | 1 | AJ224166 | | | | \vdash | | | |
| U2(RNU2) small nuclear RNA auxillary factor 1 (non-standard symbol) (U2AF1) | 1 | M96982 | | + | + | + | | + | |
| U22 snóRNA host gene (UHG) | 2 | U40580 | | | | | | | |
| U4/U6-associated RNA splicing factor (HPRP3P) | 4 | AF016370 | | + | + | + | <u> </u> | + | |
| U49 small nuclear RNA | 1 | X96649 | | + | | - | ┢ | - | |
| U5 snRNP-specific protein (220 kD), ortholog of S. cerevisiae Prp8p (PRP8) | 1 | AB007510 | + | + | + | + | | + | |
| U5 snRNP-specific protein, 116 kD (U5-116KD) | 4 | D21163 | + | + | + | + | | + | |
| U5 snRNP-specific protein, 200 kDa (DEXH RNA helicase family) (U5-200- KD) | 3 | 270200 | | | | | | | |
| Uba80 mRNA for ubiquitin | 4 | S79522 | + | + | + | + | + | + | high in ovary |
| ubiquinol-cytochrome c reductase (6.4kD) subunit (UQCR) | 1 | D55636 | + | + | + | + | + | + | high in fetal lung |
| ÜBIQUÍNOL- CYTOCHROME C REDUCTASE IRON- SULFUR SUBUNIT PRECURSOR (RIESKE IRON-SULFUR PROTEIN) (RISP) (low match) | 1 | P47985 | | | | | | | |
| ubiquitin A-52 residue ribosomal protein fusion product 1 (UBA52) | 2 | X56999 | | | | | | | |
| ubiquitin activating enzyme E1-like protein (GSA7) | 1 | AF094516 | | + | + | | | + | |
| ubiquitin C (UBC) | 5 | AB009010 | | + | + | + | + | + | high in ovary |

| | | | _ | | | | | | |
|---|---|----------|---|---|---|---|---|----------------|---|
| ubiquitin carboxyl-terminal esterase L3 (ubiquitin thiolesterase) (UCHL3) | 1 | M30496 | + | + | + | + | | + | |
| ubiquitin fusion degradation 1-like (UFD1L) | 1 | U64444 | + | + | + | + | | + | |
| ubiquitin protein ligase E3A (human papilloma virus E6- associated protein, | 1 | U84404 | В | + | + | | | + | |
| Angelman syndrome) (UBE3A) | | | | | | | | | |
| ubiquitin specific protease 10 (USP10) | 4 | D80012 | + | + | + | + | | + | |
| ubiquitin specific protease 11 (USP11) | 1 | U44839 | + | + | + | + | + | + | |
| ubiquitin specific protease 15 (USP15) | 3 | AB011101 | + | + | + | + | | L ⁺ | |
| ubiquitin specific protease 19 (USP19) ubiquitin specific protease | 1 | AB020698 | | + | | | | | |
| 4 (proto-oncogene) (USP4) ubiquitin specific protease | 1 | AF017305 | В | + | + | | + | + | |
| 4 (proto-oncogene) (USP4) (non-exact, 66%) | 1 | AF017306 | | | | | | | |
| ubiquitin specific protease 7 (herpes virus-associated) (USP7) | | Z72499 | | + | + | + | | + | |
| ubiquitin specific protease 8 (USP8) | 5 | D29956 | | + | + | + | | + | · |
| UBIQUITIN-ACTIVATING ENZYME E1 (A1S9 PROTEIN) (56%) | 1 | P22314 | | | | | | | |
| ubiquitin-activating enzyme E1 (A1S9T and BN75 temperature sensitivity complementing) (UBE1) | | M58028 | + | + | + | + | | + | |
| ubiquitin-activating enzyme E1, like (UBE1L) | 1 | L34170 | + | + | | + | | + | |
| UBIQUITIN-BINDING PROTEIN P62; phosphotyrosine independent ligand for the Lck SH2 domain p62 (P62) | 1 | U41806 | | | + | | + | | |
| ubiquitin-conjugating enzyme E2 variant 1 (UBE2V1) | 2 | U49278 | + | + | + | + | + | + | |
| ubiquitin-conjugating enzyme E2 variant 2 (UBE2V2) | 1 | X98091 | | | | | | | |
| UBIQUITIN- CONJUGATING ENZYME E2-17 KD (UBIQUITIN- PROTEIN LIGASE) | 1 | Q16781 | | | | | | | |
| ubiquitin-conjugating enzyme E2B (RAD6 homolog) (UBE2B) | f | M74525 | + | + | + | + | | + | |
| ubiquitin-conjugating enzyme E2G 2 (homologous to yeast UBC7) (UBE2G2) | 1 | AF032456 | + | + | + | + | | + | |
| ubiquitin-conjugating enzyme E2H (homologous to yeast UBC8) (UBE2H) | 1 | Z29328 | + | + | + | + | | + | |
| ubiquitin-conjugating enzyme E2L 1 (UBE2L1) | 1 | X92962 | | + | + | | | + | |
| ubiquitin-conjugating enzyme E2L 3 (UBE2L3) | 3 | AJ000519 | | + | + | + | | + | |
| ubiquitin-conjugating enzyme E2L 6 (UBE2L6) | 4 | AF031141 | | + | + | + | + | + | |
| ubiquitin-like 1 (sentrin) (UBL1) | 2 | U61397 | + | + | + | + | | + | |

| | | | | | | | | • | C1/CA00/00003 |
|--|----------|----------|--------------|----------|-------------|----|---|---|---------------|
| UDP-N-acetyl-alpha-D- galactosamine:polypeptide N- | 2 | X85019 | | | | | | | |
| acetylgalactosaminyltransf lerase 2 (GalNAc-T2) (GALNT2) | | | | | | | | | |
| UDP-N-acetyl-alpha-D- | 1-1- | X92689 | | | | ├- | ऻ | - | |
| galactosamine:polypeptide | | | | ŀ | | | | | |
| acetylgalactosaminyltransf erase 3 | | | | | | | } | | |
| (GalNAc-T3) (GALNT3) (non-exact 65%) | | · | | ļ | | | | | |
| unactive progesterone receptor, 23 Kd (P23) | 2 | L24804 | | + | + | + | | + | |
| unconventional myosin-ID (MYO1F) | 3 | U57053 | | | | | | | |
| uncoupling protein homolog (UCPH) | 1 | U94592 | | | | | | | |
| uncoupling protein homolog (UCPH) (low | 1 | U94592 | | | | | | | |
| match 67%) Unknown gene product | 1 | AC002310 | | . | | ļ | _ | | |
| unknown mRNA (clone | <u> </u> | AF070542 | | - | <u> </u> | ├ | _ | _ | |
| 24514) unknown protein (clone | 2 | | | | | | | | |
| ICRFp507L0677) | | Z70223 | | | | | | | |
| unknown protein (Hs.93832) | 1 | AF070626 | + | + | + | + | + | + | |
| unknown protein IT14 | 1 | AF040966 | | | | | | | |
| (S.cerevisiae) 6 homolog | 1 | D79984 | + | + | + | + | + | + | |
| upregulated by 1,25- dihydroxyvitamin D-3 (VDUP1) | 74 | 573591 | + | + | + | + | | + | high in heart |
| upregulated by 1,25- dihydroxyvitamin D-3 (VDUP1) (low match) | 1 | S73591 | | | | | | | |
| upregulated by 1,25- dihydroxyvitamin D-3 (VDUP1) (low match) | 1 | S73591 | | | | | ' | | |
| upregulated by 1,25- dihydroxyvitamin D-3 (VDUP1) (low score) | 1 | S73591 | | | | | | | |
| upstream binding factor (hUBF) | 1 | X53461 | + | + | | + | | + | |
| UV radiation resistance associated gene (UVRAG) | 2 | X99050 | | + | + | + | | + | |
| vacuolar proton-ATPase, subunit D; V-ATPase, subunit D (ATP6DV) | 4 | X71490 | | + | + | + | + | + | |
| v-akt murine thymoma viral oncogene homolog 1 | 1 | M63167 | + | + | + | + | | + | |
| (AKT1) Vanin 2 (VNN2) | 3 | AJ132100 | | + | | _ | | | |
| vasodilator-stimulated phosphoprotein (VASP) | 3 | Z46389 | + | | + | + | | + | |
| vav 1 oncogene (VAV1) | 1 | M59834 | | † | | | | + | |
| vav 2 oncogene (VAV2) | 1 | S76992 | + | + | | | | | |
| v-crk avian sarcoma virus CT10 oncogene homolog (CRK) | 1 | D10656 | W | + | + | | + | | |
| v-erb-b2 avian erythroblastic leukemia viral oncogene homolog 3 (ERBB3) | 1 | M29366 | | | | | | + | |
| VERSICAN CORE PROTEIN PRECURSOR | 1 | P13611 | | | | | | | |
| Vesicle-associated membrane protein 1 (synaptobrevin 1) (VAMP1) | 1 | M36196 | | + | + | + | | + | |

| vesicle-associated membrane protein 3 (cellubrevin) (VAMP3) | 1 | U64520 | | | | | | | , |
|--|----|----------|---|--------------|---|---|----------|---|------------------------|
| v-fos FBJ murine osteosarcoma viral oncogene homolog (FOS) | 26 | K00650 | | + | + | + | + | + | high in aorta |
| v-fos FBJ murine osteosarcoma viral oncogene homolog (FOS) (low match) | 1 | K00650 | | | | | | | |
| villin 2 (ezrin) (VIL2) | 1 | X51521 | + | + | + | + | | + | |
| villin-like protein | 1 | D88154 | | | | | | T | |
| vimentin (VIM) | 12 | X56134 | | + | + | + | + | + | high in many libraries |
| vinculin (VCL) | 4 | M33308 | | + | + | + | † | + | |
| vitamin A responsive; cytoskeleton related (JWA) | 6 | AF070523 | | + | + | + | | + | |
| v-jun avian sarcoma virus 17 oncogene homolog (JUN) | 2 | U65928 | + | + | + | + | | + | |
| v-myb avian myeloblastosis viral oncogene homolog (MYB) | | M15024 | | | + | | + | | |
| voltage-dependent anion channel 1 (VDAC1) | 1 | L06132 | + | + | + | + | | + | |
| voltage-dependent anion channel 3 (VDAC3) | 4 | U90943 | | + | + | + | | + | |
| von Hippel-Lindau syndrome (VHL) | 1 | L15409 | | + | + | + | | + | |
| von Willebrand factor (vWF) (low matched) | 1 | X06828 | | | | | | | |
| v-raf murine sarcoma 3611 viral oncogene homolog 1 (ARAF1) | 2 | L24038 | + | + | + | + | | | |
| v-rat-1 murine leukemia viral oncogene homolog 1 (RAF1) | 1 | X03484 | + | + | + | + | | + | |
| v-ral simian leukemia viral oncogene homolog B (ras related; GTP binding protein) (RALB) | 3 | M35416 | | | | | | | |
| V-rel avian reticuloendotheliosis viral oncogene homolog A (nuclear factor of kappa light polypeptide gene enhancer in B-cells 3 (p65)) (RELA) | 1 | L19067 | | + | + | + | | + | , |
| v-yes-1 Yamaguchi sarcoma viral related oncogene homolog (LYN) | 2 | M16038 | + | + | | + | | + | |
| WD repeat domain 1 (WDR1) | 1 | AB010427 | + | + | + | + | + | + | |
| WDR1 (=AF020260) | 1 | AF020056 | | 11 | | | | | |
| WD-repeat protein (HAN11) | 2 | U94747 | | + | + | | | + | |
| Williams-Beuren syndrome chromosome region 1 (WBSCR1) | 12 | AF045555 | + | + | + | + | + | + | |
| Wiskott-Aldrich syndrome protein interacting protein (WASPIP) | 4 | X86019 | + | + | + | | | + | |
| X (inactive)-specific transcript (XIST) | 2 | M97168 | | | | | | | |
| xeroderma pigmentosum, complementation group C (XPC) | 3 | D21089 | + | + | + | + | | | |
| XIAP associated factor-1 | 2 | X99699 | | \vdash | | + | | | |
| XIB | 1 | X90392 | | + | + | | + | + | |
| X-linked anhidroitic | 1 | AF003528 | | \vdash | | | | - | |
| ectodermal dysplasia | | | | | | | | | |

| TV | | LIBBORD | | | | | , | , | |
|---|-----------|------------|--------------|--|----------------|-------------------|--------------|--|------------------|
| X-ray repair complementing defective | 1 | M30938 | + | + | + | + | | + | high in spleen |
| repair in Chinese hamster | | | | | | ŀ | 1 | | |
| cells 5 (double-strand- | | | | | 1 | | İ | 1 | |
| break | · | | | Į. | ĺ | 1 | | | |
| rejoining; Ku autoantigen, | | | | | | 1 | | | 1 |
| 80kD) (XRCC5) | | | | | į . | İ |] | i | |
| XRP2 protein | 1 | AJ007590 | | + | - | +- | ┼ | + | |
| • | l | 1 | | | ⊢ | <u> </u> | ↓ | ↓_ | |
| yeloid differentiation primary response gene | 1 | U84408 | | + | + | + | j | + | 1 |
| (88) (MYD88) | | | | 1 | } | 1 | | | |
| zeta-chain (TCR) | | L05148 | + - | | | + | | ├— | |
| associated protein kinase | ' | L03140 | , , | 1 | | * | | | l |
| (70kD) (ZAP70) | | ļ | ì | Ì | Į | 1 | 1 | 1 | İ |
| zeta-chain (TCR) | | L05148 | | | ├ | ┼ | - | ╁ | |
| associated protein kinase | , | 2001.40 | | | 1 | ļ | | | |
| (70kD) (ZAP70) (low | | ì | 1 | | ŀ | | 1 | 1 | |
| match) | | ļ | 1 | i | l | i | ĺ | ı | |
| zinc finger protein | 2 | U69274 | + | + | + | + | 1 | + | |
| (Hs.47371) | | } | | 1 | i | 1 | l | 1 | 1 |
| zinc finger protein | 1 | U69645 | + | + | + | + | | + | |
| (Hs.78765) | | 1 | 1 | 1 | ŀ | 1 | ı | 1 | i |
| zinc finger protein 10 (KOX | 1 | X78933 | | | | | | Ī | + only |
| 1) (ZNF10) | | | <u> </u> | | L | | L | L | |
| ZINC FINGER PROTEIN | 1 | Q15973 | | | | | | | |
| 124 (HZF-16) (non-exact | | | | İ | l | | | | 1 |
| 51%) | | DE VOLUME | | \bot | | $ldsymbol{f eta}$ | <u></u> | <u> </u> | |
| zinc finger protein 124 | 1 | S54641 | | | 1 | | | | |
| (HZF-16) (ZNF124) (non- exact, 78%) | | | ! | 1 | | | l | ļ | |
| ZINC FINGER PROTEIN | 1 | P52736 | | | | <u> </u> | <u> </u> | <u> </u> | |
| 1133 | ' | P32/30 | | 1 | l | | l | 1 | |
| zinc finger protein 136 | 1 | U09367 | ļ | | + | + | ├ | <u> </u> | |
| (clone pHZ-20) (ZNF136) | ' | 009307 | l | ł | T | - | l | 1 | |
| zinc finger protein 140 | 1 | U09368 | | + | | + | ├— | + | |
| (clone pHZ-39) (ZNF140) | • | 003300 | ļ | 1 | | " | 1 | T | |
| zinc finger protein 140 | 1 | AF060865 | | 1 | \vdash | ┼ | ├── | ├ | |
| (clone pHZ-39) (ZNF140) | • | / " 00000 | | İ | i | | 1 | Į | 1 |
| (non-exact 59%) | | | | | | 1 | i | l | 1 |
| zinc finger protein 140 | 1 | U09368 | | | | | 1 | - | |
| (clone pHZ-39) (ZNF140) | | | | Ì | | Ì | | l | |
| (non-exact 73%) | | | | ł | | | | 1 | |
| zinc finger protein 140 | 1 | S66508 | | T | | | | | |
| (clone pHZ-39) (ZNF140) (non-exact 73%aa) | | | | | 1 | 1 | | | 1 |
| (non-exact 73%aa) | | | | <u> </u> | | <u>L</u> . | | | |
| zinc finger protein 140 | 1 | U09368 | | | | | | | |
| (clone pHZ-39) (ZNF140) | " | | | | | | 1 | | ! |
| (non-exact, 80%) | | TIBBBEB | | 1 | <u> </u> | Ļ. | <u> </u> | ــــــــــــــــــــــــــــــــــــــ | |
| zinc finger protein 143 (clone pHZ-1) (ZNF143) | 2 | U09850 | + | + | + | + | + | + | |
| zinc finger protein 143 | | LINNOEN | | - | | . | ┞— | | |
| (clone pHZ-1) (ZNF143) | 1 | U09850 | | | } | ł | | | i l |
| (low match) | | | | 1 | i | ł | l | | ĺ |
| zinc finger protein 148 | 1 | AF039019 | + | + | | ├ | | - | |
| (pHZ-52) (ZNF148) | • | / 000010 | · | 1 | l | 1 | ŀ | | · |
| ZINC FINGER PROTEIN | 1 | Q13105 | | + | | - | ┢─ | | |
| 151 (MIZ-1 PROTEIN) (low | · | } | | 1 | l | | 1 | | |
| match) | | | | 1 | | 1 | l | | |
| zinc finger protein 173 | 1 | U09825 | B, T | + | + | | + | | |
| (ZNF173) | | l | ' | 1 | | 1 | | 1 | |
| zinc finger protein 192 | 1 | U57796 | | 1 | | | T | | |
| (ZNF192) (non-exact, 66%) | | <u> </u> | | L | | L | l . | L | |
| zinc finger protein 198 | 1 | AJ224901 | | + | + | + | | | |
| (ZNF198) | | | | <u> </u> | | | | <u>L</u> | |
| zinc finger protein 2 (ZNF2) | 1 | X60152 | | | | | | | |
| (low match) | | | | 1 | | | <u> </u> | | |
| zinc finger protein 200 | 1 | AF060866 | | + | | + | | | |
| (ZNF200) | | AEN ANNE | | | | <u> </u> | ـــا | <u> </u> | |
| zinc finger protein 207 (ZNF207) | 6 | AF046001 | + | + | + | + | + | + | high in prostate |
| zinc finger protein 216 | 2 | AENEODTO - | | + | <u> </u> | ├ | <u> </u> | | |
| (ZNF216) | 2 | AF062072 | + | + | + | + | : | + | |
| (4141 210) | L <u></u> | i | | | L | Ц | L | L | L) |

| zinc finger protein 217 | 1 1 | AF041259 | Lacti | vated | | T | Т- | 1 + | |
|---|-------------|--|--|--------------|---|-----------|--|----------|----------------------|
| (ZNF217) ZINC FINGER PROTEIN | | | | | | | <u> </u> | L | |
| 22 (ZINC FINGER | 1 | P17026 | - | | | | | | |
| PROTEIN KOX15) (non- exact 58%) | | | | | | | | | |
| zinc finger protein 230 (ZNF230) | 1 | U95044 | | + | | \dagger | | \vdash | |
| Zinc finger protein 239 | 1 | L26914 | | + | - | + | \vdash | ╁ | |
| (ANF239) zinc finger protein 261 | 1-1- | AB002383 | | ++ | + | + | | + | |
| (ZNF261) zinc finger protein 262 | 1 | AB007885 | | <u> </u> | | | | | |
| (ANF262) | | | | + | | + | | + | |
| zinc finger protein 263 (ZNF263) | 1 | D88827 | <u> </u> | | | | | | |
| zinc finger protein 264 (ZNF264) | 1 | AB007872 | | + | + | + | | | |
| ZINC FINGER PROTEIN 33A (ZINC FINGER | 1 | Q06730 | | | | | | † | |
| PROTEIN KOX31) | | | 1 | | | | | | |
| (KIAA0065) (HA0946) zinc finger protein 42 | 1 | M58297 | + | + | + | + | - | + | |
| (myeloid-specific retinoic cid- responsive) (ZNF42) | | | | | | | | | |
| zinc finger protein 43 (HTF6) (ZNF43) (low | 1 | X59244 | | | - | 1 | | _ | |
| match) | | | | | | | | ĺ | |
| zinc finger protein 43 (HTF6) (ZNF43) (non- | 1 | X59244 | | | | | | | |
| exact, 54%) zinc finger protein 43 | 1 | X59244 | <u> </u> | | | | | | |
| (HTF6) (ZNF43) (non- exact, 71%) | ' | A35244 | | | | | | | i i |
| ZINC FINGER PROTEIN 43 (ZINC PROTEIN HTF6) | 1 | P28160 | | | | | \vdash | | |
| (non-exact 67%) | | | | | | | | | |
| Time times are 1777 | | | | | | | _ | | |
| zinc finger protein 45 (a Kruppel-associated box | 1 | L75847 | | | | | | | only found in testis |
| Kruppel-associated box (KRAB) domain | 1 | L75847 | | | | | | | only found in testis |
| Kruppel-associated box (KRAB) domain polypeptide) (ZNF45) ZINC FINGER PROTEIN | 1 | L75847 | | | | | | | |
| Kruppel-associated box (KRAB) domain polypeptide) (ZNF45) ZINC FINGER PROTEIN 46 (ZINC FINGER PROTEIN KUP) (non-exact | | | | | v | | | | |
| Kruppel-associated box (KRAB) domain polypeptide) (ZNF45) ZINC FINGER PROTEIN 46 (ZINC FINGER PROTEIN KUP) (non-exact 62%) | 1 | P24278 | | | | | | | |
| Kruppel-associated box (KRAB) domain polypeptide) (ZNF45) ZINC FINGER PROTEIN 46 (ZINC FINGER PROTEIN KUP) (non-exact 62%) zinc finger protein 6 (CMPX1) (ZNF6) | 1 | P24278 X56465 | | + | + | + | | + | |
| Kruppel-associated box (KRAB) domain polypeptide) (ZNF45) ZINC FINGER PROTEIN 46 (ZINC FINGER PROTEIN KUP) (non-exact 62%) Zinc finger protein 6 (CMPX1) (ZNF6) Zinc finger protein 74 (Cos52) (ZNF74) (non- | 1 | P24278 | | + | + | + | | + | |
| Kruppel-associated box (KRAB) domain polypeptide) (ZNF45) ZINC FINGER PROTEIN 46 (ZINC FINGER PROTEIN 46 (ZINC FINGER PROTEIN KUP) (non-exact 62%) Zinc finger protein 6 (CMPX1) (ZNF6) Zinc finger protein 74 (Cos52) (ZNF74) (non-exact, 67%) Zinc finger protein 76 | 1 | P24278 X56465 | | + | + | + | | + | |
| Kruppel-associated box (KRAB) domain polypeptide) (ZNF45) ZINC FINGER PROTEIN 46 (ZINC FINGER PROTEIN KUP) (non-exact 62%) Zinc finger protein 6 (CMPX1) (ZNF6) Zinc finger protein 74 (Cos52) (ZNF74) (non-exact, 67%) Zinc finger protein 76 (expressed in testis) (ZNF76) | 1 | P24278 X56465 X71623 | | | | | | | |
| Kruppel-associated box (KRAB) domain polypeptide) (ZNF45) ZINC FINGER PROTEIN 46 (ZINC FINGER PROTEIN 46 (ZINC FINGER PROTEIN KUP) (non-exact 62%) Zinc finger protein 6 (CMPX1) (ZNF6) Zinc finger protein 74 (Cos52) (ZNF74) (non-exact, 67%) Zinc finger protein 76 (expressed in testis) (ZNF76) ZINC FINGER PROTEIN 83 (ZINC FINGER | 1 | P24278 X56465 X71623 | | | | | | | |
| Kruppel-associated box (KRAB) domain polypeptide) (ZNF45) ZINC FINGER PROTEIN 46 (ZINC FINGER PROTEIN 46 (ZINC FINGER PROTEIN KUP) (non-exact 62%) Zinc finger protein 6 (CMPX1) (ZNF6) Zinc finger protein 74 (Cos52) (ZNF74) (non-exact, 67%) Zinc finger protein 76 (expressed in testis) (ZNF76) ZINC FINGER PROTEIN 83 (ZINC FINGER PROTEIN HPF1) (non- | 1 1 | X56465 X71623 M91592 | | | | | | | |
| Kruppel-associated box (KRAB) domain polypeptide) (ZNF45) ZINC FINGER PROTEIN 46 (ZINC FINGER PROTEIN KUP) (non-exact 62%) Zinc finger protein 6 (CMPX1) (ZNF6) Zinc finger protein 74 (Cos52) (ZNF74) (non-exact, 67%) Zinc finger protein 76 (expressed in testis) (ZNF76) ZINC FINGER PROTEIN 83 (ZINC FINGER PROTEIN HPF1) (non-exact 65%) Zinc finger protein 84 | 1 1 | X56465 X71623 M91592 | T activated | | | | | | |
| Kruppel-associated box (KRAB) domain polypeptide) (ZNF45) ZINC FINGER PROTEIN 46 (ZINC FINGER PROTEIN 46 (ZINC FINGER PROTEIN KUP) (non-exact 62%) Zinc finger protein 6 (CMPX1) (ZNF6) Zinc finger protein 74 (Cos52) (ZNF74) (non-exact, 67%) Zinc finger protein 76 (expressed in testis) (ZNF76) ZINC FINGER PROTEIN 83 (ZINC FINGER PROTEIN 83 (ZINC FINGER PROTEIN HPF1) (non-exact 65%) Zinc finger protein 84 (HPF2) (ZNF84) Zinc finger protein 85 | 1 1 | X56465 X71623 M91592 P51522 | T activated | + | + | | | + | |
| Kruppel-associated box (KRAB) domain polypeptide) (ZNF45) ZINC FINGER PROTEIN 46 (ZINC FINGER PROTEIN 46 (ZINC FINGER PROTEIN KUP) (non-exact 62%) Zinc finger protein 6 (CMPX1) (ZNF6) Zinc finger protein 74 (Cos52) (ZNF74) (non-exact, 67%) Zinc finger protein 76 (expressed in testis) (ZNF76) ZINC FINGER PROTEIN 83 (ZINC FINGER PROTEIN HPF1) (non-exact 65%) Zinc finger protein 84 (HPF2) (ZNF84) | 1 1 | P24278 X56465 X71623 M91592 P51522 M27878 | Tactivated | + | + | + | + | + | |
| Kruppel-associated box (KRAB) domain polypeptide) (ZNF45) ZINC FINGER PROTEIN 46 (ZINC FINGER PROTEIN 46 (ZINC FINGER PROTEIN KUP) (non-exact 62%) Zinc finger protein 6 (CMPX1) (ZNF6) Zinc finger protein 74 (Cos52) (ZNF74) (non-exact, 67%) Zinc finger protein 76 (expressed in testis) (ZNF76) ZINC FINGER PROTEIN 83 (ZINC FINGER PROTEIN HPF1) (non-exact 65%) Zinc finger protein 84 (HPF2) (ZNF84) Zinc finger protein 85 (ZNF85)) Zinc finger protein 9 (ZNF9) ZINC FINGER PROTEIN | 1 1 1 2 | P24278 X56465 X71623 M91592 P51522 M27878 U35376 | T activated | + | + | + | + | + | |
| Kruppel-associated box (KRAB) domain polypeptide) (ZNF45) ZINC FINGER PROTEIN 46 (ZINC FINGER PROTEIN KUP) (non-exact 62%) Zinc finger protein 6 (CMPX1) (ZNF6) Zinc finger protein 74 (Cos52) (ZNF74) (non-exact, 67%) Zinc finger protein 76 (expressed in testis) (ZNF76) ZINC FINGER PROTEIN 83 (ZINC FINGER PROTEIN HPF1) (non-exact 65%) Zinc finger protein 84 (HPF2) (ZNF84) Zinc finger protein 85 (ZNF85)) Zinc finger protein 9 (ZNF9) ZINC FINGER PROTEIN 93 (=ZINC FINGER PROTEIN 93 (=ZINC FINGER PROTEIN 93 (=ZINC FINGER PROTEIN 97 (=ZINC FINGER | 1 1 1 2 5 | P24278 X56465 X71623 M91592 P51522 M27878 U35376 M28372 | Tactivated | + | + | + | + | + | |
| Kruppel-associated box (KRAB) domain polypeptide) (ZNF45) ZINC FINGER PROTEIN 46 (ZINC FINGER PROTEIN 46 (ZINC FINGER PROTEIN KUP) (non-exact 62%) Zinc finger protein 6 (CMPX1) (ZNF6) Zinc finger protein 74 (Cos52) (ZNF74) (non-exact, 67%) Zinc finger protein 76 (expressed in testis) (ZNF76) ZINC FINGER PROTEIN 83 (ZINC FINGER PROTEIN HPF1) (non-exact 65%) Zinc finger protein 84 (HPF2) (ZNF84) Zinc finger protein 85 (ZNF85)) Zinc finger protein 9 (ZNF9) ZINC FINGER PROTEIN 93 (=ZINC FINGER PROTEIN 93 (=ZINC FINGER PROTEIN HTF34) (non-exact 70%) Zinc finger protein C2H2-25 | 1 1 1 2 5 | P24278 X56465 X71623 M91592 P51522 M27878 U35376 M28372 | Tactivated | + | + | + | + | + | |
| Kruppel-associated box (KRAB) domain polypeptide) (ZNF45) ZINC FINGER PROTEIN 46 (ZINC FINGER PROTEIN 46 (ZINC FINGER PROTEIN KUP) (non-exact 62%) Zinc finger protein 6 (CMPX1) (ZNF6) Zinc finger protein 74 (Cos52) (ZNF74) (non-exact, 67%) Zinc finger protein 76 (expressed in testis) (ZNF76) ZINC FINGER PROTEIN 83 (ZINC FINGER PROTEIN HPF1) (non-exact 65%) Zinc finger protein 84 (HPF2) (ZNF84) Zinc finger protein 85 (ZNF85)) Zinc finger protein 9 (ZNF9) ZINC FINGER PROTEIN 93 (=ZINC FINGER PROTEIN 93 (=ZINC FINGER PROTEIN 93 (=ZINC FINGER PROTEIN HTF34) (non-exact 70%) Zinc finger protein C2H2-25 (ZNF25) Zinc finger protein clone | 1 1 1 2 5 1 | P24278 X56465 X71623 M91592 P51522 M27878 U35376 M28372 P35789 | T activated | + + + + | + | +++++ | + | + | |
| Kruppel-associated box (KRAB) domain polypeptide) (ZNF45) ZINC FINGER PROTEIN 46 (ZINC FINGER PROTEIN 46 (ZINC FINGER PROTEIN KUP) (non-exact 62%) Zinc finger protein 6 (CMPX1) (ZNF6) Zinc finger protein 74 (Cos52) (ZNF74) (non-exact, 67%) Zinc finger protein 76 (expressed in testis) (ZNF76) ZINC FINGER PROTEIN 83 (ZINC FINGER PROTEIN HPF1) (non-exact 65%) Zinc finger protein 84 (HPF2) (ZNF85)) Zinc finger protein 85 (ZNF85)) Zinc finger protein 9 (ZNF9) ZINC FINGER PROTEIN 93 (=ZINC FINGER PROTEIN 93 (=ZINC FINGER PROTEIN 91 (=ZINC FINGER PROTEIN HTF34) (non-exact 70%) Zinc finger protein C2H2-25 (ZNF25) Zinc finger protein done L3-4 Zinc finger protein | 1 1 1 2 5 1 | P24278 X56465 X71623 M91592 P51522 M27878 U35376 M28372 P35789 U38904 | T activated | + + + + | + | +++++ | + | + | |
| Kruppel-associated box (KRAB) domain polypeptide) (ZNF45) ZINC FINGER PROTEIN 46 (ZINC FINGER PROTEIN KUP) (non-exact 62%) Zinc finger protein 6 (CMPX1) (ZNF6) Zinc finger protein 74 (Cos52) (ZNF74) (non-exact, 67%) Zinc finger protein 76 (expressed in testis) (ZNF76) ZINC FINGER PROTEIN 83 (ZINC FINGER PROTEIN HPF1) (non-exact 65%) Zinc finger protein 84 (HPF2) (ZNF84) Zinc finger protein 85 (ZNF85)) Zinc finger protein 9 (ZNF9) ZINC FINGER PROTEIN 93 (=ZINC FINGER PROTEIN HPF1) (2NF9) ZINC FINGER PROTEIN 93 (=ZINC FINGER PROTEIN HTF34) (non-exact 70%) Zinc finger protein C2H2-25 (ZNF25) Zinc finger protein clone L3-4 | 1 1 1 2 5 1 | P24278 X56465 X71623 M91592 P51522 M27878 U35376 M28372 P35789 U38904 AF024706 | | + + + + | + | +++++ | + | + | |

| WO 00/40749 | PCT/CA00/00005 |
|-----------------|----------------|
| 11 0 001 101 13 | 101,0.100,0000 |

| ZINC FINGER PROTEIN HRX (ALL-1) (71%a.a.) | 1 | Q03164 | | | | | | | | | |
|--|---|--------|-------------|---|---|---|--------|-----|-------------|-----|---|
| zinc finger protein HZF4 | 1 | X78927 | | | | | | | | *** | |
| zinc finger protein RIZ | 1 | D45132 | + | + | + | + | \neg | + 1 | | | |
| zinc finger protein, subfamily 1A, 1 (Ikaros) (LYF1) | 1 | U40462 | + | | | | | | | | |
| zinc finger protein, subfamily 1A, 1 (lkaros) (LYF1) (low match) | 1 | U40462 | | | | | | | | | |
| zinc finger transcriptional regulator (GOS24) | 1 | M92844 | | | | | | | | | |
| zinc-finger helicase (hZFH) | 2 | U91543 | + | + | + | + | | + | | | |
| Zn-15 related zinc finger protein (rff) | 1 | U22377 | | + | + | + | Ī | | | | · |
| Zn-15 related zinc finger protein (rlf) (non-exact 56%) | 1 | U22377 | | | | | | | | | |
| ZNF80-linked ERV9 long terminal repeat | 1 | X83497 | | | | | | | | | |
| ZW10 (Drosophila) homolog, centromere/kinetochore protein (ZW10) | 2 | U54996 | | + | | | | | | ··· | |
| zyxin (ZYX) | 4 | X95735 | | | | | | | | | |

Column 1: List of unique genes derived from 6,283 known ESTs from blood cells.

Column 2: Number of genes found in randomly sequenced ESTs from blood cells.

Column 3: Accession number. Column 4: "+" indicates the presence of the unique gene in publicly available cDNA libraries of blood (Bl), brain (Br), heart (H), kidney (K), liver (Li) and lung (Lu). **Comparison to previously identified tissue-specific genes was determined using the GenBank of the National Centre of Biotechnology Information (NCBI) Database.

10

15

5

Discussion

Every cell and tissue comprising the human body share the necessary genetic information required to maintain cellular homeostasis. These "housekeeping" genes function in basic cellular maintenance, including energy metabolism and cellular structure in all cell types. However, in certain situations, even the housekeeping genes show altered expression. Thus, it is necessary to define the use of these genes as internal controls from one investigation to another. Current results from the human blood cell EST database indicate that over 50% of the transcripts are

widely expressed throughout the human body. Most of the cell or tissue specific genes are also detectable in blood cells by RT-PCR analysis.

5

10

15

20

25

For example, isoformic myosin heavy chain genes are known to be generally expressed in cardiac muscle tissue. In the rodent, the βMyHC gene is only highly expressed in the fetus and in diseased states such as overt cardiac hypertrophy, heart failure and diabetes; the αMyHC gene is highly expressed shortly after birth and continues to be expressed in the adult heart. In the human, however, βMyHC is highly expressed in the ventricles from the fetal stage through adulthood. This highly expressed βMyHC, which harbours several mutations, has been demonstrated to be involved in familial hypertrophic cardiomyopathy (Geisterfer-Lowrance *et al.* 1990). It was reported that mutations of βMyHC can be detected by PCR using blood lymphocyte DNA (Ferrie et al., 1992). Most recently, it was also demonstrated that mutations of the myosin-binding protein C in familial hypertrophic cardiomyopathy can be detected in the DNA extracted from lymphocytes (Niimura *et al.*, 1998).

Similarly, APP and APC, which are known to be tissue specific and predominantly expressed in the brain and intestinal tract, are also detectable in the transcripts of blood. These cell- or tissue-specific transcripts are not detectable by Northern blot analysis. However, the low number of transcript copies can be detected by RT-PCR analysis. These findings strongly demonstrate that genes preferentially expressed in specific tissues can be detected by a highly sensitive RT-PCR assay. In recent years, evidence has been obtained to indicate that expression of cell or tissue-restricted genes can be detected in the peripheral blood of patients with metastatic transitional cell carcinoma (Yuasa et al. 1998) and patients with prostate cancer (Gala et al. 1998).

Atrial natriuretic factor (ANF) and zinc finger protein (ZFP), which are known to be highly expressed in heart tissue biopsies and in the plasma of heart failure patients, are also detectable in the transcripts of blood. Differential expression of zinc finger protein among the normal, diabetic and asymptomatic preclinical

subjects may have additional value as a prophylactic "early warning system". On a related note, there is now more attention/discussion in the cardiovascular disease field being focused on Syndrome X, loosely defined as a continuum of hypertension, increasing sugar levels, diabetes, kidney failure, culminating in heart failure, with the possibility of stroke and heart attack at any time in the continuum. The early identification of patients at risk of organ failure has been a challenge to the medical community for some time and the present method has the potential of resolving or, at least, ameliorating this challenge.

10

15

20

25

The present invention demonstrates that a simple drop of blood may be used to determine the quantitative expression of various mRNAs that reflect the health/disease state of the subject through the use of RT-PCR analysis. This entire process takes about three hours or less. The single drop of blood may also be used for multiple RT-PCR analyses. There is no need for large samples and/or costly and time-consuming separation of cell types within the blood for this method as compared to the methods described by Kimoto (1998) and Chelly et al. (1989; 1988). It is believed that the present finding can potentially revolutionize the way that diseases are detected, diagnosed and monitored because it provides a non-invasive, simple, highly sensitive and quick screening for tissue-specific transcripts. The transcripts detected in whole blood have potential as prognostic or diagnostic markers of disease, as they reflect disturbances in homeostasis in the human body. Delineation of the sequences and/or quantitation of the expression levels of these marker genes by RT-PCR will allow for an immediate and accurate diagnostic/prognostic test for disease or to assess the efficacy and monitor a particular therapeutic.

In addition to RT-PCR, other methods of amplifying may also be used for the purpose of measuring/quantitating tissue-specific transcripts in human blood. For example, mass spectrometry may be used to quantify the transcripts (Koster et al., 1996; Fu et al., 1998). The application of presently disclosed method for detecting tissue-specific transcripts in blood does not restrict to subjects undergoing course of

therapy or treatment, it may also be used for monitoring a patient for the onset of overt symptoms of a disease. Furthermore, the present method may be used for detecting any gene transcripts in blood. A kit for diagnosing, prognosing or even predicting a disease may be designed using gene-specific primers or probes derived from a whole blood sample for a specific disease and applied directly to a drop of blood. A cDNA library specific for a disease may be generated from whole blood samples and used for diagnosis, prognosis or even predicting a disease.

The following references were cited herein:

Claudio JO et al. (1998). Genomics 50:44-52.

5

10 Chelly J et al. (1989). Proc. Nat. Acad. Sci. USA. 86:2617-2621.

Chelly J et al. (1988). Nature 333:858-860.

Drews J & Ryser S (1997). Nature Biotech. 15:1318-9.

Ferrie RM et al. (1992). Am. J. Hum. Genet. 51:251-62.

Fu D-J et al. (1998). Nat. Biotech 16: 381-4.

15 Gala JL et al. (1998). Clin. Chem. 44(3):472-81.

Geisterfer-Lowrance AAT et al. (1990). Cell 62:999-1006.

Groden J et al. (1991). Cell 66:589-600.

Hwang DM et al. (1997). Circulation 96:4146-4203.

Jandreski MA & Liew CC (1987). Hum. Genet. 76:47-53.

20 Jin O et al. (1990). Circulation 82:8-16

Kimoto Y (1998). Mol. Gen. Genet 258:233-239.

Koster M et al. (1996). Nat. Biotech 14: 1123-8.

Liew & Jandreski (1986). Proc. Nat. Acad. Sci. USA. 83:3175-3179

Liew CC et al. (1990). Nucleic Acids Res. 18:3647-3651.

25 Liew CC (1993). J Mol. Cell. Cardiol. 25:891-894

Liew CC et al. (1994). Proc. Natl. Acad. Sci. USA. 91:10645-10649.

Liew et al. (1997). Mol. and Cell. Biochem. 172:81-87.

Niimura H et al. (1998). New Eng. J. Med. 338:1248-1257.

Ogawa M (1993). Blood 81:2844-2853.

5

10

15

Santoro IM & Groden J (1997). Cancer Res. 57:488-494.

Yuasa T et al. (1998). Japanese J. Cancer Res. 89:879-882.

Any patents or publications mentioned in this specification are indicative of the levels of those skilled in the art to which the invention pertains. Further, these patents and publications are incorporated by reference herein in their entirety to the same extent as if each individual publication was specifically and individually indicated to be incorporated by reference.

One skilled in the art will appreciate readily that the present invention is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those objects, ends and advantages inherent herein. The present examples, along with the methods, procedures, treatments, molecules, and specific compounds described herein are presently representative of preferred embodiments, are exemplary, and are not intended as limitations on the scope of the invention. Changes therein and other uses will occur to those skilled in the art which are encompassed within the spirit of the invention as defined by the scope of the claims.

WHAT IS CLAIMED IS:

5

- 1. A method for detecting expression of a gene in blood from a subject, comprising the steps of:
 - a) quantifying RNA from a subject blood sample; and
- b) detecting expression of said gene in the quantified RNA, wherein the expression of said gene in said quantified RNA indicates expression of said gene in the subject blood.
- 10 2. The method of claim 1, wherein the quantification is performed by mass spectrometry.
 - 3. A method for detecting expression of one or more genes in blood from a subject, comprising the steps of:
 - a) obtaining a subject blood sample;
 - b) extracting RNA from said blood sample;
 - c) amplifying said RNA;
 - d) generating expressed sequence tags from the amplified RNA product; and
- e) detecting expression of said genes in the expressed sequence tags, wherein the expression of said genes in said expressed sequence tags indicates expression of said genes in the subject blood.
- 4. The method of claim 3, wherein said genes are non-cancer-25 associated genes.
 - 5. The method of claim 3, wherein said genes are tissue-specific genes.

6. The method of claim 3, wherein said subject is a fetus, an embryo, a child, an adult or a non-human animal.

- 5 7. The method of claim 3, wherein the amplification is performed by RT-PCR.
- 8. The method of claim 7, wherein said RT-PCR utilizes primers selected from the group consisting of random sequence primers and gene-specific primers.
 - 9. A method for detecting expression of one or more genes in blood from a subject, comprising the steps of:
 - a) obtaining a subject blood sample:
 - b) extracting DNA fragment(s) from said blood sample;
 - c) amplifying said DNA fragment(s); and
 - d) detecting expression of said genes in the amplified DNA product, wherein the expression of said genes in said amplified DNA product indicates expression of said genes in the subject blood.

20

25

- 10. A method for monitoring a course of therapeutic treatment in an individual, comprising the steps of:
 - a) obtaining a blood sample from said individual;
 - b) extracting RNA from said blood sample;
 - c) amplifying said RNA;
- d) generating expressed sequence tags from the amplified RNA product; and

e) detecting expression of genes in said expressed sequence tags, wherein the expression of said genes is associated with the effect of said therapeutic treatment; and

- f) repeating steps a)-e), wherein the course of said therapeutic treatment is monitored by detecting the change of expression of said genes in the expressed sequence tags.
 - 11. The method of claim 10, wherein the amplification is performed by RT-PCR.

10

15

- 12. The method of claim 11, wherein the change of expression of said genes in the expressed sequence tags is monitored by sequencing the expressed sequence tags and comparing the resulting sequences at various time points.
- 13. The method of claim 11, wherein the change of expression of said genes in the expressed sequence tags is monitored by performing single nucleotide polymorphism analysis and detecting the variation of a single nucleotide in the expressed sequence tags at various time points.
- 14. The method of claim 10, wherein said individual is monitored for the onset of overt symptoms of a disease, and wherein the expression of said genes is associated with the onset of said symptoms.
- 15. A method for diagnosing a disease in a test subject, comprising the steps of:
 - a) generating a cDNA library for said disease from a whole blood sample from a normal subject;

b) generating expressed sequence tag (EST) profile from the normal subject cDNA library;

c) generating a cDNA library for said disease from a whole blood sample from a test subject;

5

15

20

- d) generating EST profile from the test subject cDNA library; and
- e) comparing the test subject EST profile to the normal subject EST profile, wherein if said test subject EST profile differs from said normal subject EST profile, said test subject might be diagnosed with said disease.
- 16. A kit for diagnosing, prognosing or predicting a disease, comprising:
 - a) gene-specific primers; wherein said primers are designed in such a way that the sequences of said primers contain the opposing ends of two adjacent exons for the specific gene with the intron sequence excluded; and
 - b) a carrier, wherein said carrier immobilizes said primer(s).
 - 17. The kit of claim 16, wherein said gene-specific primer(s) are selected from the group consisting of insulin-specific primers, atrial natriuretic factor-specific primers, zinc finger protein gene-specific primers, beta-myosin heavy chain gene-specific primers, amyloid precurser protein gene-specific primers, and adenomatous polyposis-coli protein gene-specific primers.
 - 18. The kit of claim 17, wherein the sequences of said genespecific primers are selected from the group consisting of SEQ ID Nos. 1 and 2, and SEQ ID Nos. 5 and 6.
 - 19. A method for diagnosing, prognosing or predicting a disease in a test subject, comprising the step of:

applying the kit of claim 16 to a test subject whole blood sample, wherein quantitative expression levels of specific genes associated with said disease are detected and compared to the levels of said specific genes expressed in a normal subject, therefore, said disease may be diagnosed, prognosed or predicted.

5

- 20. The method of claim 19, wherein said method is used for monitoring a course of therapeutic treatment or monitoring the onset of overt symptoms of said disease.
- 10 21. A kit for diagnosing, prognosing or predicting a disease, comprising:
 - a) probes derived from a whole blood sample for a specific disease; and
 - b) a carrier, wherein said carrier immobilizes said probes.

15


20


25

22. A method for diagnosing, prognosing or predicting a disease in a test subject, comprising the step of:

applying the kit of claim 21 to a test subject whole blood sample, wherein quantitative expression levels of specific genes associated with said disease are detected and compared to the levels of said specific genes expressed in a normal subject, therefore, said disease may be diagnosed, prognosed or predicted.

- 23. The method of claim 22, wherein said method is used for monitoring a course of therapeutic treatment or monitoring the onset of overt symptoms of said disease.
- 24. A cDNA library specific for a disease, wherein said cDNA library is generated from whole blood samples.












FIGURE 3






FIGURE 4









FIGURE 5


CC slow Sept 1999

6/7


Α

B









Total :13, 283 ESTs

Known: 6,283

Mitochondrial: 405 Ribosome: 498

Cell Division

Repeat: 868 Mis.: 156

Novel: 2,718

Human Blood

☐ Cell Signalling/Communication

Cell structure/Motility



%9

8%

22%

2%

E Gene/Protein expression

%9



26%

I Suchasimed

29%




FIGURE 7

SEQUENCE LISTING

| <110> | Liew, Choong-Chin | |
|------------------------|---|-----------|
| <120> in Blood and | Method for the Detection of Gene Tr | anscripts |
| in brood and | oses inereor | |
| <130> | 2173/0003 | |
| <150> | US Number not yet assigned | |
| <151> | 2000-01-04 | |
| <150> | US 60/115,125 | |
| <151> | 1999-01-06 | |
| <160> | 10 | |
| | | |
| <210> | 1 | |
| <211> | 18 | |
| <212> | DNA | |
| <213> | artificial sequence | |
| <220> | | |
| <221> | primer_bind | |
| <223> for quantitat | forward primer of exon 1 of insuling tive RT-PCR analysis | gene used |
| <400> | 1 | |
| gccctctggg ga | acctgac | 18 |
| <210> | 2 | |
| <211> | 18 | |
| | DNA | |
| <213> | artificial sequence | |
| <220> | a-01-01-01-01-01-01-01-01-01-01-01-01-01- | |
| | primer bind | |
| <223> | reverse primer of exons 1 and 2 or | f insulin |
| | quantitative RT-PCR analysis | |
| | | |
| <400> | 2 | |
| cccacctgca gg | tcctct | 18 |

| WO 00/40749 | 9 | PCT/CA00/00005 |
|---------------------|---|------------------|
| <210> | 3 | • |
| <211> | 24 | |
| <212> | DNA | |
| <213> | artificial sequence | |
| <220> | | |
| <221> | primer_bind | |
| <223> quantitati | forward primer of βMyHC ve RT-PCR analysis | gene used for |
| <400> | 3 | |
| gctggaacgt | agagactccc tgct | 24 |
| <210> | 4 | |
| <211> | 24 | |
| <212> | DNA | • |
| <213> | artificial sequence | |
| <220> | | |
| <221> | primer_bind | |
| <223> quantitati | reverse primer of β MyHC ve RT-PCR analysis | gene used for |
| <400> | 4 | • |
| ggateettee | agatcatcca cttg | 24 |
| <210> | 5 | |
| <211> | 20 | |
| <212> | DNA | |
| <213> | artificial sequence | |
| <220> | | |
| | <pre>primer_bind</pre> | |
| <223> RT-PCR | | for quantitative |
| | analysis | |
| <400> | 5 | |
| ggatttcaag | aatttgctgg | 20 |

| WO 00/40749 | PCT/CA00/00005 |
|--------------------------|---|
| <210> | 6 |
| <211> | 20 |
| <212> | DNA |
| <213> | artificial sequence |
| <220> | |
| <221> | primer_bind |
| <223> RT-PCR analysis | reverse primer of ANF used for quantitative |
| <400> | 6 |
| gcagatcgat cag | aggagtc 20 |
| <210> | 7 |
| <211> | 20 |
| <212> | DNA |
| <213> | artificial sequence |
| <220> | - |
| <221> | primer bind |
| <223> RT-PCR | forward primer of APP used for quantitative |
| | analysis |
| <400> | 7 |
| ggatgcttca tgtg | gaacgtg 20 |
| | |
| <210> | 8 |
| <211> | 19 |
| <212> | DNA |
| <213> | artificial sequence |
| <220> | |
| <221> | primer_bind |
| <223> RT-PCR | reverse primer of APP used for quantitative |
| | analysis |
| <400> | 8 |

19

tcattcacac cagcacatg

| WO 00/40749 | PCT/CA00/000 | 05 |
|-------------------------|--|----|
| <210> | 9 | |
| <211> | 21 | |
| <212> | DNA | |
| <213> | artificial sequence | |
| <220> | | |
| <221> | primer_bind | |
| <223> RT-PCR analys | forward primer of ZFP used for quantitative is | |
| <400> | 9 | |
| cacargagrc ar | ggtcaacg a 21 | |
| | | |
| <210> | 10 | |
| <211> | 22 | |
| <212> | DNA . | |
| <213> | artificial sequence | |
| <220> | | |
| <221> | primer_bind | |
| <223> RT-PCR analys: | reverse primer of ZFP used for quantitative is | |
| <400> | 10 | |
| ggattaaaat ga | agcaccca ga 22 | |

(19) World Intellectual Property Organization International Bureau





(43) International Publication Date 13 July 2000 (13.07.2000)

PCT

(10) International Publication Number WO 00/40749 A3

(51) International Patent Classification7:

_ -

(21) International Application Number: PCT/CA00/00005

(22) International Filing Date: 5 January 2000 (05.01.2000)

(25) Filing Language:

English

C12Q 1/68

(26) Publication Language:

English

(30) Priority Data:

60/115,125 09/477,148 6 January 1999 (06.01.1999) US 4 January 2000 (04.01.2000) US

(71) Applicant (for all designated States except US): GENE-NEWS INC. [CA/CA]; 45 Bevdale Road, Toronto, Ontario, M2R 1L8 (CA).

(72) Inventor; and

(75) Inventor/Applicant (for US only): LIEW, Choong-Chin [CA/CA]; 81 Millersgrove Drive, Willowdale, Ontario M2R 3S1 (CA).


- (74) Agent: DEETH WILLIAMS WALL; National Bank Building, Suite 400, 150 York Street, Toronto, Ontario M5H 3S5 (CA).
- (81) Designated States (national): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LY, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).


Published:

- with international search report
- (88) Date of publication of the international search report: 19 July 2001

[Continued on next page]

(54) Title: METHOD FOR THE DETECTION OF GENE TRANSCRIPTS IN BLOOD AND USES THEREOF





00/40749 A3

(57) Abstract: The present invention is directed to detection and measurement of gene transcripts in blood. Specifically provided is a RT-PCR analysis performed on a drop of blood for detecting, diagnosing and monitoring diseases using tissue-specific primers. The present invention also describes methods by which delineation of the sequence and/or quantitation of the expression levels of disease-associated genes allows for an immediate and accurate diagnostic/prognostic test for disease or to assess the effect of a particular treatment regimen.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Inter anal Application No PCT/CA 00/00005

| A. CLASSIFI | CATION OF SUB. | ECT MATTER |
|-------------|----------------|------------|
| IPC 7 | C12Q1/68 | |

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, MEDLINE, CHEM ABS Data, BIOSIS, EMBASE, EMBL

| Category ° | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|--|-----------------------|
| X | WO 98 49342 A (COCKBAIN JULIAN R M ;FORSKNINGSPARKEN I AAS AS (NO); LOENNEBORG AN) | 1-15, 21-24 |
| Y | 5 November 1998 (1998-11-05) the whole document | 19,20 |
| X | WO 98 24935 A (AN GANG ;HARA MARK O (US); RALPH DAVID (US); VELTRI ROBERT (US); U) 11 June 1998 (1998-06-11) | 1-15, 21-24 |
| Υ | the whole document | 19,20 |
| X | EP 0 534 640 A (PFIZER) | 16 |
| Υ | 31 March 1993 (1993-03-31) the whole document | 17-20 |
| | -/ | |
| | | |
| | | |

| X Further documents are listed in the continuation of box C. | Patent family members are listed in annex. |
|---|--|
| "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filling date "L" document which may throw doubts on priority claim(s) or | "T" later document published after the international filling date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone |
| which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed | "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family |
| Date of the actual completion of the international search 27 June 2000 | ≟ate of maliting of the international search report 12/07/2000 |
| Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL – 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016 | Authorized officer Hagenmaier, S |

Inter xial Application No PCT/CA 00/00005

| Category ° | ation) DOCUMENTS CONSIDERED TO BE RELEVANT Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|--|-----------------------|
| Y | PATENT ABSTRACTS OF JAPAN vol. 1997, no. 11, 28 November 1997 (1997-11-28) & JP 09 187299 A (NIPPON BIO SERAPII KK), 22 July 1997 (1997-07-22) abstract & DATABASE GCG_GENESEQ 'Online! AC:V00198, March 1998 (1998-03) NIPPON BIOTHERAPY: "HUMAN INSULIN SENSE PRIMER INS-7" abstract | 17,18 |
| Y | DATABASE GENBANK 'Online! AC:V00565, March 1995 (1995-03) BELL ET AL.: "HUMAN GENE FOR PREPROINSULIN" XP002141055 abstract | 17,18 |
| Y | DATABASE GENBANK 'Online! AC:M54947, April 1993 (1993-04) SEIDMAN ET AL.: "HUMAN ATRIAL NATRIURETIC FACTOR GENE" XP002141054 abstract | 17,18 |
| Y | DATABASE GENBANK 'Online! AC:X52889, September 1993 (1993-09) LIEW: "HUMAN GENE FOR CARDIAC BETA MYOSIN HEAVY CHAIN" XP002141056 abstract | 17 |
| Y | DATABASE GENBANK 'Online! AC:2808656, December 1998 (1998-12) BERNOT ET AL.: "A TRANSCRIPTIONAL MAP OF THE FMF REGION/ZINC FINGER PROTEIN" XP002141057 abstract | 17 |
| Υ | YOSHIKAI ET AL.: "GENOMIC ORGANIZATION OF THE HUMAN AMYLOID BETA-PROTEIN PRECURSOR GENE" GENE, vol. 87, 1990, pages 257-263, XP002141053 the whole document | 17 |
| Y | DATABASE GENBANK 'Online! AC:M73548, January 1995 (1995-01) JOSLYN ET AL.: "HUMAN POLYPOSIS LOCUS mRNA" XP002141058 abstract | 17 |
| | -/ | |

Inter onal Application No PCT/CA 00/00005

| | ation) DOCUMENTS CONSIDERED TO BE RELEVANT | Relevant to claim No. |
|------------|--|-----------------------|
| Category ° | Citation of document, with indication, where appropriate, of the relevant passages | TOOTON O GONT, 140. |
| A | WO 98 18906 A (CLAYCOMB WILLIAM CREIGHTON; PROCTER & GAMBLE (US)) 7 May 1998 (1998-05-07) the whole document | |
| A | US 5 352 775 A (MARKHAM ALEXANDER F ET AL) 4 October 1994 (1994-10-04) the whole document | |
| 4 | DE 44 35 919 C (DEUTSCHES KREBSFORSCH) 7 December 1995 (1995-12-07) the whole document | |
| 4 | US 5 837 449 A (ECKER DAVID J ET AL) 17 November 1998 (1998-11-17) the whole document | |
| 4 | WO 98 33942 A (NAKAO KOICHI ;BRISTOW MICHAEL R (US); LEINWAND LESLIE A (US); MINO) 6 August 1998 (1998-08-06) the whole document | · |
| | ` | |
| | | |
| | | |
| | | |
| | | , |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |

information on patent family members

Inter onal Application No
PCT/CA 00/00005

| , - | tent document I in search report | | Publication date | | atent family member(s) | Publicatio date | n |
|------------|---|-------|------------------|------|---------------------------|----------------------|-----------|
| шо | 9849342 | Α | 05-11-1998 | AU | 7222698 | A 24-11- | 1998 |
| WU | 7047346 | n | 00 11 1000 | EP | 0979308 | | |
| | | | | NO | 995296 | | 1999 |
| MU | 9824935 | Α | 11-06-1998 | AU | 5515198 | A 29-06- | 1998 |
| WU | 3024333 | ,, | | EP | 0960214 | | 1999 |
| FP | 0534640 | Α | 31-03-1993 | AT | 143700 | | |
| | 0301010 | • • | | CA | 2078703 | | |
| | | | | DE | 69214243 | | |
| | | | | DE | | T 06-02- | |
| | | | | DK | 534640 | T 17-03- | |
| | | | | ES | 2092056 | T 16-11- | |
| | | | | FΙ | 924242 | | |
| | | | | GR | 3021721 | | |
| | | | | JP | 2703156 | | |
| | | | | JP | 5192199 | A 03-08 - | 1993 |
| | | | | US | 5643730 | A 01-07- | 1997 |
| JP | 09187299 | A | 22-07-1997 | NONE | | | |
| MU | 9818906 | A | 07-05-1998 | AU | 5149998 | A 22-05- | |
| NO | 3010300 | • • | | EP | 0956341 | | |
| | | | | NO | 991894 | A 24-06- | 1999 |
| IIS | 5352775 | Α | 04-10-1994 | US | 5648212 | | |
| • | •••• | | | US | 5691454 | | |
| | | | | US | 5783666 | | |
| | | | | US | RE36713 | | |
| | | | | AU | 1366992 | | |
| | | | | EP | 056 9 527 | | |
| | | | | WO | 9213103 | A 06-08- | -1992 |
| DF | 4435919 | | 07-12-1995 | WO | 9611267 | | |
| <i>.</i> | , | - | | EP | 0784680 | | |
| | | | | JP | 10506789 | T 07-07- | -1998 |
| IIS | 5837449 | А | 17-11-1998 | AU | 3249793 | | |
| J J | , 300.445 | | | .CA | 2126451 | | |
| | | | | EP | 0644889 | | |
| | | | | JP | 6511387 | | |
| | | | | WO | 9313114 | A 08-07- | -199: |
| | 9833942 | Α | 06-08-1998 | AU | 6141098 | A 25-08- | -199 |

CORRECTED VERSION

(19) World Intellectual Property Organization International Bureau



(43) International Publication Date 13 July 2000 (13.07.2000)

PCT

(10) International Publication Number WO 00/40749 A3

- (51) International Patent Classification7:
- -
- (21) International Application Number: PCT/CA00/00005
- (22) International Filing Date: 5 January 2000 (05.01.2000)
- (25) Filing Language:

English

C12Q 1/68

(26) Publication Language:

English

(30) Priority Data:


60/115,125 09/477,148 6 January 1999 (06.01.1999) US 4 January 2000 (04.01.2000) US

(71) Applicant (for all designated States except US): GENE-NEWS INC. [CA/CA]; 45 Bevdale Road, Toronto, Ontario, M2R 1L8 (CA).


- (72) Inventor; and
- (75) Inventor/Applicant (for US only): LIEW, Choong-Chin [CA/CA]; 81 Millersgrove Drive, Willowdale, Ontario M2R 3S1 (CA).
- (74) Agent: DEETH WILLIAMS WALL; National Bank Building, Suite 400, 150 York Street, Toronto, Ontario M5H 3S5 (CA).
- (81) Designated States (national): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.


[Continued on next page]

(54) Title: METHOD FOR THE DETECTION OF GENE TRANSCRIPTS IN BLOOD AND USES THEREOF



(57) Abstract: The present invention is directed to detection and measurement of gene transcripts in blood. Specifically provided is a RT-PCR analysis performed on a drop of blood for detecting, diagnosing and monitoring diseases using tissue-specific primers. The present invention also describes methods by which delineation of the sequence and/or quantitation of the expression levels of disease-associated genes allows for an immediate and accurate diagnostic/prognostic test for disease or to assess the effect of a particular treatment regimen.





WO 00/40749 A3



(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published

- with international search report
- (88) Date of publication of the international search report: 19 July 2001
- (48) Date of publication of this corrected version: 23 August 2001
- (15) Information about Correction: see PCT Gazette No. 34/2001 of 23 August 2001, Section II

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

METHOD FOR THE DETECTION OF GENE TRANSCRIPTS IN BLOOD AND USES THEREOF

BACKGROUND OF THE INVENTION

Cross-Reference to Related Application

This application claims the benefit of priority of provisional patent application U.S. Serial Number 60/115,125, filed January 6, 1999 and of a U.S. application entitled "Method for the Detection of Gene Transcripts in Blood and uses Thereof" filed on January 4, 2000 (application number not yet assigned).

Field of the Invention

5

10

15

20

25

The present invention relates generally to the molecular biology of human diseases. More specifically, the present invention relates to a process using the genetic information contained in human peripheral whole blood for the diagnosis, prognosis and monitoring of genetic and infectious disease in the human body.

Description of the Related Art

The blood is a vital part of the human circulatory system for the human body. Numerous cell types make up the blood tissue including monocytes, leukocytes, lymphocytes and erythrocytes. Although many blood cell types have been described, there are likely many as yet undiscovered cell types in the human blood. Some of these undiscovered cells may exist transiently, such as those derived from tissues and organs that are constantly interacting with the circulating blood in health and disease. Thus, the blood can provide an immediate picture of what is happening in the human body at any given time.

The turnover of cells in the hematopoietic system is enormous. It was reported that over one trillion cells, including 200 billion erythrocytes and 70 billion neutrophilic leukocytes, turn over each day in the human body (Ogawa 1993). As a consequence of continuous interactions between the blood and the body, genetic changes that occur within the cells or tissues of the body will trigger specific changes in gene expression within blood. It is the goal of the present invention that these genetic alterations be harnessed for diagnostic and prognostic purposes, which may lead to the development of therapeutics for ameliorating disease.

5

10

15

20

25

The complete profile of gene expression in the circulating blood remains totally unexplored. It is hypothesized that gene expression in the blood is reflective of body state and, as such, the resultant disruption of homeostasis under conditions of disease can be detected through analysis of transcripts differentially expressed in the blood alone. Thus, the identification of several key transcripts or genetic markers in blood will provide information about the genetic state of the cells, tissues, organs and systems of the human body in health and disease.

The prior art is deficient in non-invasive methods of screening for tissue-specific diseases. The present invention fulfills this long-standing need and desire in the art.

SUMMARY OF THE INVENTION

This present invention discloses a process of using the genetic information contained in human peripheral whole blood in the diagnosis, prognosis and monitoring of genetic and infectious disease in the human body. The process described herein requires a simple blood sample and is, therefore, non-invasive compared to conventional practices used to detect tissue specific disease, such as biopsies.

One object of the present invention is to provide a non-invasive method for the diagnosis, prognosis and monitoring of genetic and infectious disease in humans and animals.

In one embodiment of the present invention, there is provided a method for detecting expression of a gene in blood from a subject, comprising the steps of: a) quantifying RNA from a subject blood sample; and b) detecting expression of the gene in the quantified RNA, wherein the expression of the gene in quantified RNA indicates the expression of the gene in the subject blood.

5

10

15

20

25

In another embodiment of the present invention, there is provided a method for detecting expression of one or more genes in blood from a subject, comprising the steps of: a) obtaining a subject blood sample; b) extracting RNA from the blood sample; c) amplifying the RNA; d) generating expressed sequence tags (ESTs) from the amplified RNA product; and e) detecting expression of the genes in the ESTs, wherein the expression of the genes in the ESTs indicates the expression of the genes in the subject blood. Preferably, the genes are tissue-specific genes.

In still another embodiment of the present invention, there is provided a method for detecting expression of one or more genes in blood from a subject, comprising the steps of: a) obtaining a subject blood sample; b) extracting DNA fragments from the blood sample; c) amplifying the DNA fragments; and d) detecting expression of the genes in the amplified DNA product, wherein the expression of the genes in the subject blood.

In yet another embodiment of the present invention, there is provided a method for monitoring a course of a therapeutic treatment in an individual, comprising the steps of: a) obtaining a blood sample from the individual; b) extracting RNA from the blood sample; c) amplifying the RNA; d) generating expressed sequence tags (ESTs) from the amplified RNA product; e) detecting expression of genes in the ESTs, wherein the expression of the genes is associated with the effect of

the therapeutic treatment: and f) repeating steps a)-e), wherein the course of the therapeutic treatment is monitored by detecting the change of expression of the genes in the ESTs. Such a method may also be used for monitoring the onset of overt symptoms of a disease, wherein the expression of the genes is associated with the onset of the symptoms.

5

10

15

20

25

In still yet another embodiment of the present invention, there is provided a method for diagnosing a disease in a test subject, comprising the steps of:

a) generating a cDNA library for the disease from a whole blood sample from a normal subject; b) generating expressed sequence tag (EST) profile from the normal subject cDNA library; c) generating a cDNA library for the disease from a whole blood sample from a test subject; d) generating EST profile from the test subject cDNA library; and e) comparing the test subject EST profile to the normal subject EST profile, wherein if the test subject EST profile differs from the normal subject EST profile, the test subject might be diagnosed with the disease.

In still yet another embodiment of the present invention, there is provided a kit for diagnosing, prognosing or predicting a disease, comprising: a) genespecific primers; wherein the primers are designed in such a way that their sequences contain the opposing ends of two adjacent exons for the specific gene with the intron sequence excluded; and b) a carrier, wherein the carrier immobilizes the primer(s). Such a kit may be applied to a test subject whole blood sample to diagnose, prognose or predict a disease.

In yet another embodiment of the present invention, there is provided a kit for diagnosing, prognosing or predicting a disease, comprising: a) probes derived from a whole blood sample for a specific disease; and b) a carrier, wherein the carrier immobilizes the probes. Such a kit may be applied to a test subject whole blood sample to diagnose, prognose or predict a disease.

Furthermore, the present invention provides a cDNA library specific for a disease, wherein the cDNA library is generated from whole blood samples.

Other and further aspects, features, and advantages of the present invention will be apparent from the following description of the presently preferred embodiments of the invention. These embodiments are given for the purpose of disclosure.

5

10

15

20

25

BRIEF DESCRIPTION OF THE DRAWINGS

So that the matter in which the above-recited features, advantages and objects of the invention, as well as others which will become clear, are attained and can be understood in detail, more particular descriptions of the invention briefly summarized above may be had by reference to certain embodiments thereof which are illustrated in the appended drawings. These drawings form a part of the specification. It is to be noted, however, that the appended drawings illustrate preferred embodiments of the invention and therefore are not to be considered limiting in their scope.not be considered to limit the scope of the invention.

Figure 1 shows the following RNA samples prepared from human blood; Figure 1A: Lane 1, Molecular weight marker; Lane 2, RT-PCR on APP gene; Lane 3, PCR on APP gene; Lane 4, RT-PCR on APC gene; Lane 5, PCR on APC gene; Figure 1B: Lanes 1 and 2, RT-PCR and PCR of βMyHC, respectively; Lanes 3 and 4, RT-PCR of βMyHC from RNA prepared from human fetal and human adult heart, respectively; Lane 5, Molecular weight marker.

Figure 2 shows quantitative RT-PCR analysis performed on RNA samples extracted from a drop of blood. Forward primer (5'-GCCCTCTGGGGACCTGAC-3', SEQ ID No. 1) of exon 1 and reverse primer (5'-CCCACCTGCAGGTCCTCT-3", SEQ ID No. 2) of exons 1 and 2 of insulin gene. Blood samples of 4 normal subjects were assayed. Lanes 1, 3, 5 and 7 represent overnight "fasting" blood sample and lanes 2, 4, 6 and 8 represent "non-fasting" samples.

Figure 3 shows quantitative RT-PCR analysis performed on RNA samples extracted from a drop of blood. Lanes 1 and 2 represent normal healthy person and lane 3 represents late-onset diabetes (Type II) and lane 4 represents asymptomatic diabetes.

Figure 4 shows multiple RT-PCR assay in a drop of blood. Primers were derived from insulin gene (INS), zinc-finger protein gene (ZFP) and house-keeping gene (GADH). Lane 1 represents normal person. Lane 2 represents late-onset diabetes and lane 3 represents asymptomatic diabetes.

Figure 5 shows standardized levels of insulin gene (Figure 5A) and ZFP gene (Figure 5B) expressed in a drop of blood. The first three subjects were normal, second two subjects showed normal glucose tolerance, and the last subject had late onset diabetes type II. Figure 5C shows standardized levels of insulin gene expressed in each fractionated cell from whole blood.

Figure 6 shows the differential screening of human blood cell cDNA library with different cDNA probes of heart and brain tissue. Figure 6A shows blood cell cDNA probes vs. adult heart cDNA probes. Figure 6B shows blood cell cDNA probes vs. human brain cDNA probes.

Figure 7 graphically shows the 1,800 unique genes in human blood and in the human fetal heart grouped into seven cellular functions.

20

25

5

10

15

DETAILED DESCRIPTION OF THE INVENTION

In accordance with the present invention, there may be employed conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. See, e.g., Sambrook, Fritsch & Maniatis, "Molecular Cloning: A Laboratory Manual (1982); "DNA Cloning: A Practical Approach," Volumes I and II (D.N. Glover ed. 1985); "Oligonucleotide Synthesis" (M.J. Gait ed. 1984); "Nucleic Acid

Hybridization" [B.D. Hames & S.J. Higgins eds. (1985)]; "Transcription and Translation" [B.D. Hames & S.J. Higgins eds. (1984)]; "Animal Cell Culture" [R.I. Freshney, ed. (1986)]; "Immobilized Cells And Enzymes" [IRL Press, (1986)]; B. Perbal, "A Practical Guide To Molecular Cloning" (1984). Therefore, if appearing herein, the following terms shall have the definitions set out below.

5

10

15

20

25

A "cDNA" is defined as copy-DNA or complementary-DNA, and is a product of a reverse transcription reaction from an mRNA transcript. "RT-PCR" refers to reverse transcription polymerase chain reaction and results in production of cDNAs that are complementary to the mRNA template(s).

The term "oligonucleotide" is defined as a molecule comprised of two or more deoxyribonucleotides, preferably more than three. Its exact size will depend upon many factors which, in turn, depend upon the ultimate function and use of the oligonucleotide. The term "primer" as used herein refers to an oligonucleotide, whether occurring naturally as in a purified restriction digest or produced synthetically, which is capable of acting as a point of initiation of synthesis when placed under conditions in which synthesis of a primer extension product, which is complementary to a nucleic acid strand, is induced, i.e., in the presence of nucleotides and an inducing agent such as a DNA polymerase and at a suitable temperature and pH. The primer may be either single-stranded or double-stranded and must be sufficiently long to prime the synthesis of the desired extension product in the presence of the inducing agent. The exact length of the primer will depend upon many factors, including temperature, source of primer and the method used. For example, for diagnostic applications, depending on the complexity of the target sequence, the oligonucleotide primer typically contains 15-25 or more nucleotides, although it may contain fewer nucleotides. The factors involved in determining the appropriate length of primer are readily known to one of ordinary skill in the art.

As used herein, random sequence primers refer to a composition of primers of random sequence, i.e. not directed towards a specific sequence. These

sequences possess sufficient complementary to hybridize with a polynucleotide and the primer sequence need not reflect the exact sequence of the template.

"Restriction fragment length polymorphism" refers to variations in DNA sequence detected by variations in the length of DNA fragments generated by restriction endonuclease digestion.

5

10

15

20

25

A standard Northern blot assay can be used to ascertain the relative amounts of mRNA in a cell or tissue obtained from plant or other tissue, in accordance with conventional Northern hybridization techniques known to those persons of ordinary skill in the art. The Northern blot uses a hybridization probe, e.g. radiolabelled cDNA, either containing the full-length, single stranded DNA or a fragment of that DNA sequence at least 20 (preferably at least 30, more preferably at least 50, and most preferably at least 100 consecutive nucleotides in length). The DNA hybridization probe can be labelled by any of the many different methods known to those skilled in this art. The labels most commonly employed for these studies are radioactive elements, enzymes, chemicals which fluoresce when exposed to untraviolet light, and others. A number of fluorescent materials are known and can be utilized as labels. These include, for example, fluorescein, rhodamine, auramine, Texas Red, AMCA blue and Lucifer Yellow. A particular detecting material is antirabbit antibody prepared in goats and conjugated with fluorescein through an isothiocyanate. Proteins can also be labeled with a radioactive element or with an enzyme. The radioactive label can be detected by any of the currently available counting procedures. The preferred isotope may be selected from ³H, ¹⁴C, ³²P, ³⁵S, ³⁶Cl, ⁵¹Cr, ⁵⁷Co, ⁵⁸Co, ⁵⁹Fe, ⁹⁰Y, ¹²⁵I, ¹³¹I, and ¹⁸⁶Re. Enzyme labels are likewise useful, and can be detected by any of the presently utilized colorimetric, gasometric fluorospectrophotometric, amperometric or spectrophotometric, techniques. The enzyme is conjugated to the selected particle by reaction with bridging molecules such as carbodiimides, diisocyanates, glutaraldehyde and the like. Many enzymes which can be used in these procedures are known and can be utilized.

The preferred are peroxidase, β -glucuronidase, β -D-glucosidase, β -D-galactosidase, urease, glucose oxidase plus peroxidase and alkaline phosphatase. U.S. Patent Nos. 3,654,090, 3,850,752, and 4,016,043 are referred to by way of example for their disclosure of alternate labeling material and methods.

As used herein, "individual" refers to human subjects as well as non-human subjects. The examples herein are not meant to limit the methodology of the present invention to human subjects only, as the instant methodology is useful in the fields of veterinary medicine, animal sciences and such.

5

10

15

20

25

In one embodiment of the present invention, there is provided a method for detecting expression of a gene in blood from a subject, comprising the steps of: a) quantifying RNA from a subject blood sample; and b) detecting expression of the gene in the quantified RNA, wherein the expression of the gene in quantified RNA indicates the expression of the gene in the subject blood. An example of the quantifying method is by mass spectrometry.

In another embodiment of the present invention, there is provided a method for detecting expression of one or more genes in blood from a subject, comprising the steps of: a) obtaining a subject blood sample; b) extracting RNA from the blood sample; c) amplifying the RNA; d) generating expressed sequence tags (ESTs) from the amplified RNA product; and e) detecting expression of the genes in the ESTs, wherein the expression of the genes in the ESTs indicates the expression of the genes in the subject blood. Preferably, the subject is a fetus, an embryo, a child, an adult or a non-human animal. The genes are non-cancer-associated and tissue-specific genes. Still preferably, the amplification is performed by RT-PCR using random sequence primers or gene-specific primers.

In still another embodiment of the present invention, there is provided a method for detecting expression of one or more genes in blood from a subject, comprising the steps of: a) obtaining a subject blood sample; b) extracting DNA fragments from the blood sample; c) amplifying the DNA fragments; and d) detecting

expression of the genes in the amplified DNA product, wherein the expression of the genes in the amplified DNA product indicates the expression of the genes in the subject blood.

5

10

15

20

25

In yet another embodiment of the present invention, there is provided a method for monitoring a course of a therapeutic treatment in an individual, comprising the steps of: a) obtaining a blood sample from the individual; b) extracting RNA from the blood sample; c) amplifying the RNA; d) generating expressed sequence tags (ESTs) from the amplified RNA product; e) detecting expression of genes in the ESTs, wherein the expression of the genes is associated with the effect of the therapeutic treatment; and f) repeating steps a)-e), wherein the course of the therapeutic treatment is monitored by detecting the change of expression of the genes in the ESTs. Such a method may also be used for monitoring the onset of overt symptoms of a disease, wherein the expression of the genes is associated with the onset of the symptoms. Preferably, the amplification is performed by RT-PCR, and the change of the expression of the genes in the ESTs is monitored by sequencing the ESTs and comparing the resulting sequences at various time points; or by performing single nucleotide polymorphism analysis and detecting the variation of a single nucleotide in the ESTs at various time points.

In still yet another embodiment of the present invention, there is provided a method for diagnosing a disease in a test subject, comprising the steps of:

a) generating a cDNA library for the disease from a whole blood sample from a normal subject; b) generating expressed sequence tag (EST) profile from the normal subject cDNA library; c) generating a cDNA library for the disease from a whole blood sample from a test subject; d) generating EST profile from the test subject cDNA library; and e) comparing the test subject EST profile to the normal subject EST profile, wherein if the test subject EST profile differs from the normal subject EST profile, the test subject might be diagnosed with the disease.

In still yet another embodiment of the present invention, there is provided a kit for diagnosing, prognosing or predicting a disease, comprising: a) genespecific primers; wherein the primers are designed in such a way that their sequences contain the opposing ends of two adjacent exons for the specific gene with the intron sequence excluded; and b) a carrier, wherein the carrier immobilizes the primer(s). Preferably, the gene-specific primers are selected from the group consisting of insulinspecific primers, atrial natriuretic factor-specific primers, zinc finger protein genespecific primers, beta-myosin heavy chain gene-specific primers, amyloid precurser protein gene-specific primers, and adenomatous polyposis-coli protein gene-specific primers. Further preferably, the gene-specific primers are selected from the group consisting of SEQ ID Nos. 1 and 2; and SEQ ID Nos. 5 and 6. Such a kit may be applied to a test subject whole blood sample to diagnose, prognose or predict a disease by detecting the quantitative expression levels of specific genes associated with the disease in the test subject and then comparing to the levels of same genes expressed in a normal subject. Such a kit may also be used for monitoring a course of therapeutic treatment or monitoring the onset of overt symptoms of a disease.

5

10

15

20

25

In yet another embodiment of the present invention, there is provided a kit for diagnosing, prognosing or predicting a disease, comprising: a) probes derived from a whole blood sample for a specific disease; and b) a carrier, wherein the carrier immobilizes the probes. Such a kit may be applied to a test subject whole blood sample to diagnose, prognose or predict a disease by detecting the quantitative expression levels of specific genes associated with the disease in the test subject and then comparing to the levels of same genes expressed in a normal subject. Such a kit may also be used for monitoring a course of therapeutic treatment or monitoring the onset of overt symptoms of a disease.

Furthermore, the present invention provides a cDNA library specific for a disease, wherein the cDNA library is generated from whole blood samples.

11

The following examples are given for the purpose of illustrating various embodiments of the invention and are not meant to limit the present invention in any fashion.

5

10

15

EXAMPLE 1

Construction of a cDNA library

RNA extracted from human tissues (including fetal heart, adult heart, liver, brain, prostate gland and whole blood) were used to construct unidirectional cDNA libraries. The first mammalian heart cDNA library was constructed as early as 1982. Since then, the methodology has been revised and optimal conditions have been developed for construction of human heart and hematopoietic progenitor cDNA libraries (Liew et al., 1984; Liew 1993, Claudio et al., 1998). Most of the novel genes which were identified by sequence annotation can now be obtained as full length transcripts.

EXAMPLE 2

Catalogue of blood cell ESTs

20

Random partial sequencing of expressed sequence tags (ESTs) of cDNA clones from the blood cell library was carried out to establish an EST database of blood. The known genes as derived from the ESTs were categorized into seven major cellular functions (Hwang, Dempsey et al., 1997).

EXAMPLE 3

Differential screening of cDNA library

5

10

15

20

25

cDNA probes generated from transcripts of each tissue were used to hybridize the blood cell cDNA clones (Liew et al., 1997). The "positive" signals which were hybridized with P-labelled cDNA probes were defined as genes which shared identity with blood and respective tissues. The "negative" spots which were not exposed to P-labelled cDNA probes were considered to be blood-cell-enriched or low frequency transcripts.

EXAMPLE 4

Reverse transcriptase-polymerase chain reaction (RT-PCR) assay

RNA extracted from samples of human tissue was used for RT-PCR analysis (Jin et al. 1990). Three pairs of forward and reverse primers were designed for human cardiac beta-myosin heavy chain gene (βMyHC), amyloid precurser protein (APP) gene and adenomatous polyposis-coli protein (APC) gene. The PCR products were also subjected to automated DNA sequencing to verify the sequences as derived from the specific transcripts of blood.

EXAMPLE 5

Detection of tissue specific gene expression in human blood using RT-PCR

The beta-myosin heavy chain gene (βMyHC) transcript (mRNA) is known to be highly expressed in ventricles of the human heart. This sarcomeric protein is important for heart muscle contraction and its presence would not be expected in other non-muscle tissues and blood. In 1990, the gene for human cardiac

βMyHC was completely sequenced (Liew et al. 1990) and was comprised of 4 exons and 42 introns.

The method of reverse transcription polymerase chain reaction (RT-PCR) was used to determine whether this cardiac specific mRNA is also present in human blood. A pair of primers was designed; the forward primer (SEQ ID No. 3) was on the boundary of exons 21 and 22, and the reverse primer (SEQ ID No. 4) was on the boundary of exons 24 and 25. This region of mRNA is only present in β MyHC and is not found in the alpha-myosin heavy chain gene (α MyHC).

A blood sample was first treated with lysing buffer and then undergone centrifuge. The resulting pellets were further processed with RT-PCR. RT-PCR was performed using the total blood cell RNA as a template. A nested PCR product was generated and used for sequencing. The sequencing results were subjected to BLAST and the identity of exons 21 to 25 was confirmed to be from βMyHC (Figure 1A).

Using the same method just described, two other tissue specific genes - amyloid precursor protein (APP, forward primer, SEQ ID No. 7; reverse primer, SEQ ID No. 8) found in the brain and associated with Alzheimer's disease, and adenomatous polyposis coli protein (APC) found in the colon and rectum and associated with colorectal cancer (Groden *et al.* 1991; Santoro and Groden 1997) - were also detected in the RNA extracted from human blood (Figure 1B).

20

25

5

10

15

EXAMPLE 6

Multiple RT-PCR analysis on a drop of blood from a normal/diseased individual

A drop of blood was extracted to obtain RNA to carry out quantitative RT-PCR analysis. Specific primers for the insulin gene were designed: forward primer (5'-GCCCTCTGGGGACCTGAC-3', SEQ ID No. 1) of exon 1 and reverse primer (5'-CCCACCTGCAGGTCCTCT-3", SEQ ID No. 2) of exons 1 and 2 of insulin gene. Such reverse primer was obtained by deleting the intron between the

exons 1 and 2. Blood samples of 4 normal subjects were assayed. It was found that the insulin gene is expressed in the blood and the quantitative expression of the insulin gene in a drop of blood is influenced by fasting and non-fasting states of normal healthy subjects (Figure 2). This very low level of expression of the insulin gene reflects the phenotypic status of a person and strongly suggests that there is a physiological and pathological role for its expression, contrary to the basal or illegitimate theory of transcription suggested by Chelly *et al.* (1989) and Kimoto (1998).

5

10

15

20

25

Same quantitative RT-PCR analysis was performed using insulin specific primers on RNA samples extracted from a drop of blood from a normal healthy person, a person having late-onset diabetes (Type II) and a person having asymptomatic diabetes. It was found that the insulin gene is expressed differentially amongst subjects that are healthy, diagnosed as type II diabetic, and also in an asymptomatic preclinical patient (Figure 3).

Similarly, specific primers for the atrial natriuretic factor (ANF) gene were designed (forward primer, SEQ ID No. 5; reverse primer, SEQ ID No. 6) and RT-PCR analysis was performed on a drop of blood. ANF is known to be highly expressed in heart tissue biopsies and in the plasma of heart failure patients. However, atrial natriuretic factor was observed to be expressed in the blood and the expression of the atrial natriuretic factor gene is significantly higher in the blood of patients with heart failure as compared to the blood of a normal control patient.

Specific primers for the zinc finger protein gene (ZFP, forward primer, SEQ ID No. 9; reverse primer, SEQ ID No. 10) were also designed and RT-PCR analysis was performed on a drop of blood. ZFP is known to be high in heart tissue biopsies of cardiac hypertrophy and heart failure patients. In the present study, the expression of ZFP was observed in the blood as well as differential expression levels of ZFP amongst the normal, diabetic and asymptomatic preclinical subjects (Figure 4); although neither of the non-normal subjects has been specifically diagnosed as

suffering from cardiac hypertrophy and/or heart failure, the higher expression levels of the ZFP gene in their blood may indicate that these subjects are headed in that general direction.

It was hypothesized that a housekeeping gene such as glyceraldehyde dehydrogenase (GADH) which is required and highly expressed in all cells would not be differentially expressed in the blood of normal vs. disease subjects. This hypothesis was confirmed by RT-PCR using GADH specific primers (Figure 4). Thus, GADH is useful as an internal control.

Standardized levels of insulin gene or ZFP gene expressed in a drop of blood were estimated using a housekeeping gene as an internal control relative to insulin or ZFP expressed (Figures 5A & 5B). The levels of insulin gene expressed in each fractionated cell from whole blood were also standardized and shown in Figure 5C.

EXAMPLE 7

Human blood cell cDNA library

5

10

15

20

25

In order to further substantiate the present invention, differential screening of the human blood cell cDNA library was conducted. cDNA probes derived from human blood, adult heart or brain were respectively hybridized to the human blood cDNA library clones. As shown in Figure 7, more than 95% of the "positively" identified clones are identical between the blood and other tissue samples.

DNA sequencing of randomly selected clones from the human whole blood cell cDNA library was also performed. This allowed information regarding the cellular function of blood to be obtained concurrently with gene identification. More than 20,000 expressed sequence tags (ESTs) have been generated and characterized to date, 17.6% of which did not result in a statistically significant match to entries in the

GenBank databases and thus were designated as "Novel" ESTs. These results are summarized in Figure 7 together with the seven cellular functions related to percent distribution of known genes in blood and in the fetal heart.

From 20,000 ESTs, 1,800 have been identified as known genes which may not all appear in the hemapoietic system. For example, the insulin gene and the atrial natriuretic factor gene have not been detected in these 20,000 ESTs but their transcripts were detected in a drop of blood, strongly suggesting that all transcripts of the human genome can be detected by performing RT-PCR analysis on a drop of blood.

In addition, approximately 400 novel genes have been identified from the 20,000 ESTs characterized to date, and these will be subjected to full length sequencing and open reading frame alignment to reduce the actual number of novel ESTs prior to screening for disease markers.

Analysis of the approximately 6,283 ESTs which have known matches in the GenBank databases revealed that this dataset represents over 1,800 unique genes. These genes have been catalogued into seven cellular functions. Comparisons of this set of unique genes with ESTs derived from human brain, heart, lung and kidney demonstrated a greater than 50% overlap in expression (Table 1).

20 <u>TABLE 1</u>

5

10

15

25

Overlap of Genes Expressed in Blood *

| Tissues | ESTs** | Overlap in Blood |
|---------|---------|------------------|
| brain | 134,000 | 60% |
| heart | 65,000 | 59% |
| lung | 60,200 | 58% |
| kidney | 32,300 | 54% |

* Estimated from limited known genes of about 1,800 as derived from the database of 6,297 ESTs from human blood cell library.

** Obtained from the National Centre of Biotechnology Information (NCBI), U.S.A.

5

10

15

20

EXAMPLE 8

Blood cell ESTs

The results from the differential screening clearly indicate that the transcripts expressed in the whole blood are reflective of genes expressed in all cells and tissues of the body. More than 95% of detectable spots were identical from two different tissues. The remaining 5% of spots may represent cell- or tissue-specific transcripts; however, results obtained from partial sequencing to generate ESTs of these clones revealed most of them not to be cell- or tissue-specific transcripts. Therefore, the negative spots are postulated to be reflective of low abundance transcripts in the tissue from which the cDNA probes were derived.

An alternative approach that was employed to identify transcripts expressed at low levels is the large-scale generation of expressed sequence tags (ESTs). There is substantial evidence regarding the efficiency of this technology to detect previously characterized (known) and uncharacterized (unknown or novel) genes expressed in the cardiovascular system (Hwang & Dempsey *et al.*. 1997). In the present invention, 20,000 ESTs have been produced from a human blood cell cDNA library and resulted in the identification of approximately 1,800 unique known genes (Table 2)

25

In the most recent GenBank release, analysis of more than 300,000 ESTs in the database (dbESTs) generated more than 48,000 gene clusters which are thought to represent approximately 50% of the genes in the human genome. Only 4,800 of the dbESTs are blood-derived. In the present invention, 20,000 ESTs have

been obtained to date from a human blood cDNA library, which provides the world's most informative database with respect to blood cell transcripts. From the limited amount of information generated so far (i.e. 1,800 unique genes), it has already been determined that more than 50% of the transcripts are found in other cells or tissues of the human body (Table 2). Thus, it is expected that by increasing the number of ESTs generated, more genes will be identified that have an overlap in expression between the blood and other tissues. Furthermore, the transcripts for several genes which are known to have tissue-restricted patterns of expression (i.e. βMyHC, APP, APC, ANF, ZFP) have also been demonstrated to be present in blood.

5

10

15

20

Most recently, a cDNA library of human hematopoietic progenitor stem cells has also been constructed. From the limited set of 1,000 ESTs, there are at least 200 known genes that are shared with other tissue related genes (Claudio *et al.* 1998).

Table 2 demonstrates the expression of known genes of specific tissues in blood cells. Previously, only the presence of "housekeeping" genes would have been expected. Additionally, the presence of at least 25 of the currently known 500 genes corresponding to molecular drug targets was detected. These molecular drug targets are used in the treatment of a variety of diseases which involve inflammation, renal and cardiovascular function, neoplastic disease, immunomodulation and viral infection (Drews & Ryser, 1997). It is expected that additional novel ESTs will represent future molecular drug targets.

TABLE 2

Comparison of 1,800 Unique Genes Identified in the Blood Cell cDNA Library to Genes Previously Identified in Specific Tissues

5

| Gene Identification | | Accession No. | Tissue Distribution | | | | | | | | |
|---|--------------|---------------|---------------------|---|----------|----|----------|-----|--|--|--|
| Gene identification | | | В | Br | П | IK | | Lui | | | |
| 100 kDa coactivator | | U22055 | | + | - | | _ | + | | | |
| OkD protein (BC10) | | AF053470 | | + | + | | + | + | | | |
| | | U54778 | | + | + | | | + | | | |
| 14-3-3 epsilon | - | U28964 | | ļ <u>, </u> | + | | + | | | | |
| 14-3-3 protein | 11 | 1 | | l ' | + | | <u> </u> | - | | | |
| 15 kDa selenoprotein (SEP15) | 1 | AF051894 | | + | | | | | | | |
| 1-phosphatidylinositol-4- phosphate 5-kinase soform C | 1 | 578798 | | | | | | | | | |
| 23 kD highly basic protein | 21 | X56932 | + | + | + | + | + | + | | | |
| 2-5A-dependent RNase | 1 | L10381 | | | | | | | | | |
| 2'-5'oligoadenylate synthetase 2 (OAS2) | 4 | M87284 | В | | | | | | | | |
| 26S proteasome subunit 11 | 1 | AF086708 | | | | | | | | | |
| 36 kDa phosphothyrosine protein | 2 | AJ223280 | | | + | | | | | | |
| 3-7 gene product (non- exact 86%aa) | 1 | D64159 | | | | | | | | | |
| 3-phosphoglycerate dehydrogenase (PGAD) | 1 | AF006043 | T | + | + | | | + | | | |
| 3-prime-phosphoadenosine 5-prime-phosphosulfate synthase 1 (PAPSS1) | 2 | U53447 | + | + | Ť | + | | + | | | |
| 46kd mannose 6- phosphate receptor (MPR46) (low match) | 1 | X56257 | | | | | | | | | |
| 5-aminoimidazole 4- carboxamide ribonucleotide transformylase | 1 | D89976 | | | | | | | | | |
| 5'-nucleotidase | 3 | D38524 | T | + | | Ī | + | | | | |
| 6-phosphofructo-2- kinase/fructose-2,6- biphosphatase 4 (PFKFB4) | 1 | D49818 | | + | | | | | | | |
| 6-phosphofructo-2- kinase/fructose-2,6- bisphosphatase (PF2K) | 1 | AF041829 | | | | | | | | | |
| 71 kd heat shock cognate protein hsc70 | 23 | Y00371 | | | | | | | | | |
| 76 kDa membrane protein (P76) | 2 | U81006 | | + | + | + | + | + | | | |
| 8-oxoguanine DNA glycosylase (OGG1) | 1 | U96710 | В | | | | + | + | | | |
| a disintegrin and metalloprotease domain 10 (ADAM10) | 1 | AF009615 | T | | | | + | | | | |
| à disintegrin and metalloprotease domain 8 (ADAM8) | 1 ~ | D26579 | В | + | | | | | | | |
| A kinase anchor protein 95 (AKAP95) | 2 | Y11997 | B, T activated | | + | | | + | | | |
| A kinase anchor protein, 149kD (AKAP149) | 2 | X97335 | | + | + | + | | + | | | |

| A4 differentiation- | 1 | U93305 | | | | | | | |
|--|-----------------------|--|---|---------|-----------|--|--|-------------|------------------------|
| dependent protein (A4), triple LIM domain protein | | | | | | ŀ | | | |
| (LMO6), and | | 1 | | | | | | | |
| synaptophysin (SYP); | | | | | | | | | |
| calcium channel alpha-1 | | | | | | | | | |
| ABL and putative M8604 | | U07561 | | | | | \vdash | | |
| Met protein | , | | | | | | | | |
| Absent in melanoma 1 | 1 | U83115 | + | + | | | | + | |
| (AIM1) | | Z31696 | | + | + | <u> </u> | | | |
| accessory proteins BAP31/BAP29 | 2 | 231090 | | т . | * | | | İ | |
| (DXS1357E) | | | | | ŀ | | | ŀ | |
| acetyl-Coenzyme A | 2 | X12966 | + | + | + | + | + | + | |
| acyltransferase (peroxisomal 3-oxoacyl- | | | | | | 1 | | | |
| Coenzyme A thiolase) | | | | | | | | | |
| (ACAA) | | | | | | l | | | |
| acetyl-Coenzyme A | 1 | D88152 | T lymphoma | + | + | | | | |
| transporter (ACATN) acidic 82 kDa protein | 4 | U15552 | | | | | | ├ | |
| · | · · | 1 | <u> </u> | | <u> </u> | | ļ., | <u> </u> | |
| acidic protein rich in leucines (SSP29) | 1 | Y07969 | В | + | + | | + | + | |
| Aconitase 2. mitochondrial | 1 | U80040 | + | + | + | + | | + | |
| (ACO2) | | _ | <u> </u> | | <u>L</u> | <u></u> | L | | |
| actin binding protein | 1 | AF059569 | | | | | | | |
| MAYVEN actin, beta (ACTB) | 158 | X04098 | Т, В | + | + | ├ | + | | ļ |
| | | M10277 | 1, 5 | , | <u>'</u> | | <u> </u> | | |
| actin, beta (ACTB) (non- exact, low match 73%) | 1 | M102// | | | | | | l | |
| actin, gamma (low score) | 1 | K00791 | | | | ! | | | |
| actin, gamma 1 (ACTG1) | 4 | X04098 | + | + | + | + | + | + | high in many libraries |
| actin-binding LIM protein | 4 | D31883 | <u> </u> | + | + | + | | + | , |
| (ABLIM) | T | 55,000 | | , | ' | | | ' | |
| Actinin, alpha 1 (ACTN1) | 8 | M95178 | | + | + | + | | + | |
| actinin, alpha 4 (ACTN4) | 1 | D89980 | | + | + | | + | | |
| - | | | 1 | | | | | + | |
| activated p21cdc42Hs | 1 | L13738 | В | + | | | 1 | T . | · |
| activated p21cdc42Hs kinase (ACK) | | | | | | | | | |
| kinase (ACK) activated RNA polymerase | 1 | L13738 X79805 | B + | + | + | + | | + | |
| kinase (ACK) activated RNA polymerase II transcription cofactor 4 | | | | | + | + | | | |
| kinase (ACK) activated RNA polymerase II transcription cofactor 4 (PC4) | | | | | + | + | | | |
| kinase (ACK) activated RNA polymerase II transcription cofactor 4 (PC4) activating transcription factor 1 (ATF1) | 1 | X79805 X55544 | | + | + | + | | | |
| kinase (ACK) activated RNA polymerase II transcription cofactor 4 (PC4) activating transcription factor 1 (ATF1) activating transcription | 1 | X79805 | | | | + | + | | |
| kinase (ACK) activated RNA polymerase II transcription cofactor 4 (PC4) activating transcription factor 1 (ATF1) activating transcription factor 2 (ATF2) | 1 | X79805 X55544 X15875 | | + | + | + | | + | |
| kinase (ACK) activated RNA polymerase II transcription cofactor 4 (PC4) activating transcription factor 1 (ATF1) activating transcription factor 2 (ATF2) activating transcription | 1 | X79805 X55544 | | + | + | + | + | | |
| kinase (ACK) activated RNA polymerase II transcription cofactor 4 (PC4) activating transcription factor 1 (ATF1) activating transcription factor 2 (ATF2) activating transcription factor 4 (tax-responsive enhancer element B67) | 1 | X79805 X55544 X15875 | | + | + | + | | + | |
| kinase (ACK) activated RNA polymerase II transcription cofactor 4 (PC4) activating transcription factor 1 (ATF1) activating transcription factor 2 (ATF2) activating transcription factor 4 (tax-responsive enhancer element B67) (ATF4) | 1 1 2 | X79805 X55544 X15875 M86842 | + | + | + | | | + | |
| kinase (ACK) activated RNA polymerase II transcription cofactor 4 (PC4) activating transcription factor 1 (ATF1) activating transcription factor 2 (ATF2) activating transcription factor 4 (tax-responsive enhancer element B67) (ATF4) active BCR-related gene | 1 | X79805 X55544 X15875 | | + | + | + | | + | |
| kinase (ACK) activated RNA polymerase II transcription cofactor 4 (PC4) activating transcription factor 1 (ATF1) activating transcription factor 2 (ATF2) activating transcription factor 4 (tax-responsive enhancer element B67) (ATF4) active BCR-related gene (ABR) | 1 1 2 | X79805 X55544 X15875 M86842 | + | + | + | | | + | |
| kinase (ACK) activated RNA polymerase II transcription cofactor 4 (PC4) activating transcription factor 1 (ATF1) activating transcription factor 2 (ATF2) activating transcription factor 4 (tax-responsive enhancer element B67) (ATF4) active BCR-related gene (ABR) acyl-CoA oxidase (AOX) | 1 1 2 | X79805 X55544 X15875 M86842 U01147 U03254 | + | + | + | | | + | |
| kinase (ACK) activated RNA polymerase II transcription cofactor 4 (PC4) activating transcription factor 1 (ATF1) activating transcription factor 2 (ATF2) activating transcription factor 4 (tax-responsive enhancer element B67) (ATF4) active BCR-related gene (ABR) acyl-CoA oxidase (AOX) acyl-Coenzyme A dehydrogenase. C-4 to C- | 1 1 2 | X79805 X55544 X15875 M86842 | + | + | + | | | + | |
| kinase (ACK) activated RNA polymerase II transcription cofactor 4 (PC4) activating transcription factor 1 (ATF1) activating transcription factor 2 (ATF2) activating transcription factor 4 (tax-responsive enhancer element B67) (ATF4) active BCR-related gene (ABR) acyl-CoA oxidase (AOX) acyl-Coenzyme A dehydrogenase, C-4 to C- 12 straight chain (ACADM) | 1 1 2 | X79805 X55544 X15875 M86842 U01147 U03254 M16827 | + | + | + | + | + | + | |
| kinase (AČK) activated RNA polymerase II transcription cofactor 4 (PC4) activating transcription factor 1 (ATF1) activating transcription factor 2 (ATF2) activating transcription factor 4 (tax-responsive enhancer element B67) (ATF4) active BCR-related gene (ABR) acyl-Coenzyme A dehydrogenase, C-4 to C- 12 straight chain (ACADM) acyl-Coenzyme A | 1 1 2 | X79805 X55544 X15875 M86842 U01147 U03254 | + | + | + | | | + | |
| kinase (ACK) activated RNA polymerase II transcription cofactor 4 (PC4) activating transcription factor 1 (ATF1) activating transcription factor 2 (ATF2) activating transcription factor 4 (tax-responsive enhancer element B67) (ATF4) active BCR-related gene (ABR) acyl-CoA oxidase (AOX) acyl-Coenzyme A dehydrogenase, C-4 to C- 12 straight chain (ACADM) acyl-Coenzyme A dehydrogenase, very long | 1 1 2 | X79805 X55544 X15875 M86842 U01147 U03254 M16827 | + | + | + | + | + | + | |
| kinase (AČK) activated RNA polymerase II transcription cofactor 4 (PC4) activating transcription factor 1 (ATF1) activating transcription factor 2 (ATF2) activating transcription factor 4 (tax-responsive enhancer element B67) (ATF4) active BCR-related gene (ABR) acyl-Coanzyme A dehydrogenase, C-4 to C- 12 straight chain (ACADM) acyl-Coenzyme A dehydrogenase, very long chain (ACADVL) acyloxyacyl hydrolase | 1 1 2 | X79805 X55544 X15875 M86842 U01147 U03254 M16827 | + | + | + | + | + | + | |
| kinase (AČK) activated RNA polymerase II transcription cofactor 4 (PC4) activating transcription factor 1 (ATF1) activating transcription factor 2 (ATF2) activating transcription factor 4 (tax-responsive enhancer element B67) (ATF4) active BCR-related gene (ABR) acyl-CoA oxidase (AOX) acyl-Coenzyme A dehydrogenase, C-4 to C- 12 straight chain (ACADM) acyl-Coenzyme A dehydrogenase, very long chain (ACADVL) acyloxyacyl hydrolase (neutrophil) (AOAH) | 1 1 2 3 3 | X79805 X55544 X15875 M86842 U01147 U03254 M16827 D43682 M62840 | + | + + | + + + | + | + + | + | |
| kinase (ACK) activated RNA polymerase II transcription cofactor 4 (PC4) activating transcription factor 1 (ATF1) activating transcription factor 2 (ATF2) activating transcription factor 4 (tax-responsive enhancer element B67) (ATF4) active BCR-related gene (ABR) acyl-CoA oxidase (AOX) acyl-Coenzyme A dehydrogenase, C-4 to C- 12 straight chain (ACADM) acyl-Coenzyme A dehydrogenase, very long chain (ACADVL) acyloxyacyl hydrolase (neutrophil) (AOAH) adaptin, delta (ADTD) | 1 1 2 1 1 2 3 3 3 2 | X79805 X55544 X15875 M86842 U01147 U03254 M16827 D43682 M62840 U91930 | + | + | + + | + | + | + | |
| kinase (AČK) activated RNA polymerase II transcription cofactor 4 (PC4) activating transcription factor 1 (ATF1) activating transcription factor 2 (ATF2) activating transcription factor 4 (tax-responsive enhancer element B67) (ATF4) active BCR-related gene (ABR) acyl-CoA oxidase (AOX) acyl-Coenzyme A dehydrogenase, C-4 to C- 12 straight chain (ACADM) acyl-Coenzyme A dehydrogenase, very long chain (ACADVL) acyloxyacyl hydrolase (neutrophil) (AOAH) adaptin, delta (ADTD) | 1 1 2 3 3 | X79805 X55544 X15875 M86842 U01147 U03254 M16827 D43682 M62840 | + | + + | + + + | + | + + | + | |
| kinase (AČK) activated RNA polymerase II transcription cofactor 4 (PC4) activating transcription factor 1 (ATF1) activating transcription factor 2 (ATF2) activating transcription factor 4 (tax-responsive enhancer element B67) (ATF4) active BCR-related gene (ABR) acyl-Coenzyme A dehydrogenase, C-4 to C- 12 straight chain (ACADM) acyl-Coenzyme A dehydrogenase, very long chain (ACADVL) acyloxyacyl hydrolase (neutrophil) (AOAH) adaptin, delta (ADTD) adaptin, delta (ADTD) (non-exact 59%) | 1 1 2 3 3 3 2 1 | X79805 X55544 X15875 M86842 U01147 U03254 M16827 D43682 M62840 U91930 AC005328 | + | + + + | + + + + + | + | + + | + + + | |
| kinase (AČK) activated RNA polymerase II transcription cofactor 4 (PC4) activating transcription factor 1 (ATF1) activating transcription factor 2 (ATF2) activating transcription factor 4 (tax-responsive enhancer element B67) (ATF4) active BCR-related gene (ABR) acyl-CoA oxidase (AOX) acyl-Coenzyme A dehydrogenase, C-4 to C- 12 straight chain (ACADM) acyl-Coenzyme A dehydrogenase, very long chain (ACADVL) acyloxyacyl hydrolase (neutrophil) (AOAH) adaptin, delta (ADTD) adaptin, delta (ADTD) (non-exact 59%) adaptin, gamma (ADTG) | 1 1 2 2 3 3 3 2 1 1 1 | X79805 X55544 X15875 M86842 U01147 U03254 M16827 D43682 M62840 U91930 AC005328 Y12226 | + | + + + + | + + + | + | + + | + + + + | |
| kinase (AČK) activated RNA polymerase II transcription cofactor 4 (PC4) activating transcription factor 1 (ATF1) activating transcription factor 2 (ATF2) activating transcription factor 4 (tax-responsive enhancer element B67) (ATF4) active BCR-related gene (ABR) acyl-Coenzyme A dehydrogenase, C-4 to C- 12 straight chain (ACADM) acyl-Coenzyme A dehydrogenase, very long chain (ACADVL) acyloxyacyl hydrolase (neutrophil) (AOAH) adaptin, delta (ADTD) (non-exact 59%) adaptin, gamma (ADTG) adaptor complex sigma3B | 1 1 2 3 3 3 2 1 | X79805 X55544 X15875 M86842 U01147 U03254 M16827 D43682 M62840 U91930 AC005328 | + | + + + | + + + + + | + | + + | + + + | |
| kinase (AČK) activated RNA polymerase II transcription cofactor 4 (PC4) activating transcription factor 1 (ATF1) activating transcription factor 2 (ATF2) activating transcription factor 4 (tax-responsive enhancer element B67) (ATF4) active BCR-related gene (ABR) acyl-CoA oxidase (AOX) acyl-Coenzyme A dehydrogenase, C-4 to C- 12 straight chain (ACADM) acyl-Coenzyme A dehydrogenase, very long chain (ACADVL) acyloxyacyl hydrolase (neutrophil) (AOAH) adaptin, delta (ADTD) adaptin, delta (ADTD) (non-exact 59%) adaptin, gamma (ADTG) | 1 1 2 2 3 3 3 2 1 1 1 | X79805 X55544 X15875 M86842 U01147 U03254 M16827 D43682 M62840 U91930 AC005328 Y12226 | + | + + + + | + + + + + | + | + + | + + + + | |
| kinase (AČK) activated RNA polymerase II transcription cofactor 4 (PC4) activating transcription factor 1 (ATF1) activating transcription factor 2 (ATF2) activating transcription factor 4 (tax-responsive enhancer element B67) (ATF4) active BCR-related gene (ABR) acyl-CoA oxidase (AOX) acyl-Coenzyme A dehydrogenase, C-4 to C- 12 straight chain (ACADM) acyl-Coenzyme A dehydrogenase, very long chain (ACADVL) acyloxyacyl hydrolase (neutrophil) (AOAH) adaptin, delta (ADTD) (non-exact 59%) adaptin, gamma (ADTG) adaptor complex sigma3B (AP3S3) | 1 1 2 3 3 3 2 1 1 2 2 | X79805 X55544 X15875 M86842 U01147 U03254 M16827 D43682 M62840 U91930 AC005328 Y12226 X99459 | + | + + + + | + + + + + | + | + + | + + + + | |

| adducin 1 (alpha) (add1) adducin 3 (gamma) (ADD3) adenine nucleotide translocator 2 (fibroblast) (ANT2) adenine nucleotide translocator 2 (fibroblast) (ANT2) (non-exact 81%) adenine nucleotide translocator 2 (fibroblast) (ANT2) (non-exact, 79%) adenine nucleotide translocator 2 (fibroblast) (ANT2) (non-exact, 79%) adenine nucleotide translocator 2 (fibroblast) (ANT2) (non-exact, 86%) adenine nucleotide translocator 3 (liver) (ANT3) adenosine deaminase, RNA-specific (ADAR) adenylate cyclase 3 (ADCY3) adenylate cyclase 7 (ADCY7) adenylate kinase 2 (AK2) adenylate kinase 3 (AK3) (non-exact, 67%) adenylyl cyclase- | 3 | | + | + | + | + | | + | · · · · · · · · · · · · · · · · · · · |
|---|----|----------|-----------|-----|---|---|---|---|---------------------------------------|
| adenine nucleotide translocator 2 (fibroblast) (ANT2) adenine nucleotide translocator 2 (fibroblast) (ANT2) (non-exact 81%) adenine nucleotide translocator 2 (fibroblast) (ANT2) (non-exact, 79%) adenine nucleotide translocator 2 (fibroblast) (ANT2) (non-exact, 79%) adenine nucleotide translocator 2 (fibroblast) (ANT2) (non-exact, 86%) adenine nucleotide translocator 3 (liver) (ANT3) adenosine deaminase, RNA-specific (ADAR) adenylate cyclase 3 (ADCY3) adenylate cyclase 7 (ADCY7) adenylate kinase 2 (AK2) adenylate kinase 3 (AK3) (non-exact, 67%) | | L29296 | 1 1 | | | _ | | | |
| translocator 2 (fibroblast) (ANT2) adenine nucleotide translocator 2 (fibroblast) (ANT2) (non-exact 81%) adenine nucleotide translocator 2 (fibroblast) (ANT2) (non-exact, 79%) adenine nucleotide translocator 2 (fibroblast) (ANT2) (non-exact, 86%) adenine nucleotide translocator 3 (liver) (ANT3) adenosine deaminase, RNA-specific (ADAR) adenylate cyclase 3 (ADCY3) adenylate cyclase 7 (ADCY7) adenylate kinase 2 (AK2) adenylate kinase 3 (AK3) (non-exact, 67%) | 3 | U37122 | B, W | + | + | | + | + | |
| translocator 2 (fibroblast) (ANT2) (non-exact 81%) adenine nucleotide translocator 2 (fibroblast) (ANT2) (non-exact, 79%) adenine nucleotide translocator 2 (fibroblast) (ANT2) (non-exact, 86%) adenine nucleotide translocator 3 (liver) (ANT3) adenosine deaminase, RNA-specific (ADAR) adenylate cyclase 3 (ADCY3) adenylate cyclase 7 (ADCY7) adenylate kinase 2 (AK2) adenylate kinase 3 (AK3) (non-exact, 67%) | 2 | M57424 | | + | + | | + | | |
| adenine nucleotide translocator 2 (fibroblast) (ANT2) (non-exact, 79%) adenine nucleotide translocator 2 (fibroblast) (ANT2) (non-exact, 86%) adenine nucleotide translocator 3 (liver) (ANT3) adenosine deaminase, RNA-specific (ADAR) adenylate cyclase 3 (ADCY3) adenylate cyclase 7 (ADCY7) adenylate kinase 2 (AK2) adenylate kinase 3 (AK3) (non-exact, 67%) | 1 | J02683 | | | | | | | |
| adenine nucleotide translocator 2 (fibroblast) (ANT2) (non-exact, 86%) adenine nucleotide translocator 3 (liver) (ANT3) adenosine deaminase, RNA-specific (ADAR) adenylate cyclase 3 (ADCY3) adenylate cyclase 7 (ADCY7) adenylate kinase 2 (AK2) adenylate kinase 3 (AK3) (non-exact, 67%) | 1 | J02683 | | | | | | | |
| adenine nucleotide translocator 3 (liver) (ANT3) adenosine deaminase, RNA-specific (ADAR) adenylate cyclase 3 (ADCY3) adenylate cyclase 7 (ADCY7) adenylate kinase 2 (AK2) adenylate kinase 3 (AK3) (non-exact, 67%) | 1 | J02683 | | | | | | | |
| adenosine deaminase, RNA-specific (ADAR) adenylate cyclase 3 (ADCY3) adenylate cyclase 7 (ADCY7) adenylate kinase 2 (AK2) adenylate kinase 3 (AK3) (non-exact, 67%) | 3 | J03592 | | + | + | | + | + | |
| adenylate cyclase 3 (ADCY3) adenylate cyclase 7 (ADCY7) adenylate kinase 2 (AK2) adenylate kinase 3 (AK3) (non-exact, 67%) | 6 | U18121 | | + | + | | + | | |
| adenylate cyclase 7 (ADCY7) adenylate kinase 2 (AK2) adenylate kinase 3 (AK3) (non-exact, 67%) | 2 | AF033861 | | + | + | + | + | + | |
| adenylate kinase 2 (AK2) adenylate kinase 3 (AK3) (non-exact, 67%) | 1 | D25538 | | | | | | | |
| (non-exact, 67%) | 2 | U39945 | | + | + | | + | + | |
| adenylyl cyclase | 1 | X60673 | | | | | | | |
| associated protein (CAP) | 28 | M98474 | T | | + | | + | | |
| adipose differentiation- related protein; adipophilin (ADFP) | 1 | X97324 | | | + | | + | + | |
| ADP-ribosylation factor 1 (ARF1) | 13 | M84326 | | + | + | | + | + | |
| ADP-nbosylation factor 3 (ARF3) | 2 | M33384 | | + | + | | + | | |
| ADP-ribosylation factor 4 (ARF4) | 1 | M36341 | Tlymphoma | + | + | | | + | |
| ADP-ribosylation factor 5 (ARF5) | 1 | M57567 | | | + | + | + | + | |
| ADP-nbosylation factor domain protein 1, 64kD (ARFD1) | 1 | L04510 | | + | | | | | |
| ADP-ribosyltransferase (NAD+; poly (ADP-ribose) polymerase) (ADPRT) | 4 | M32721 | + | + | + | + | + | + | |
| adrenergic, beta, receptor kinase 1 (ADRBK1) | 2 | X61157 | В | + | | | + | | |
| adrenoleukodystrophy-like 1 (ALDL1) | 1 | AJ000327 | | | | | | | |
| AE-binding protein 1 (AEBP1) (non-exact, 62%) | 1 | D86479 | | | | | | | |
| AF-17 | 1 | U07932 | | | | | | | |
| A-gamma-globin | 1 | V00514 | | | | | | | |
| A-gamma-globin (chromosome 11 allele) | 1 | J00176 | | | | | | | |
| agammaglobulinaemia tyrosine kinase (ATK) | 1 | U78027 | | *** | | | | | |
| AHNAK nucleoprotein (desmoyokin) (AHNAK) | 4 | M80899 | + | + | + | + | | + | |
| alanyl (membrane) aminopeptidase (aminopeptidase N, aminopeptidase M, microsomal aminopeptidase, CD13, p150) (ANPEP) | 1 | X13276 | | | + | | + | | |
| alcohol dehydrogenase 5 (class III), chi polypeptide (ADH5) | 1 | M29872 | · | | | | | | |
| aldehyde dehydrogenase 1, soluble (ALDH1) | 1 | AF003341 | | + | | | + | + | |

| WO 00/40749 | | | | | | | | | |
|--|---|----------|---|---|----------|----------|----------|----------|--|
| aldehyde dehydrogenase 10 (fatty aldehyde | 2 | U75286 | | | | | | | |
| dehydrogenase) (ALDH10) aldehyde reductase 1 (low | 3 | J04795 | В | + | + | + | + | | |
| Km aldose reductase) (ALDR1) | | | | | | | | | |
| aldo-keto reductase family 1, member A1 (aldehyde | 2 | J04794 | В | + | + | | + | | |
| reductase) (AKR1A1) aldo-keto reductase family | | D17793 | | + | + | + | | + | |
| 1, member C3 (3-alpha hydroxysteroid dehydrogenase, type II) (AKR1C3) | | | | | | | | | |
| aldo-keto reductase family 7, member A2 (aflatoxin aldehyde reductase) (AKR7A2) | 1 | Y16675 | | + | + | | + | + | |
| aldolase A, fructose- bisphosphate (ALDOA) | 7 | X12447 | | + | + | | + | | |
| aldolase C, fructose- bisphosphate (ALDOC) | 2 | X05196 | | + | + | | + | | |
| alkaline phosphatase, liver/bone/kidney (ALPL) | 1 | 4502062 | | | | | | | |
| ALL-1 (=L04731;L04284 HRX) | 4 | Z69780 | | | | | | | |
| alpha mannosidase II isozyme | 1 | D55649 | | + | | | + | | |
| alpha thalassemia/mental retardation syndrome X- linked (ATRX) | 3 | U75653 | + | + | + | + | | + | |
| alpha-2 macroglobulin | 1 | Z11711 | | | | | | <u> </u> | |
| alpha-2-globin | 2 | V00516 | | | | ļ | <u> </u> | ļ | |
| alpha-2-macroglobulin receptor/lipoprotein receptor protein (A2MR/LRP) | 1 | U06985 | | | | | | | |
| alpha-polypeptide of N- acetyl-alpha- glucosaminidase (HEXA) | 1 | M13520 | | | | | | | |
| alpha-spectrin | 1 | X86901 | | + | | T | | | |
| alpha-subunit of Gi2 a (GTP-binding signal transduction protein) | 1 | X07854 | | | | | | | |
| aminin receptor 1 (67kD); Ribosomal protein SA (LAMR1) | 2 | J03799 | Τ | + | + | | + | + | |
| aminolevulinate, delta-, dehydratase (ALAD) | 1 | X64467 | | + | <u> </u> | <u> </u> | | + | |
| amino-terminal enhancer of split (AES) | | X73358 | + | + | + | + | <u> </u> | | |
| amino-terminal enhancer of split (AES) | | U04241 | В | + | + | <u> </u> | L | + | |
| AMP deaminase isoform L (AMPD2) | 8 | M91029 | | + | | _ | | + | |
| amphiphysin (Stiff-Mann syndrome with breast cancer 128kD autoantigen) (AMPH) | | U07616 | В | + | | | | _ | |
| amphiphysin (Stiff-Mann syndrome with breast cancer 128kD autoantigen) (AMPH)(non-exact, 68%) | | U07616 | | | | | | | |
| amphiphysin (Stiff-Mann syndrome with breast cancer 128kD autoantigen) (AMPH)(non-exact, 68%) | 1 | U07616 | | | | | | | |
| amphiphysin II | 4 | U87558 | | + | + | | + | | |
| amphiphysin II (67%aa | 1 | AF068915 | | | | | | | |
| amphiphysin II (non-exact 69% aa) | 1 | AF001383 | | | | | | | |
| | | | | | | | | | |

| amphiphysin-like (AMPHL) | 1 | U68485 | | + | + | | | | |
|--|----|----------|------------|---|---|---|---|----|------------------------|
| amphiphysin-like (AMPHL) (low match) | 1 | AF068918 | | | | | | | |
| AMY-1 | | D50692 | В, Т | | | | + | | |
| amyloid beta (A4) precursor protein-binding, family B, member 1 (Fe65) (APBB1) | 1 | L77864 | | + | + | + | | + | |
| amyloid beta (A4) precursor-like protein 2 (APLP2) | 6 | L27631 | T lymphoma | + | + | | + | + | |
| ankyrin 3, node of Ranvier (ankyrin G) (ANK) (non- exact, 50%) | 1 | U43965 | | | | | | | |
| annexin I (lipocortin I) (ANX1) | 1 | X05908 | | + | + | + | | + | |
| annexin II | 1 | D28364 | | | | | | | |
| annexin II (lipocortin II; calpactin I, heavy polypeptide) (ANX2) | 7 | D00017 | + | + | + | + | + | | high in many libraries |
| annexin IV (placental anticoagulant protein II) (ANX4) | 1 | M19383 | | + | + | + | + | + | |
| annexin V (endonexin II) (ANX5) | 2 | M21731 | | + | + | + | | + | |
| annexin V (endonexin II) (ANXV) | 1, | M19384 | | + | + | + | | + | |
| annexin VI (p68) (ANX6) | 6 | Y00097 | | + | + | + | | + | |
| annexin VII (synexin) (ANX7) | 1 | J04543 | | + | + | + | | + | |
| antigen identified by monoclonal antibodies 12E7, F21 and O13 (MIC2) | 2 | M16279 | | + | + | + | | + | |
| antigen identified by monoclonal antibodies 4F2, TRA1.10, TROP4, and T43 (MDU1) | 3 | J02939 | | + | + | + | + | + | |
| antigen TQ1 | 1 | | | | | | | | |
| anti-oxidant protein 2 (non- selenium glutathione peroxidase, acidic calcium- independent phospholipase A2) (KIAA0106) | 1 | D14662 | | + | + | + | + | + | |
| APEX nuclease (multifunctional DNA repair enzyme) (APEX) | 5 | X66133 | | + | + | | + | + | |
| Apolipoprotein L (APOL) (59%aa) | 1 | Z82215 | | | | | | | |
| apoptosis inhibitor 1 (API1) | | L49431 | † i | + | + | + | + | + | |
| apoptosis inhibitor 4 (survivin) (API4) | 1 | U75285 | B, W | + | + | | + | | |
| apoptosis inhibitor 5 (API5) | 1 | U83857 | T lymphoma | + | | | + | L_ | |
| apoptosis specific protein (ASP) | 1 | Y11588 | В | + | | | + | + | |
| apoptotic protease activating factor (APAF1) | 1 | AF013263 | В | + | + | | + | | |
| aquaporin 3 (AQP3) | 1 | AB001325 | T | | | | + | | |
| aquaponn 9 (AQP9) | 7 | AB008775 | Tactivated | | | | + | | |
| arachidonate 12- lipoxygenase (ALOX12) | 1 | M58704 | Т | | | | + | + | |
| arachidonate 5- lipoxygenase-activating protein (ALOX5AP) | 3 | X52195 | + | + | | + | | + | |
| ariadne homolog (ARI) | 1 | AJ009771 | + | + | + | + | | + | |
| ariadne-2 (D. melanogaster) homolog (all-trans retinoic acid inducible RING finger) (ARI2) | 1 | AF099149 | + | + | + | + | | + | |

| VV 00/40/49 | | | | | | | | | | |
|--|----|----------|------------------------|---|---|---|---|---|---------------|------|
| ARP1 (actin-related protein 1, yeast) homolog A (centractin alpha) | 1 | X82206 | | + | | | + | | | |
| (ACTR1A) ARP2 (actin-related protein | 9 | AF006082 | | + | + | | + | + | – | |
| 2, yeast) homolog (ACTR2) ARP2/3 protein compex subunit 34 (ARC34) | 5 | AF006085 | T activated, | + | + | | + | | | |
| Arp2/3 protein compex subunit p41 (ARC41) | 6 | AF006084 | monocyte stimulated | + | + | | + | | | |
| Arp2/3 protein compex subunit p41 (ARC41)) (low match) | 1 | AF006084 | | | | | | | | |
| Arp2/3 protein complex subunit p16 (ARC16) | 20 | AF017807 | | + | + | | + | + | | |
| Arp2/3 protein complex subunit p20 (ARC20) | 2 | AF006087 | | + | + | | + | + | | |
| Arp2/3 protein complex subunit p21(ARC21) | 3 | AF006086 | W | | | | + | + | | |
| ARP3 (actin-related protein 3, yeast) homolog (ACTR3) | 11 | AF006083 | W | | + | | + | + | | |
| arrestin, beta 2 (ARRB2) | 1 | AF106941 | B, T, W | + | + | | + | | | |
| arsA (bacterial) arsenite transporter, ATP-binding, homolog 1 (ASNA1) | 1 | AF047469 | В, Т | + | | | + | | | |
| aryl hydrocarbon receptor nuclear translocator-like (ARNTL) | 2 | AF044288 | В | + | + | | + | | | |
| aryl hydrocarbon receptor- interacting protein (AIP) | 1 | U31913 | + | + | + | + | | + | | |
| arylsulfatase A (ARSA) | 1 | X52151 | Tactivated | + | | | + | | | |
| asialoglycoprotein receptor 2 (ASGR2) | 1 | M11025 | | | | | + | + | | |
| asparaginyl-tRNA synthetase (NARS) | 3 | D84273 | | + | + | | + | | | |
| aspartyl-IRNA synthetase (DARS) | 1 | J05032 | В | + | + | | + | | | |
| ataxia telangiectasia mutated (includes complementation groups A, C and D) (ATM) | 1 | U82828 | В, Т | | + | | + | | | |
| ataxin-2-like protein A2LP (A2LG) | 1 | AF034373 | B, T activated | + | + | | | + | | |
| ÀTF6 | 1 | AF005887 | | + | | | + | | | |
| ATP binding cassette transporter (ABCR) (non-exact 80%) | 1 | U88667 | | | | | | | | |
| ATP synthase (F1-ATPase) alpha subunit, mitochondrial | 1 | X59066 | | | | | | | | |
| ATP synthase beta subunit gene | 1 | M19482 | | | | | | | | |
| ATP synthase, H+ transporting, mitochondrial F0 complex, subunit b, isoform 1 (ATP5F1) | 1 | X60221 | + | + | + | + | | + | | |
| ATP synthase, H+ transporting, mitochondrial F0 complex, subunit c (subunit 9), isoform 1 (ATP5G1) | 1 | X69907 | T activated | + | + | | + | + | | |
| ATP synthase, H+ transporting, mitochondrial F1 complex, alpha subunit, isoform 1, cardiac muscle (ATP5A1) | 3 | D14710 | | | | | | | | |
| ATP synthase, H+ transporting, mitochondrial F1 complex, alpha subunit, isoform 1, cardiac muscle (ATP5A1) (low match) | 1 | D14710 | | | | | | | | |

| WU 00/40/49 | | | | | | | | | |
|--|---|----------|------------|---|---|----------|-----|----------|----------------|
| ATP synthase, H+ transporting, mitochondrial | 2 | M27132 | | | | | | | |
| F1 complex, beta polypeptide (ATP5B) | | | |] | | | | | |
| ATP synthase, H+ transporting, mitochondrial F1 complex, gamma polypeptide 1 (ATP5C1) | 1 | D16563 | W | + | + | + | + | | |
| ATP synthase, H+ transporting, mitochondrial F1F0, subunit g (ATP5JG) | 1 | AF092124 | + | + | + | + | + | + | |
| ATP/GTP-binding protein (HEAB) | 2 | U73524 | + | + | + | + | | + | |
| ATPase, Ca++ transporting, ubiquitous (ATP2A3) | 5 | Z69881 | | + | | | | | |
| ATPase, H+ transporting, lysosomal (vacuolar proton pump) 21kD (ATP6F) | 2 | D89052 | + | + | + | + | | + | |
| ATPase, H+ transporting, lysosomal (vacuolar proton pump) 31kD (ATP6E) | 1 | X76228 | | + | + | + | | + | |
| ATPase, H+ transporting, lysosomal (vacuolar proton pump) 42kD; Vacuolar proton-ATPase, subunit C; V-ATPase, subunit C (ATP6D) | 5 | X69151 | | + | + | + | | + | |
| ATPase, H+ transporting, lysosomal (vacuolar proton pump), alpha polypeptide, 70kD, isoform 1 (ATP6A1) | 3 | L09235 | | + | | + | | | |
| ATPase, H+ transporting, lysosomal (vacuolar proton pump), beta polypeptide, 56/58kD, isoform 2 (ATP6B2) | 6 | X62949 | + | + | + | + | | + | |
| ATPase, H+ transporting, lysosomal (vacuolar proton pump), member J (ATP6J) | 2 | AF038954 | + | + | + | + | | + | high in testis |
| ATPase, H+ transporting, lysosomal (vacuolar proton pump), subunit 1 (ATP6S1) | 1 | D16469 | | + | + | + | | + | |
| ATP-binding cassette 50 (TNF-alpha stimulated) (ABC50) | 1 | AF027302 | + | + | + | + | | + | : |
| ATP-binding cassette protein M-ABC1 (mitochondrial) | 1 | AF047690 | | | | | | | , A |
| ATP-dependent RNA helicase | 1 | AJ010840 | Tiymphoma | | + | | + | | |
| autoantigen (Hs.75528) | 2 | L05425 | Tactivated | | + | | | | |
| autoantigen (Hs.75528) (non-exact 84%) | 1 | L05425 | | | | | | | |
| autoantigen (Hs.75682) | 1 | U17474 | В | + | | | | + | |
| autoantigen La/SS-B | 1 | Z35127 | | | | | | | |
| axin (AXIN1) | 1 | AF009674 | Т | + | | <u> </u> | | <u> </u> | |
| axonemal dynein heavy chain (DNAH17) | 1 | AJ000522 | | | | | | + | |
| BAI1-associated protein 3 (BAIAP3) (non-exact 54%) | 1 | AB017111 | | | | | | | |
| basement membrane- | 1 | AF044896 | | | | | | | |
| induced gene (ICB1) | ļ | <u></u> | 1 | | | | | | |
| basic leucine zipper nuclear factor 1 (JEM-1) (BLZF1) | 2 | U79751 | | | | | | | |
| basic leucine zipper nuclear factor 1 (JEM-1) | 5 | X74070 | + | + | + | + | + | + | |
| basic leucine zipper nuclear factor 1 (JEM-1) (BLZF1) basic transcription factor 3 | 2 | | + | + | + | + | + + | + | |

| (zinc finger protein 51) (BCL6) B-cell translocation gene 1, anti-proliferative (BTG) BCL2/adenovirus E1B 1 U15173 B + + + 1 19kD-interacting protein 2 (BNIP2) BCL2/adenovirus E1B 2 AF067396 + + + + + 19kD-interacting protein 3-like (BNIP3L) | WO 00/40/49 | | | | | | | | | |
|--|---|---------------|----------|---|---|---|--|----------|---|------------------------|
| BCL6 Security S | B-cell CLL/lymphoma 6 | 1 | U00115 | | + | + | | | | |
| anti-proliferative (81'G) BCL/2adenovirus E1B 1 | (BCL6) | | | | | | | | | |
| ## 1015173 B * * * * * * * * * * * * * * * * * * | B-cell translocation gene 1. | 1 | X61123 | | | + | | | + | |
| BCL12/adenovirus E1B | BCL2/adenovirus E1B 19kD-interacting protein 2 | 1 | U15173 | В | + | - | | + | + | |
| Section Tocoled-cool, myosin-like BCL2- interacting protein) | BCL2/adenovirus E1B 19kD-interacting protein 3- | 2 | AF067396 | | + | + | + | | + | |
| inferacting protein) (BECN1) bota-1,2-N- acetylglucosaminyltransfer ase il (MGAT2) beta-2-microglobulin (B2M) beta-2-microglobulin (B2M) beta-beta-2-microglobulin (B2M) beta-beta-2-microglobulin (B2M) beta-beta-2-microglobulin (B2M) beta-beta-2-microglobulin (B2M) beta-beta-beta-2-microglobulin (B2M) beta-beta-beta-beta-beta-beta-beta-beta- | beclin 1 (coiled-coil, | 1 | AF077301 | В | + | + | | + | | |
| acetylglucosaminytransfer ase it iMGAT2) Felia-Z-microglobulin (B2M) Fel | interacting protein) (BECN1) | | | | | | | | | |
| Dela-Excosaminidase alpha Chain (HEXM) Esta-Excosaminidase alpha Chain (HEXA) Chain (H | acetylglucosaminyltransfer ase il (MGAT2) | _ | | | | | | | | |
| Chain (HEXA) | beta-2-microglobulin (B2M) | 63 | S82297 | + | + | + | + | + | + | |
| Deta-tubulin (non-exact | chain (HEXA) | 1 | | | | | | | | |
| Test | beta-tubulin | 7 | V00599 | + | + | + | + | + | + | high in many libraries |
| Deta-Libbulin, pseudogene | | 1 | AF070561 | | | | | | | |
| District | | 1 | J00315 | | | | | | | |
| 1 | BING4 | 1 | Z97184 | | | | | | | |
| Distring ase (BTD) (non-exact, 58%) Distring ase (BTD) (non-exact, 58%) Distring ase (BTD) (non-exact, 58%) P43251 P432 | | | U03274 | | | | | | | |
| Diotinidase (BTD) (non-exact, 56%) Diotinidase (BTD) (non-exact, 56%) P43251 | biotinidase (BTD) (non- | 1 | | | | | | | | |
| BIOTINIDASE 1 | biotinidase (BTD) (non- | 1 | U03274 | | | | | | | |
| (serine hydrolase) (BPHL) bone marrow stromal cell antigen 1 (BST1) box-dependent myc- interacting protein isoform BIN1-10 (BIN1) box-dependent myc- interacting protein isoform BIN1-10 (BIN1) (non-exact, 64%) brain my047 protein 1 AF063605 | BIOTINIDASE | 1 | P43251 | | | | | | | |
| antigen 1 (BST1) Dox-dependent myc- interacting protein isoform BIN1-10 (BIN1) Dox-dependent myc- interacting protein isoform BIN1-10 (BIN1) (non-exact, 64%) Drain my047 protein I AF043900 Interacting protein isoform BIN1-10 (BIN1) (non-exact, 64%) Drain my047 protein I AF063605 I + + + Thermoly of the protein isoform BIN1-10 (BIN1) (non-exact, 64%) Drain my047 protein I AF063605 I + + + Thermoly of the protein isoform BIN1-10 (BIN1) (non-exact, 64%) Drain my047 protein I AF063605 I + + + Thermoly of the protein isoform BIN1-10 (BIN1) (non-exact, 64%) Drain my047 protein I AF063605 I + + + Thermoly of the protein isoform BIN1-10 (BIN1) (non-exact, 1 + + Thermoly of the protein isoform BIN1-10 (BIN1) (non-exact, 1 + + Thermoly of the protein isoform BIN1-10 (BIN1) (non-exact, 1 + + Thermoly of the protein isoform BIN1-10 (BIN1) (non-exact, 1 - + + Thermoly of the protein isoform BIN1-10 (BIN1) (non-exact, 1 - + + Thermoly of the protein isoform BIN1-10 (BIN1) (non-exact, 1 - + + Thermoly of the protein isoform BIN1-10 (BIN1) (non-exact, 1 - + + Thermoly of the protein isoform BIN1-10 (BIN1) (BIN10) | biphenyl hydrolase-like (serine hydrolase) (BPHL) | 1 | | | + | | | <u> </u> | | · |
| Dox-dependent myc-interacting protein isoform BIN1-10 (BIN1) Dox-dependent myc-interacting protein isoform BIN1-10 (BIN1) (non-exact, 64%) Brain my047 protein 1 | bone marrow stromal cell | 1 | D21878 | | | | | + | | |
| interacting protein isoform BIN1-10 (BIN1) (non-exact, 64%) brain my047 protein 1 | box-dependent myc- interacting protein isoform BIN1-10 (BIN1) | | | | | | | | | |
| branched chain keto acid dehydrogenase E1, alpha polypeptide (maple syrup urine disease) (BCKDHA) BRCA1 associated protein- 1 (ubiquitin carboxy- terminal hydrolase) (BAP1) BRCA1, Rho7 and vatl genes, and ipf35 breakpoint cluster region protein, uterine leiomyoma, 1; barrier to autointegration factor (BCRP1) breakpoint cluster region protein, uterine leiomyoma, 2 (BCRP2) breast cancer anti-estrogen resistance 3 (BCAR3) (non-exact 73%) bromodomain-containing protein, 140kD (peregrin) (BR140) Bruton's agammaglobulinemia | interacting protein isoform BIN1-10 (BIN1) (non-exact, | 1 | | | | | | | | |
| dehydrogenase E1, alpha polypeptide (maple syrup urine disease) (BCKDHA) BRCA1 associated protein- 1 (ubiquitin carboxy- terminal hydrolase) (BAP1) BRCA1, Rho7 and vatl genes, and ipf35 breakpoint cluster region protein, uterine leiomyoma, 1; barrier to autointegration factor (BCRP1) breakpoint cluster region protein, uterine leiomyoma, 2 (BCRP2) breast cancer anti-estrogen resistance 3 (BCAR3) (non-exact 73%) bromodomain-containing protein, 140kD (peregrin) (BR140) Bruton's agammaglobulinemia | 1 | | | I | | | | Í | | |
| BRCA1 associated protein- 1 (ubiquitin carboxy- terminal hydrolase) (BAP1) BRCA1, Rho7 and vatl genes, and ipf35 breakpoint cluster region protein, uterine leiomyoma, 1; barrier to autointegration factor (BCRP1) breakpoint cluster region protein, uterine leiomyoma, 2 (BCRP2) breast cancer anti-estrogen resistance 3 (BCAR3) (non-exact 73%) bromodomain-containing protein, 140kD (peregrin) (BR140) Bruton's agammaglobulinemia | dehydrogenase E1, alpha | 3 | Z14093 | Т | + | + | | + | | , |
| BRCA1, Rho7 and vatl genes, and ipf35 breakpoint cluster region protein, uterine leiomyoma, 1; barrier to autointegration factor (BCRP1) breakpoint cluster region protein, uterine leiomyoma, 2 (BCRP2) breast cancer anti-estrogen resistance 3 (BCAR3) (non-exact 73%) bromodomain-containing protein, 140kD (peregrin) (BR140) Bruton's agammaglobulinemia | BRCA1 associated protein- 1 (ubiquitin carboxy- | 1 | D87462 | + | + | + | + | | | |
| breakpoint cluster region protein, uterine leiomyoma, 1; barrier to autointegration factor (BCRP1) breakpoint cluster region 2 AF044774 + + + + + + + + + + + + + + + + + | BRCA1, Rho7 and vatl | 1 | L78833 | | | | | _ | | |
| 1; barrier to autointegration factor (BCRP1) breakpoint cluster region protein, uterine leiomyoma, 2 (BCRP2) breast cancer anti-estrogen resistance 3 (BCAR3) (non-exact 73%) bromodomain-containing protein, 140kD (peregrin) (BR140) Bruton's agammaglobulinemia | breakpoint cluster region | 2 | AF044773 | | + | + | - | - | | |
| breakpoint cluster region protein, uterine leiomyoma, 2 (BCRP2) breast cancer anti-estrogen resistance 3 (BCAR3) (non-exact 73%) bromodomain-containing protein, 140kD (peregrin) (BR140) Bruton's agammaglobulinemia | 1; barrier to autointegration | | | | | | | | | |
| breast cancer anti-estrogen 1 U92715 resistance 3 (BCAR3) (non-exact 73%) bromodomain-containing 2 M91585 + protein, 140kD (peregrin) (BR140) Bruton's 1 U13424 agammaglobulinemia | breakpoint cluster region protein, uterine leiomyoma, | 2 | AF044774 | | + | + | | + | + | |
| bromodomain-containing 2 M91585 + protein, 140kD (peregrin) (BR140) Bruton's 1 U13424 agammaglobulinemia | breast cancer anti-estrogen resistance 3 (BCAR3) | 1 | U92715 | | | | | | | |
| Bruton's 1 U13424 | bromodomain-containing protein, 140kD (peregrin) | 2 | | | + | | | | | |
| | Bruton's agammaglobulinemia | 1 | U13424 | | | | | | | |

| 1 | U78027 | | | | | | | |
|---|------------|------|----------|----------|----------|---|----------|---|
| 1 | U78027 | | | - | | | | |
| 1 | | | | | | | | |
| 6 | Y09943 | + | + | + | + | | + | |
| 1 | U78027 | + | 1 + 1 | + | + | | + | |
| 1 | U01923 | | + | + | | + | | |
| 4 | AF053304 | + | + | + | + | | + | |
| · | | + | | | + | | + | |
| 7 | U90543 | | + | + | | + | | |
| 1 | AB020625.1 | | | | | | | |
| 2 | U80744 | | + | + | | | | |
| 2 | 1 | | + | | | + | | |
| 1 | | | | | | | | |
| 1 | | | | | + | | + | |
| | | В | + | + | | + | | |
| | | | | | | | | |
| | | В | | + | | | | |
| | | Т | + | <u> </u> | | | | |
| 5 | X04366 | | + | + | | + | + | |
| 5 | M23254 | | + | + | İ | | | |
| 1 | X04106 | | + | + | | + | + | |
| 3 | D16217 | | | | | + | | |
| 2 | D83735 | | + | | + | | + | |
| 1 | D83735 | В, Т | + | | | + | | |
| 1 | D83735 | | | | | | | |
| 3 | AF013759 | В | | + | | + | + | |
| 4 | L05912 | | | | | | | |
| 1 | M90360 | | | | | | | |
| 1 | | | | | | + | | |
| | | | <u> </u> | | | | _ | |
| | | | | | <u> </u> | _ | <u> </u> | |
| 1 | U03271 | + | | | _ | | | |
| | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |

| capping protein (actin filament), gelsolin-like (CAPG) | 8 | M94345 | + | + | | + | | + | |
|---|----|------------|---------------------|--------|------------|------|---|---|--|
| carbamoyl-phosphate synthetase 2, aspartate transcarbamylase, and dihydroorotase (CAD) | 1 | D78586 | + | + | + | + | | + | |
| carbonic anhydrase V, mitochondrial (CA5) | 1 | L19297 | | + | | | + | | |
| carboxypeptidase D (CPD) | 3 | U65090 | В | + | + | - | | | |
| camitine/acylcamitine | 1 | Y10319 | | + | + | | + | | |
| translocase (CACT) Cas-Br-M (murine) | 2 | X57110 | | | | | + | | |
| ecotropic retroviral transforming sequence (cbl) | 2 | | | | | | , | | |
| casein kinase 1, alpha 1 (CSNK1A1) | 1 | L37042 | + | + | +" | + | | + | |
| casein kinase 2, alpha 1 polypeptide (CSNK2A1) | 2 | M55265 | В | + | | | + | + | |
| casein kinase I gamma 3L (CSNK1G3L) | 1 | AF049090.1 | | | | | | | |
| casein kinase II alpha | 1 | X69951 | | | | | | | |
| subunit(=S72393) CASP8 and FADD-like | 4 | AF015450 | | + | + | + | + | + | |
| apoptosis regulator (CFLAR) | • | | | | | | | | |
| caspase 1, apoptosis- related cysteine protease (interleukin 1, beta, | 7 | U13697 | + | | | + | | | |
| convertase) (CASP1) caspase 10, apoptosis- related cysteine proteas (CASP10) | 1 | U60519 | B, T activ lymph | | Т | | + | | |
| caspase 3, apoptosis- related cysteine protease (CASP3) | 3 | U13737 | В, Т | + | + | + | + | | |
| caspase 4, apoptosis- related cysteine protease (CASP4) | 6 | U25804 | + | + | + | + | | + | |
| Caspase 5, apoptosis- related cysteine protease (CASP5) | 1 | U28015 | | · | + | | | | |
| caspase 8, apoptosis- related cysteine protease (CASP8) | 2 | X98173 | | + | | + | | + | |
| caspase 9, apoptosis- related cysteine protease (CASP9) | 1 | U56390 | В | | | + | + | | |
| catalase (CAT) | 5 | X04076 | В | + | + | | + | | |
| catechol-O- methyltransferase (COMT) | - | M65213 | | + | + | | + | | |
| catenin (cadherin- associated protein), alpha 1 (102kD) (CTNNA1) | 6 | D14705 | | + | + | | | | |
| cathelicidin antimicrobial peptide (CAMP) | 1 | X89658 | В | | | | | | |
| cathepsin B (CTSB) | 4 | L16510 | | 1 | + | | + | + | |
| cathepsin C (CTSC) | 3 | U79415 | | + | + | + | | + | |
| cathepsin D (lysosomal aspartyl protease) (CTSD) | 4 | M11233 | | + | + | | + | | |
| cathepsin E (CTSE) | 1 | J05036 | | | | | + | | |
| cathepsin G (CTSG) | 1 | M16117 | T, W | | + | | | | |
| cathepsin S (CTSS) | 34 | M86553 | B, Monocyt | e stim | ulate a | d, T | + | + | |
| cathepsin W (lymphopain) (CTSW) | 4 | AF013611 | | | | | | + | |
| CBF1 interacting corepressor CIR (=U03644 recepin) | | AF098297 | | | | | | | |

| CCAAT/enhancer binding protein (C/EBP), alpha (CEBPA) | 3 | X87248 | | + | + | + | | + | |
|--|------|----------|--|---|---------------------------------------|----------|----------|---|---|
| CCAAT/enhancer binding protein (C/EBP), delta | 1 | S63168 | | | + | | + | + | |
| (CEBPB) CCAAT-box-binding | 2 | M37197 | T lymphoma | | - | + | + | | |
| transcription factor (CBF2) CCR5 receptor (CCR5) | 1 | AF011504 | | | · · · · · · · · · · · · · · · · · · · | | | | |
| (non-exact?) CD14 antigen (CD14) | - 11 | M86511 | + | + | + | + | | + | |
| CD18 (=M95293) | 4 | X64071 | | | | - | | _ | |
| CD1C antigen, c | 2 | M28827 | | | | | | + | |
| polypeptide (CD1C) CD2 antigen (cytoplasmic | 1 | AF104222 | | | | | | | |
| tail)-binding protein 2 (CD2BP2) | | | | | | | | + | |
| CD2 antigen (p50), sheep red blood cell receptor (CD2) | 4 | M14362 | + | | + | + | + | T | |
| CD2 cytoplasmic tail- binding protein 1 (CD2BP1) | 2 | AF038602 | | | | | T | | |
| CD20 antigen (CD20) | 1 | X12530 | | | | | | | |
| CD20 receptor (S7) | | X07203 | <u> </u> | | | | Щ | | |
| CD22 antigen (CD22) | 1 | U62631 | В | | | | | | |
| CD24 signal transducer | 7 | M58664 | <u> </u> | | | <u> </u> | | | |
| CD33 antigen (gp67) (CD33) | 1 | M23197 | ļ | | | | + | | |
| CD33 antigen-like 2; OB binding protein-2 (CD33L2) (non-exact, 68%) | 1 | U71383 | | | | | İ | | |
| CD33L2 (61% aa) | 1 | D86359 | | | | <u></u> | | | |
| CD36 antigen (collagen type I receptor, thrombospondin receptor) (CD36) | 7 | M98398 | Tlymphoma | | + | | + | + | |
| CD37 antigen (CD37) | 5 | X14046 | + | + | | + | | + | |
| CD38 alt | 1 | D84277 | | | | | | | |
| CD39 antigen (CD39) | 1 | U87967 | В | + | | T | + | + | - |
| CD3D antigen, delta polypeptide (TiT3 complex) (CD3D) | . 1 | X03934 | | | + | + | | + | |
| CD3E antigen, epsilon polypeptide (TiT3 complex) (CD3E) | 1 | X03884 | + | | | + | | | |
| CD3G antigen, gamma polypeptide (TiT3 complex) (CD3G) | 2 | X06026 | W | | | | + | | |
| CD3Z antigen, zeta polypeptide (TiT3 complex) (CD3Z) | 2 | J04132 | + | | | + | | | |
| CD3-zeta (clone pBS NK1) | 1 | X55510 | | | | | | | |
| CD4 (low match) | 1 | S68043 | | | | | | | |
| CD4 antigen (p55) (CD4) | 4 | M12807 | | + | + | | + | | |
| CD44 antigen (homing function and Indian blood group system (CD44) | 6 | X56794 | W | | | | + | + | |
| CD48 antigen (B-cell membrane protein) (CD48) | 3 | X06341 | + | + | + | + | | + | |
| CD53 antigen (CD53) | 10 | L11670 | + | + | | + | | + | |
| CD53 antigen (CD53) (low match) | 1 | M60871 | | | | | | | |
| CD63 antigen (melanoma 1 antigen) (CD63) | | M59907 | | | | | | | |
| CD68 antigen (CD68) | 2 | S57235 | | + | + | | + | + | |

| polypeptide of major histocompatibility complex, class II antigen-associated) (CD74) CD794 antigen | WO 00/40/49 | | | | | | | | | |
|---|---|-----|----------|-------------|---|---|---|---|---|------------------------|
| CD794 sartigen | polypeptide of major histocompatibility complex, class II antigen-associated) | 72 | K01144 | + | + | + | + | + | + | high in many libraries |
| CD798 antigen | CD79A antigen | 2 | M80462 | | | + | | | | |
| CDB antigen, alpha | CD79B antigen (immunoglobulin- | 2 | М89957 | + | | | | | | |
| CD8 antigen (cd8) | CD8 antigen, alpha | 2 | M27161 | + | | | + | | + | |
| CD81 antigen (activated B 1 CD1151 B + + + + + + + + + + | CD8 antigen, beta polypeptide 1 (p37) | 1 | X13445 | W | | | | | | |
| CDB3 antigen (activated B 1 C01151 B + + + + + | CD81 antigen (target of antiproliferative antibody 1 | 1 | M33680 | | + | + | | | + | |
| CD84 antigen (leukocyte antigen) (CD84) | CD83 antigen (activated B lymphocytes, mmunoglobulin | 1 | Q01151 | В | | + | | | | |
| CD8 antigen (p24) (CD9) | CD84 antigen (leukocyte | 1 | U82988 | | | + | | | + | |
| CD97 antigen (CD97) 12 X84700 + + + + CD97 antigen (CD97) 12 X84700 + + + + CD97 antigen (CD97) 1 P48960 1 | | 1 | L25259 | | + | l | | L | L | |
| CD97 antigen (CD97) | CD9 antigen (p24) (CD9) | 2 | M38690 | | | + | | + | + | |
| Description CD97 CD97 CD97 CD97 CD97 antigen (CD97) (non-exact 59%) CD97 antigen (CD97) (non-exact 62%) CD23 (cell division cycle 1 | CD97 antigen (CD97) | 12 | X84700 | + | + | | + | | | |
| CD97 antigen (CD97) (non-exact 62%) | CD97 antigen (CD97) (noin-exact 59%) | 1 | P48960 | | | | | | | |
| CDC23 (cell division cycle 1 | | - 1 | X94630 | + | + | | + | | | |
| CDC37 homolog | CDC23 (cell division cycle 23, yeast, homolog) | 1 | AF053977 | | + | | | + | + | |
| Cdc42 effector protein 3 | | 1 | U63131 | В | + | + | | + | + | |
| CDC-like kinase (CLK) | Cdc42 effector protein 3 | 2 | AF104857 | В | + | + | | + | | ** |
| CDUS2 antigen 13 | | 1 | L29219 | | + | + | + | | + | |
| (CMPATH-1 antigen) (CDW52) cell cycle progression restoration 8 protein(CPR8) cell division cycle 10 (homologous to CDC10 of S. cerevisiae) (CDC10) cell division cycle 20, S. cerevisiae homolog (CDC20) cell division cycle 25B (CDC25B) cell division cycle 2-like 1 (CPITSLE proteins) (CDC2L1) (non-exact 42%) cell division cycle 42 (GTP-binding protein, 25kD) (CDC42) cell division protein (non-exact 68%) CELL-CYCLE NUCLEAR AUTOANTIGEN SG2NA (S/G2 NUCLEAR ANTIGEN) centromere protein B (B0kD) (CENPB) cep250 centrosome 1 AF01794 AF01794 + + + + + + + + + + + + + + + + + + + | CDC-like kinase 2 (CLK2) | 1 | AF023268 | В | + | + | П | | | |
| restoration 8 protein(CPR8) | (CAMPATH-1 antigen) (CDW52) | | | T activated | + | + | | + | | |
| Cell division cycle 20, CDC10 Cell division cycle 20, CDC20 Cell division cycle 25B Cell division cycle 25B Cell division cycle 25B Cell division cycle 2-like 1 AF087514 CDC25B) Cell division cycle 2-like 1 AF087514 CPITSLRE proteins) CDC2L1) (non-exact 42%) Cell division cycle 42 (GTP-binding protein, 25kD) CDC42) Cell division protein (non-exact 68%) CELL-cycle NUCLEAR AF063015 CELL-cycle NUCLEAR AUTOANTIGEN SG2NA (S/G2 NUCLEAR ANTIGEN) Centromere protein B 1 X55039 + | restoration 8 protein(CPR8) | | | | | | | | L | |
| Cell division cycle 20, 1 | (homologous to CDC10 of | 4 | S72008 | + | + | † | + | | _ | |
| Cell division cycle 25B | cell division cycle 20, S.cerevisiae homolog | 1 | U05340 | | + | + | + | | | |
| Cell division cycle 2-like 1 | cell division cycle 25B | 6 | Z68092 | + | + | + | + | | + | |
| Cell division cycle 42 (GTP-binding protein, 25kD) (CDC42) | cell division cycle 2-like 1 (PITSLRE proteins) | 1 | AF067514 | | | | | | | |
| Cell division protein (non- exact 68%) | cell division cycle 42 (GTP- binding protein, 25kD) | 5 | M35543 | + | + | + | + | | + | |
| CELL-CYCLE NUCLEAR | cell division protein (non- | 1 | AF063015 | | | | | | | |
| Centromere protein B | CELL-CYCLE NUCLEAR AUTOANTIGEN SG2NA (S/G2 NUCLEAR | 1 | Q13033 | | | | | | | |
| cep250 centrosome 3 AF022655 B + + + | centromere protein B | 1 | X55039 | | + | | | + | | |
| | cep250 centrosome | 3 | AF022655 | В | + | | T | + | | |

| W O 00/40/43 | | | | | | | | | |
|---|----------|------------|--|----------|--------------|------------|-------------|--------------|--------------|
| ceroid-lipofuscinosis, | 7 | AF017456 | + | + | + | + | + | + | high in bone |
| neuronal 2, late infantile | | | | | | | | | |
| (Jansky-Bielschowsky | | | | | | | | | |
| disease) (CLN2) | | | | | | | | | |
| c-fgr (=M63877 | 6 | X52206 | | | | | | | 1 |
| nonreceptor protein- | | | | | | | | | |
| tyrosine kinase (fgr)) | | | | | | | | | |
| CGI-19 protein | 3 | AF132953.1 | | | | | | | j |
| chaperonin containing | ·····1 | X74801 | | + | + | | | + | |
| TCP1, subunit 3 (gamma) | | | | 1 | | | | | 1 1 |
| (CCT3) | | | | | | | | | |
| chaperonin containing | 1 | AF026291 | Ì | + | + | | + | + | |
| TCP1, subunit 4 (delta) | | 1 | | | į | | | ı | |
| (CCT4) | | | | | | | | | |
| chaperonin containing | 4 | L27706 | В | + | + | | | | |
| TCP1, subunit 6A (zeta 1) | | | | | | 1 | | 1 | Į i |
| (CCT6A) | | | | | | | | | |
| chaperonin containing | 4 | AF026292 | 8 | + | | | | + | |
| TCP1, subunit 7 (eta) | | | | | | | | | |
| (CCT7) | | | | | | | L | | |
| Chediak-Higashi syndrome | 1 | U67615 | B, T | + | + | | + | 1 | |
| 11 (CHS1) | | | lymphoma | | | | | <u> </u> | |
| Chediak-Higashi syndrome | 1 | U67615 | | | | | | | |
| 1 (CHS1) (low score) | | <u></u> | | | | | | <u> </u> | <u> </u> |
| chemokine (C-C motif) | 4 | U03905 | | | | | | |] |
| receptor 2 (CCR2) | | | L | <u> </u> | <u> </u> | | | | |
| chemokine (C-C motif) | 1 | X85740 | | 1 | | | l | | |
| receptor 4 (CCR4) (low | | | | 1 | | | | | |
| match) (may contain | | } | | l | | | | | |
| repeat) | | 1 24504 | ļ | | ļ | | - | | |
| chemokine (C-C motif) | 6 | L31581 | | | 1 | 1 | ļ | | 1 |
| receptor 7 (CCR7) | | LIDASEA | | + | | ⊢ | - | | |
| chemokine (C-X3-C) | 5 | U20350 | | * | į . | 1 | ŀ | | |
| receptor 1 (CX3CR1) | | M99293 | + | + | + | + | | + | |
| chemokine (C-X-C motif), | 5 | M99293 | T | T | * | T . | | T | Į. |
| receptor 4 (fusin) (CXCR4) | 2 | M80927 | ļ | + | ├ | + | - | + | |
| chitinase 3-like 1 (cartilage | | IVIOU827 | I | " | İ | ' | | Ι' | |
| glycoprotein-39) (CHI3L1) | 2 | U49835 | | + | | + | - | + | |
| chitinase 3-like 2 (CHI3L2) | _ | | | <u> </u> | <u> </u> | L. | | Ļ. | |
| chloride channel 1, | 1 | G18280 | | ļ | | | | 1 | |
| skeletal muscle (CLCN1) | | | | └ | | <u> </u> | | ļ | |
| chloride channel 6 | 1 | D28475 | | + | + | | | | |
| (CLCN6) | | | | + | + | + | | + | ļ |
| Chloride intracellular | 1 | U93205 | + | T | T . | T | ļ | * | 1 |
| channel 1 (CLIC1) | | V45000 | | ļ | + | | | ├ | |
| chondroitin sulfate | 5 | X15998 | | İ | * | i | | | |
| proteoglycan 2 (versican) | | | | l | | | 1 | | |
| (CSPG2) | | J02814 | ļ | | + | | | + | |
| chondroitin sulfate | 2 | JU2814 | | ì | * | | l | • | 1 |
| proteoglycan core protein | | Q09028 | | - | | - | | | |
| chromatin assembly factor | • | 203020 | | | 1 | 1 | l | 1 | |
| 1 p48 subunit (CAF-1 P48 subunit) (retinoblastoma | | | | 1 | | | l | 1 | |
| | | | | ļ | i | | l | | |
| binding protein p48) (retinoblastoma-binding | | | 1 | | | | l | 1 | |
| protein 4) (MSI1 protein | ì | 1 | 1 | | | | l | | |
| (homolog) | | | 1 | 1 | | 1 | 1 | | |
| chromodomain helicase | 2 | AF006513 | † | 1 | | 1 | | | |
| DNA binding protein 1 | _ | | 1 | 1 | | İ | l | | |
| (CHD1) | | | | | | | 1. | | L |
| chromodomain helicase | 1 | AF054177 | | | | | <u> </u> | Π | |
| DNA binding protein 1-like | | 1 | 1 | | | 1 | | 1 | |
| (CHD1L) | | | 1 | L . | <u> </u> | <u>L</u> _ | <u></u> | L_ | |
| chromodomain helicase | 1 | AF006514 | В | + | + | | + | | |
| DNA binding protein 2 | 1 | | 1 | 1 | | 1 | l | | |
| (CHD2) | | 1 | <u> </u> | | L | <u>L</u> _ | L | 1_ | |
| chromodomain helicase | 1 | AF006515 |] | | | | | | |
| DNA binding protein 3 | | | | 1 | | 1 | 1 | 1 | |
| (CHD3) | | | | L | L | | <u> </u> | ـــــ | <u> </u> |
| chromodomain helicase | 5 | X86691 | + | + | + | + | 1 | + | |
| DNA binding protein 4 | | | 1 | 1 | } | | 1 | 1 | |
| (CHD4) | <u> </u> | <u></u> | <u> </u> | <u> </u> | <u> </u> | <u> </u> | Ц_ | 1 | <u></u> |
| | | | | | | | | | |

| WO 00/40749 | | | | | | | | | |
|---|---|-------------|--------------|--------------|---|---|---|---|--|
| chromosome 1 open reading frame 7 (C1ORF7) | 1 | AF054176 | | | | | | | |
| chromosome 1 specific transcript KIAA0493 | 1 | AB007962 | | | | | | [| |
| chromosome 17 open reading frame 1B (C17ORF1B) | 1 | AJ008112 | | + | | | | | |
| chromosome 4 open reading frame 1 (C4ORF1) | 1 | AF006621 | | + | + | + | | + | |
| chromosome condensation 1-like (CHC1L) | 2 | AF060219 | | + | + | + | | + | |
| chromosome X open reading frame 5 (CXORF5) | 1 | Y15164 | В | + | + | | + | | |
| chromosome-associated polypeptide C(CAP-C) | 2 | AF092564 | В | + | + | | + | + | |
| cig42 | | AF026944 | | | | | - | | |
| cig5 | 3 | AF026941 | | 1 - | | | | | |
| citrate synthase (CS) | 2 | AF047042 | В | +++ | + | | + | + | |
| | | U31372 | | 1 | | - | | | |
| class I major histocompatibility antigen (HLA-Cw3) | 2 | | | | | | | | |
| class I major histocompatibility antigen (HLA-Cw3) (low match) | 1 | U31372 | | | _ | | | | |
| clathrin assembly protein lymphoid myeloid leukemia (CALM) | 3 | U45976 | В | + | + | | | + | |
| clathrin heavy chain | 1 | X55878 | | 1 1 | | | | | |
| clathrin, heavy polypeptide- like 2 (CLTCL2) | 1 | D21260 | | | | | | | |
| clathrin, light polypeptide (Lca) (CLTA) (low match) | 1 | M20472 | | | | | | | |
| clathrin- | 3 | D63475 | _ | ++ | + | + | + | + | |
| associated/assembly/adapt or protein, medium 1 (CLAPM1) | J | | | | | | | | |
| cleavage stimulation factor, 3' pre-RNA, subunit 2 64kD (CSTF2) (non-exact 82%) | 1 | M85085 | | | | | + | | |
| Cleavage stimulation factor, 3' pre-RNA, subunit 3, 77kD (CSTF3) | 1 | U15782 | 8 | + | + | | * | | |
| clk3 | 1 | L29220 | В | + | + | | | | |
| clone 23815 (Hs.82845) | | U90916 | | + | + | 1 | | + | |
| clone 24592 mRNA | 1 | D88378 | + | + | + | + | | + | |
| clq/MBL/SPA receptor | 1 | U94333 | | | | | | | |
| C1qR(p) () Clusterin (complement lysis inhibitor, SP-40,40, sulfated glycoprotein 2, testosterone-repressed prostate message 2, apolipoprotein J) (CLU) | 1 | M64722 | + | + | + | + | + | + | |
| CMP-sialic acid transporter (CMPST) | 1 | D87969 | В | + | + | | | | |
| CMRF35 | 3 | X66171 | | | | | | 1 | |
| c-myc oncogene containing | 1 | X54629 | | | | | Г | Γ | |
| coagulation factor II (thrombin) receptor (F2R) | 1 | M62424 | | + | + | | | + | |
| coagulation factor V (proaccelerin, labile factor) (F5) | 1 | M14335 | | + | | + | + | | |
| coagulation factor XIII a | 3 | M21998 | | | | | | | |
| coagulation factor XIII, A1 polypeptide (F13A1) | 6 | M14354 | | + | + | + | | + | |
| coated vesicle membrane protein (RNP24) | 1 | X92098 | + | + | + | + | + | + | |
| E | | | | | | | | | |

| WO 00/40749 | | | | | | | | | |
|---|----|----------|------|---|---|----------|---|---|---------------------|
| coatomer protein complex, subunit alpha (COPA) | 5 | U24105 | T | + | | | + | | |
| Cofilin 1 (non-muscle) (CFL1) | 13 | X95404 | + | + | + | + | + | + | high in fetal brain |
| cold inducible RNA-binding protein (CIRBP) | 7 | D78134 | | + | + | | | + | |
| cold shock domain protein A (CSDA) | 3 | X95325 | | + | + | | | | |
| collagen, type IX, alpha 2 (COL9A2) | 3 | AF019406 | В | | | | | | |
| colony stimulating factor 1 receptor, formerly McDonough feline sarcoma viral (v-fms) oncogene homolog (CSF1R) | 3 | X03663 | | + | | | + | + | |
| colony stimulating factor 2 receptor, beta, low-affinity (granulocyte-macrophage) (CSF2RB) | 5 | M59941 | | | | | | | · |
| colony stimulating factor 2 receptor, beta, low-affinity (granulocyte-macrophage) (CSF2RB) (low match) | 1 | M59941 | | | | | | | |
| colony stimulating factor 3 receptor (granulocyte) (CSF3R) | 16 | X55720 | | + | | | | | |
| complement component 5 receptor 1 (C5a ligand) (C5R1) | 1 | M62505 | L | | | | | | |
| conserved gene amplified in osteosarcoma (OS4) | 2 | AF000152 | | + | + | + | | + | |
| COP9 (constitutive photomorphogenic, Arabidopsis, homolog) subunit 3 (COPS3) | 2 | AF031647 | | + | + | | | + | |
| COP9 homolog (HCOP9) | 2 | U51205 | В | + | + | + | + | + | |
| COPII protein, homolog of s. cerevisiae SEC23p (SEC23A) | 4 | X97064 | | + | + | | | | |
| copine I (CPNE1) | 2 | U83246 | В | + | + | | + | | |
| copine I (CPNE1) (low score) | 1 | U83246 | | | | ļ | | | |
| coproporphyrinogen oxidase (coproporphyria, harderoporphyria) (CPO) | 1 | D16611 | | | + | | + | + | |
| core-binding factor, beta subunit (CBFB) | 1 | L20298 | | + | | | | | |
| coronin | 22 | X89109 | T, W | + | + | | + | | |
| coronin (low match) | 1 | U34690 | | | | | | | |
| coronin (non-exact, 71%) | 1 | X89109 | | | | | | | |
| cot (cancer Osaka thyroid) oncogene (COT) | 1 | D14497 | + | + | + | + | | + | |
| cryptochrome 1 (photolyase-like) (CRY1) | | D84657 | | + | + | <u> </u> | | + | |
| CTD (carboxy-terminal domain, RNA polymerase III, polypeptide A) phosphatase, subunit 1 (CTDP1) | 1 | AF081287 | | + | + | + | | + | |
| C-terminal binding protein 1 (CTBP1) | 1 | U37408 | В | + | + | | + | | |
| C-terminal binding protein 2 (CTBP2) | 2 | AF016507 | | + | + | | + | | |
| CUG triplet repeat, RNA- binding protein 1 (CUGBP1) | 3 | U63289 | | + | + | + | | + | |
| cullin 1 (CUL1) | 3 | U58087 | | + | + | + | | + | |
| cullin 3 (CUL3) | 2 | U58089 | | + | + | + | | + | |
| cut (Drosophila)-like 1 (CCAAT displacement protein) (CUTL1) | 1 | M74099 | В | + | | | | | |
| | | 2 | | | | | | | |

| WO 00/40/49 | | | | | | | | | |
|--|-------------|----------|------------------|----------|----------|----|----------|----------|--|
| cyclin D2 (CCND2) | 2 | D13639 | | + | + | + | | + | |
| cyclin D3 (CCND3) | 5 | M92287 | В, Т | - | + | | + | | |
| | | | lymphoma | + | + | - | | + | |
| cyclin G1 (CNNG1) | 1 | D78341 | В | | | | | _ | |
| cyclin I | 3 | D50310 | В | + | | لط | + | | |
| cyclin T2 (CNNT2) | 1 | AF048732 | B, T lymphoma | В | | | | | |
| cyclin-dependent kinase 2 (CDK2) | 1 | X62071 | | | | | | | |
| cyclin-dependent kinase inhibitor (p27Kip1) | 1 | S76986 | | | | | | | |
| cyclin-dependent kinase inhibitor 1A (p21, Cip1) (CDKN1A) | 2 | 567388 | + | + | + | + | + | + | |
| CYP2D7-CYP2D6 intergenic region (partial) | 1 | X90926 | | | | | | | |
| cystatin B (stefin B) (CSTB) | 1 | L03558 | | | + | | + | + | |
| cysteine and glycine-rich protein 3 (cardiac LIM protein) (CSRP3) | 5 | L54057 | | | + | | | | |
| cytidine deaminase (CDA) | 2 | L27943 | | | | | + | <u> </u> | |
| cytochrome b | 1 | AF042500 | | | | | | | |
| cytochrome b (CYTB) (isolate Aus5) | 1 | AF042518 | | | | | | | |
| cytochrome b(-245) beta chain N-terminal region (X- linked granulomatous disease gene) | 2 | X05895 | | | | | | | |
| cytochrome b-245, beta polypeptide (chronic granulomatous disease) (CYBB) | 2 | X04011 | + | | | + | | + | |
| cytochrome C | 1 | P00001 | | | | | | | |
| cytochrome c oxidase subunit IV (COX4) | 1 | U90915 | T | + | + | | + | + | |
| cytochrome c oxidase subunit Vb (COX5B) | 2 | M59250 | | | | | + | | |
| cytochrome c oxidase subunit VII-related protein (COX7RP) | 6 | AB007618 | + | + | + | + | | + | |
| cytokine suppressive anti- inflammatory drug binding protein 1 (p38 MAP kinase) (CSBP1) | 1 | L35263 | lymphocyte | + | + | | + | | |
| Cytoplasmic antiproteinase=38 kda intracellular serine proteinase inhibitor | 1 | S69272 | | | + | | | | |
| cytotoxic granule- associated RNA-binding protein p40-TIA-1 | 1 | 570114 | | | | | : | | |
| D123 (D123) | 1 | D14878 | + | + | | + | | + | |
| D2-2 | 1 | AF019226 | T | | | 1 | | | |
| D38 | 1 | X74802 | | - | 1 | 1 | <u> </u> | | |
| damage-specific DNA binding protein 1 (127kD) | 2 | AJ002955 | + | + | + | + | + | + | |
| (DDB1) DCHT (low match) | | AF017635 | | _ | _ | | _ | - | |
| | | U78524 | | + | + | + | + | + | |
| DEAD/H (Asp-Glu-Ala- Asp/His) box binding protein 1 (DDXBP1) | | | | | <u> </u> | Ĺ | | | |
| DEAD/H (Asp-Glu-Ala- Asp/His) box polypeptide (72KD) (P72) | 2 | U59321 | T | + | + | | +. | + | |
| DEAD/H (Asp-Glu-Ala- Asp/His) box polypeptide 1 (DDX1) | 1 | X70649 | | + | + | | | Ť | |

| WO 00/40/49 | | | | | | | | | .I/CA | | |
|--|----|----------|---------------------------|-----|---|---|---|---|-------|------|--|
| DEAD/H (Asp-Glu-Ala- Asp/His) box polypeptide 15 (DDX15) | 2 | AB001636 | | | | | | | | | |
| DEAD/H (Asp-Glu-Ala- Asp/His) box polypeptide 16 (DDX16) | 2 | AB011149 | + | + | + | + | | + | | | |
| DEAD/H (Asp-Glu-Ala- Asp/His) box polypeptide 3 (DDX3) | 3 | U50553 | + | + " | + | + | | + | | | |
| DEAD/H (Asp-Glu-Ala- Asp/His) box polypeptide 5 (RNA helicase, 68kD) (DDX5) | 37 | X15729 | + | + | + | + | | + | | | |
| DEAD/H (Asp-Glu-Ala- Asp/His) box polypeptide 5 (RNA helicase, 68kD) (DDX5) (low match) | 1 | AF015812 | | | | | | | | | |
| DEAD/H (Asp-Glu-Ala- Asp/His) box polypeptide 6 (RNA helicase, 54kD) (DDX6) | 2 | D17532 | + | + | | | | | _ | | |
| DEAD/H (Asp-Glu-Ala- Asp/His) box polypeptide 8 (RNA helicase, 54kD) (DDX8) | 1 | D50487 | | + | + | + | | + | | | |
| DEAD/H (Asp-Glu-Ala- Asp/His) box polypeptide 9 (RNA helicase A, nuclear DNA helicase II; lleukophysin) (DDX9) | 3 | L13848 | + | + | + | + | | + | | | |
| DEAD/H (Asp-Glu-Ala- Asp/His) box polypeptide, Y chromosome (DBY) | 1 | AF000985 | | + | + | | + | | | | |
| Death associated protein 3 (DAP3) | 2 | X83544 | +. | + | + | + | + | + | | | |
| death effector domain- containing protein (DEDD) | 1 | AF083236 | | + | + | | | + | | | |
| death-associated protein 6 (DAXX) | 2 | AF039136 | | + | + | + | | + | | | |
| dedicator of cyto-kinesis 2 (DOCK2) | 4 | D86964 | + | + | | + | | + | | : | |
| defender against cell death 1 (DAD1) | 1 | D15057 | | | + | | + | + | | | |
| Defensin, alpha 1, myeloid- related sequence (DEFA1) | 4 | L12690 | | | | + | + | + | | | |
| DEK gene (D6S231E) | 1 | X64229 | В | | + | | + | | | | |
| delta sleep inducing peptide, immunoreactor (DSIPI) | 4 | Z50781 | + | + | + | + | | + | | | |
| dendritic cell protein (GA17) | 3 | AF064603 | + | + | + | + | | + | | | |
| deoxycytidine kinase (DCK) | 1 | M60527 | | | | | | | | | |
| deoxyribonuclease II, lysosomal (DNASE2) | 3 | AB004574 | | | | | | | | | |
| ĎGS-I | 2 | L77566 | | + | | | | L | | | |
| diacylglycerol kinase | 3 | D16440 | 1 | | | | | | | | |
| diacylglycerol kinase alpha (DAGK1) (clone 24) | 3 | AF064771 | | + | | | | | | | |
| diacylglycerol kinase alpha (DAGK1) (clone 24) (low match) | 1 | AF064771 | | | | | | | | | |
| diaphanous (Drosophila, homolog) 1 (DIAPH1) | 1 | AF051782 | B, monocyte stimulated | + | + | | + | + | | | |
| diaphorase (NADH) (cytochrome b-5 reductase) (DIA1) | | Y09501 | + | + | + | + | + | + | | | |
| differentiated Embryo Chondrocyte expressed gene 1 (DEC1) | | AB004066 | | + | | | + | + | | | |

| WO 00/40/49 | | | | | | | | | 21/CA00/00003 |
|---|---|----------------------|--------------------------|--------------|--|--------------|--------------|---|------------------------|
| differentiated Embryo Chondrocyte expressed gene 1 (DEC1) (low match) | 1 | AB004066 | | | | | | | |
| differentiation antigen | 1 | L23415 | | | | | | | |
| DiGeorge syndrome critical | 1 | X84076 | | + | + | | | + | |
| region gene 2 (DGCR2) dihydrolipoamide | 2 | J03620 | | + | | \vdash | + | + | |
| dehydrogenase (E3 component of pyruvate | | | | Ì | | | | | |
| dehydrogenase complex. | | | | | | | | | |
| 2-oxo-glutarate complex, branched chain keto acid | | | | | | | | | |
| dehydrogenase complex) | | | | | | } | | | |
| dihydrolipoamide S- acetyltransferase (E2 | 1 | Y00978 | В | + | | | + | | |
| component of pyruvate | | | | ŀ | | | | | |
| dehydrogenase complex) (DLAT) | | | | | | | | | |
| dihydropyrimidinase-like 2 (DPYSL2) | 1 | D78013 | | + | + | | + | + | |
| dinG gene | 1 | Y10571 | | | | | | | |
| diptheria toxin resistance protein required for | 3 | AF053003 | В | + | + | | + | + | |
| diphthamide biosynthesis | | | | | | | | | |
| (Saccharomyces)-like 2 (DPH2L2) | | l | | | | | | | |
| disintegrin-protease (non- exact 72%) | 1 | Y13323 | | | | | | | |
| DJ-1 protein | 2 | AF021819 | + | + | + | + | | + | |
| Dmx-like 1 (DMXL1) | 1 | AJ005821 | + | | + | + | | | |
| DNA (cytosine-5-)- methyltransferase 1 (DNMT1) | 3 | X63692 | T activated, lymphoma | + | | | + | + | |
| DNA fragmentation factor, 40 kD, beta subunit (DFFB) | 1 | AF064019 | | | | | | | |
| DNA fragmentation factor, 45 kD, alpha subunit | 2 | U91985 | 1 | + | + | | | + | |
| (DFFA) | | | | | | | | | |
| DNA mismatch repair protein (hMLH1) | 1 | U17840 | | | | | İ | | |
| DNA segment on | 3 | M64241 | + | + | + | + | + | + | high in many libraries |
| chromosome X (unique) 648 expressed sequence | | | | | | | | | |
| DNA segment, single copy probe LNS-CAI/LNS-CAII | 3 | M73547 | | + | + | + | | + | |
| (deleted in polyposis (D5S346) | | | | | | | | | |
| DNA-damage-inducible transcript 1 (DDIT1) (low | 1 | L24498 | | | | | | | |
| match) | | A 1004300 | | <u> </u> | | _ | | | |
| DnaJ protein | 1 | AJ001309 AJ001309 | | <u> </u> | | - | | | |
| docking protein 2, 56kD | 1 | AF034970 | | | | - | | _ | |
| (DOK2) dolichyl- | 1 | D89060 | + | + | + | + | + | + | activated T cell |
| diphosphooligosaccharide- protein glycosyltransferase (DDOST) | ' | 000000 | | | | , | | | don't diod ' oon |
| dolichyl-phosphate | 1 | D86198 | Tactivated | + | + | | + | - | |
| mannosyltransferase polypeptide 1, catalytic subunit (DPM1) | | |] | | | | | | |
| down-regulated by | 1 | AJ223183 | | | | | + | - | |
| activation (immunoglobulin superfamily) (DORA) | | | | | | | ĺ | | |
| down-regulated in | 1 | P40879 | | | | | | | |
| adenoma DRA (low match) D-type cyclin-interacting | 1 | AF082569 | В | | | \vdash | + | + | |
| protein 1 (DIP1) | | <u>L</u> | <u> </u> | <u> </u> | <u> </u> | | <u> </u> | | |

| WO 00/40749 | | | | | | | | | .17CA00/00005 |
|--|-----|----------|-----|---|----------|---|----------|----------|---------------|
| dual specificity phosphatase 1 (DUSP1) | 4 | X68277 | + | + | + | + | + | + | |
| dual specificity phosphatase 11 (RNA/RNP) complex 1-interacting) | 1 | AF023917 | + | + | + | + | | + | |
| (dusp11) dual specificity phosphatase 3 (vaccinia virus phosphatase VH1- related) (DUSP3) | 1 - | L05147 | | + | + | | + | + | |
| dual specificity phosphatase 6 (DUSP6) | 6 | X93920 | + | + | + | + | + | + | |
| dynactin 1 (p150, Glued (Orosophila) homolog) (DYTN1) | 3 | X98801 | | | | | | | |
| dynactin 1 (p150, Glued (Drosophila) homolog) (DYTN1) (low match) | 1 | X98801 | В | + | + | | | | |
| dynamin 2 (DNM2) | 1 | L36983 | | | | | | | |
| dynamitin (dynactin complex 50 kD subunit) (DCTN-50) (non-exact 88%) | 1 | U50733 | | | | | | | |
| dynein, axonemal, heavy polypeptide 17-like (non- exact, 57%aa) | 1 | X99947 | | | | | | | |
| dynein, cytopiasmic, light intermediate polypeptide 2 (DNCLI2) | 1 | AF035812 | В | + | + | | | + | |
| dynein, cytoplasmic, light intermediate polypeptide 2 (DNCLI2) (non-exact, 69%) | 1 | AF035812 | | | | | | | |
| dyskeratosis congenita 1, dyskerin (DKC1) | 1 | U59151 | В | + | | | + | + | |
| dystonia 1, torsion (autosomal dominant) (DYT1) | 1 | AF007871 | | + | + | + | | + | |
| dystrobrevin, beta (DTNB) | 1 | AF022728 | | + | | 1 | | | |
| dystrophia myotonica- containing WD repeat motif (DMWD) | 1 | L19267 | | + | + | | + | + | · |
| dystrophia myotonica- protein kinase (DMPK) | 1 | L08835 | + | + | + | | | + | |
| dystrophin (muscular dystrophy, Duchenne and Becker types) (DMD) (low match, 59%aa) | 1 | X14298 | | | | | | | |
| E1B-55kDa-associated | 1 | AJ007509 | - W | + | + | | + | + | |
| E2F transcription factor 3 (E2F3) | 2 | D38550 | | + | + | + | + | + | |
| E2F transcription factor 4, p107/p130-binding (E2F4) | 1 | X86096 | В | + | | | + | <u> </u> | |
| E2F transcription factor 5, p130-binding (E2F5) | 2 | U15642 | + | + | <u> </u> | + | <u>_</u> | + | |
| E74-like factor 1 (ets domain transcription factor) (ELF1) | ļ | M82882 | В | | + | | + | + | |
| È74-like factor 4 (ets domain transcription factor) (ELF4) | | U32645 | | + | + | | | + | |
| E74-like factor 4 (ets domain transcription factor) (ELF4) (non-exact, 71%) | | U32645 | | | | | _ | | |
| early development regulator 2 (homolog of polyhomeotic 2) (EDR2) | 4 | U89278 | + | + | + | + | | + | , |
| EBV induced G-protein coupled receptor (EBI2) | 7 | L08177 | W | | | | | | |
| ecotropic viral integration site 2B (EVI2B) | 3 | M60830 | | + | | + | | | |

| | | | | | | | | | | | |
|--|----|------------|---|---|---|---|---|---|------|-----|---|
| ectin, galactoside-binding, soluble, 1 (galectin 1) (LGALS1) | 1 | J04456 | | | | | | + | | | |
| EGF-like-domain, multiple 4 (EGFL4) | 1 | AB011541 | | | | | | | | |] |
| eIF-2-associated p67 | 3 | U13261 | В | + | | | | + | | | 1 |
| elastin (supravalvular aortic stenosis, Williams-Beuren syndrome) (ELN) (low match) | 1 | M24782 | | + | + | | | | | | |
| elav-type RNA-binding protein (ETR-3) | 3 | U69546 | | | | | | _ | | | |
| electron-transfer- flavoprotein, alpha polypeptide (glutaric aciduria II) (ETFA) | 2 | J04058 | | + | | | | | | | |
| ELK3, ETS-domain protein (SRF accessory protein 2) (ELK3) | 2 | Z36715 | | | + | | | + | | | |
| elongation factor 1-beta | 1 | L26404 | | | | | | | | | |
| elongation factor Ts (mitochondrial protein) | 1 | AF110399 | | | | | | | | | |
| elongation factor Tu- nuclear encoded mitochondrial | 1 | X84694 | | ä | | | | | | |] |
| eMDC II protein | 1 | AJ242015.1 | | 1 | | | | | | | 1 |
| ems1 sequence (mammary tumor and squamous cell carcinoma-associated (p80/85 src substrate) | 1 | M98343 | | + | + | | + | + | | *** | |
| (EMS1) endogenous retroviral element HC2 | 1 | Z70664 | | + | | | | | | | 1 |
| endosulfine alpha (ENSA) | 1 | X99906 | T | + | | | | | | | 1 |
| endothelial differentiation, sphingolipid G-protein- coupled receptor, 1 (EDG1) | 2 | M31210 | | + | + | + | | + | | | |
| endothelial differentiation, sphingolipid G-protein- coupled receptor, 1 (EDG1) (low match 66%) | 1 | M31210 | | | | | | | | | |
| endothelial monocyte- activating polypeptide (EMAPII) | 1 | U10117 | + | + | + | + | | + | | | |
| enolase 1, (alpha) (ENO1) | 12 | M14328 | + | + | + | + | + | + | | |] |
| enolase 2, (gamma, neuronal) (ENO2) | 1 | X51956 | | + | | | | | | | |
| enolase-alpha | 1 | D28437 | | | | | | | | | |
| enoyl Coenzyme A hydratase 1, peroxisomal (ECH1) | 2 | U16660 | | | | | | | | | |
| enoyl Coenzyme A hydratase, short chain, 1, mitochondrial (ECHS1) | 1 | D13900 | + | + | + | + | + | + | | | |
| ENOYL-COA HYDRATASE, MITOCHONDRIAL PRECURSOR (SHORT CHAIN ENOYL-COA HYDRATASE) (SCEH) (ENOYL-COA HYDRATASE 1) (low match, non-exact 56%) | 1 | P30084 | | | | | | | | | - |
| epidermal growth factor receptor pathway substrate 15 (EPS15) | 2 | U07707 | | + | | + | | + | | | |

| WO 00/40/49 | | | | | | | | | |
|---|-----|----------|------|---|---|---|---|---|------------------------------|
| EPIDIDYMAL SECRETORY PROTEIN E1 PRECURSOR (EPI-1) (HE1) (EPIDIDYMAL SECRETORY PROTEIN 14.6) (ESP14.6) | 2 | Q15668 | | | | | | | |
| epithelial membrane protein 3 (EM[P3) | 1 | U87947 | + | + | + | + | | + | |
| Epoxide hydrolase 1, microsomal (xenobiotic) (EPHX1) | 1 | L29766 | | | | | | | + only |
| ERCC2 (=L47234) | 1 | X52221 | - | | | | | | |
| ERF-2 | 3 | U07802 | + | + | + | + | | + | high in gall bladder |
| ERp28 protein | 1 | X94910 | + | + | + | + | | + | |
| erythrocyte membrane protein | 2 | M81635 | | | | | | | |
| erythroleukemic cells K562 | 2 | L25343 | | | | | | | |
| EST (Hs.189509) | 2 | U24166 | | | | | | | |
| estrogen receptor-related protein (hERRa1) | 1 | L38487 | | | | | | | |
| ESTs, Highly similar to ADENYLOSUCCINATE SYNTHETASE | 1 | X66503 | В, Т | + | + | | | | |
| ESTs, Moderately similar to cysteine-rich fibroblast growth factor receptor | 1 | U28811 | + | + | + | + | | + | |
| ET binding factor 1 (SBF1) | 1 | U93181 | + | + | | | | + | |
| ets domain protein ERF | 1 | U15655 | + | + | + | + | | + | |
| eukaryotic translation elongation factor 1 alpha 1 (EEF1A1) | 326 | X03558 | Ī | + | + | | | + | |
| eukaryotic translation elongation factor 1 alpha 1 (EEF1A1) (low match) | 1 | X03558 | | | | | | | |
| eukaryotic translation elongation factor 1 alpha 1 (EEF1A1) (low match) | 1 | X03558 | | | | | _ | | |
| eukaryotic translation elongation factor 1 beta 2 (EEF1B2) | 5 | X60489 | + | + | + | + | | + | |
| eukaryotic translation elongation factor 1 delta (guanine nucleotide exchange protein) (EEF1D) | 1 | Z21507 | + | + | + | + | + | + | |
| eukaryotic translation elongation factor 1 gamma (EEF1G) | 31 | Z11531 | | | | | | | |
| eukaryotic translation elongation factor 2 (EEF2) | 2 | X51466 | | + | | | | + | |
| eukaryotic translation initiation factor 2, subunit 1 (alpha, 35kD) (EIF2S1) | | J02645 | | | | | | | |
| eukaryotic translation initiation factor 2, subunit 2 (beta, 38kD) (EIF2S2) | 1 | M29536 | | | | | | | |
| eukaryotic translation initiation factor 2, subunit 3 (gamma, 52kD) (EIF2S3) | 3 | L19161 | | + | + | | | | |
| eukaryotic translation initiation factor 3, subunit 10 (theta, 150/170kD) (EIF3S10) | 2 | U78311 | | | | | | | |
| eukaryotic translation initiation factor 3, subunit 2 (beta, 36kD) (EIF3S2) | 3 | U36764 | + | + | + | + | + | + | high in white blood cells |
| eukaryotic translation initiation factor 3, subunit 3 (gamma, 40kD) (EIF3S3) | 6 | U54559 | + | + | + | + | | + | high in spleen |
| eukaryotic translation initiation factor 3, subunit 4 (delta, 44kD) (EIF3S4) | 9 | AF020833 | | + | + | + | | + | |

| WU 00/40/49 | | | | | | | | _ | C1/CA00/00003 |
|--|----|----------|------|---|---|---|---|---|---|
| eukaryotic translation initiation factor 3, subunit 6 (48kD) (EIF3S6) | 4 | U94175 | + | + | + | + | | + | high in bladder |
| eukaryotic translation initiation factor 3, subunit 6 (EIF3S6) | 1 | U62962 | | + | + | + | | + | Highly represented (1.4833 pct) in library 36 human gall bladder |
| eukaryotic translation initiation factor 3, subunit 7 (zeta, 66/67kD) (EIF3S7) | 3 | U54558 | + | + | + | + | | + | |
| eukaryotic translation initiation factor 3, subunit 8, 110KD (EIF3S8) | 5 | U46025 | + | + | + | + | + | + | high in testis |
| eukaryotic translation initiation factor 4 gamma, 1 (EIF4G) | ſ | AF012088 | | | | | | | |
| eukaryotic translation initiation factor 4 gamma, 1 (EIF4G) (low match) | 1 | AF012088 | | | | | | | |
| eukaryotic translation initiation factor 4 gamma, 1 (EIF4G1) | 2 | D12686 | | | | | | | |
| eukaryotic translation initiation factor 4 gamma, 2 (EIF4G2) | 6 | U73824 | + | + | + | + | + | + | |
| eukaryotic translation initiation factor 4 gamma, 2 (EIFG2) | 2 | U76111 | + | + | + | + | + | + | |
| eukaryótic translation initiation factor 4A, isoform 1 (EIF4A1) | 29 | D13748 | | | | | | | |
| eukaryotic translation initiation factor 4A, isoform 2 (EIF4A2) | 11 | D30655 | + | + | + | + | + | + | |
| eukaryotic translation initiation factor 4B (EIF4B) | 18 | X55733 | + | + | + | + | | + | |
| eukaryotic translation initiation factor 4E (EIF4E) | 1 | P06730 | | | | | | | |
| Eukaryotic translation initiation factor 4E binding protein 2 (EIF4EBP2) | 3 | L36056 | Т, В | | | | + | + | , |
| eukaryotic translation initiation factor 4H (EIF4H) | 2 | Q15056 | | | | | | | |
| eukaryotic translation initiation factor 5 (EIF5) | 2 | U49436 | + | + | + | + | + | + | |
| eukaryotic translation termination factor 1 (ETF1) | 2 | U90176 | + | + | + | + | | + | |
| EV12 protein | 1 | M55266 | | + | | ł | | 1 | |
| Ewing sarcoma breakpoint region 1 (EWSR1) | 1 | X66899 | + | + | + | + | | + | |
| EWS/FLI1 activated transcript 2 homolog (EAT- 2) | 2 | AF020264 | | | | | | | |
| EWS-E1A-F chimeric protein | 1 | U35622 | | | | | | | |
| excision repair cross- complementing rodent repair deficiency, complementation group 1 (includes overlapping antisense sequence) (ERCC1) | 1 | M28650 | + | + | + | + | | + | |
| excision repair cross- complementing rodent repair deficiency, complementation group 5 (xeroderma pigmentosum, complementation group G (Cockayne syndrome)) (ERCC5) | 1 | X69978 | | + | + | + | | + | |
| exostoses (multiple)-like 3 (EXTL3) | 1 | AF001690 | | + | + | + | | + | |
| F11 | 1 | X77744 | | | | + | | | |
| | | | | | | | | | |

| WO 00/40743 | | | | | | | | | |
|---|--------------|----------|----|---|--------------|----------|--------------------|----------|------------------------|
| F1-ATPase beta subunit | 2 | X03559 | | | | | | | |
| (F-1 beta) Fanconi anaemia group A | 2 | Z83095 | | | - | \vdash | | | |
| Fanconi anemia, | - | X99226 | + | + | + | + | | | |
| complementation group A (FANCA) | | 11222 | | | | | | + | |
| far upstream element (FUSE) binding protein 1 (FUBP1) | 2 | U05040 | + | | + | | | + | |
| farnesyl diphosphate synthase (farnesyl | 1 | J05262 | + | + | + | + | | + | |
| pyrophosphate synthetase, dimethylallyltra nstransferase, geranyltranstransferase) (FDPS) | | | | | | | | | |
| farnesyl-diphosphate farnesyltransferase 1 (FDFT1) | 2 | X69141 | + | + | + | + | + | + | |
| farnesyltransferase, CAAX box, beta (FNTB) | 2 | L00635 | | + | + | | | | |
| Fas ligand (gene and promoter region) | 1 | AF044583 | | | | | | | |
| Fas-ligand associated factor 1 | 1 | U70667 | | | | | | | |
| fatty-acid-Coenzyme A ligase, long-chain 1 (FACL1) | 4 | D10040 | + | + | + | + | + | + | |
| Fc fragment of IgA, receptor for (FCAR) | 1 | X54150 | | | | | | | |
| Fc fragment of IgE, high affinity I, receptor for; gamma polypeptide (FCER1G) | 1 | M33195 | + | + | + | + | | + | |
| Fc fragment of IgE, low affinity II, receptor for (CD23A) (FCER2) | 2 | X04772 | + | + | | | | | |
| Fc fragment of IgG, low affinity IIa, receptor for (CD32) | 6 | M31932 | + | + | + | + | + | + | |
| Fc fragment of IgG, low affinity IIa, receptor for (CD32) (FCGR2A) | 1 | X62572 | + | + | + | + | + | + | |
| Fc fragment of IgG, low affinity Illa, receptor for (CD16) (FCGR3A) | 34 | X07934 | + | + | + | + | | + | |
| Fc fragment of IgG, receptor, transporter, alpha (FCGRT) | 3 | U12255 | | + | + | + | + | + | high in many libraries |
| fc-fgr | 1 | Z13983 | | | | | | | |
| Fc-gamma-receptorIIIB (FCGR3B) | 2 | M90746 | | | | | | | |
| feline sarcoma (Snyder- Theilen) viral (v- fes)/Fujinami avian sarcoma (PRCII) viral (v- fps) oncogene homolog(FES) c-fes/fps) | 3 | X06292 | | | | | | | |
| female sterile homeotic- related gene 1 (mouse homolog) (FSRG1) | 2 | X96670 | + | + | + | + | | + | |
| ferritin L-chain | 9 | Y09188 | | | | | $oxedsymbol{oxed}$ | | |
| ferritin, heavy polypeptide 1 (FTH1) | 4 | M11146 | + | + | + | + | + | + | |
| fertilin alpha pseudogene | 1 | Y09232 | | 1 | | ↓_ | 4- | - | |
| fetal Alzheimer antigen (FALZ) | 2 | U05237 | | + | | | | <u></u> | |
| fetal Ig heavy chain variable region | 1 | M34024 | | | | | | | |
| fibrillarin (FBL) | 1 | X56597 | + | + | + | + | + | + | |
| fibrinogen-like protein 2 (T49) | 3 | Z36531 | | | | + | | <u> </u> | |
| | | | 42 | | | | | | |

42

| WO 00/40/49 | | | | | | | | • | |
|---|----|----------|---|---|---|---|----------|---|---|
| fibroblast growth factor receptor 2 (bacteria- | 1 | M35718 | + | + | + | + | + | + | |
| expressed kinase, keratinocyte growth factor receptor, craniofacial | | | | | | | | | |
| dysostosis 1, Crouzon syndrome, | | | | | | | | | |
| Pfeiffer syndrome, Jackson-Weiss) (FGFR2) | | , | | | | | | | |
| ficolin (collagen/fibrinogen domain-containing) 1 (FCN1) | 19 | D83920 | | - | | + | | + | |
| filamin A, alpha (actin- binding protein-280) (FLNA) | 2 | X53416 | | | | | | | |
| filamin B, beta (actin- binding protein-278) (FLNB) | 1 | AF043045 | | + | + | | + | | |
| Finkel-Biskis-Reilly munne sarcoma virus (FBR-MuSV) ubiquitously expressed (fox derived); ribosomal protein S30 (FAU) | 2 | X65923 | + | + | + | + | + | + | Highly represented in intraepithelial neoplasia and invasive prostate tumor |
| FK-506 binding protein | | M80199 | + | + | + | + | | + | |
| FK506-binding protein 1A (12kD) (FKBP1A) | 2 | M34539 | | | | | | | |
| FK506-binding protein 18 (12.6 kD) (FKBP1B) | 1 | M92423 | | + | | + | | + | |
| FK506-binding protein 5 (FKBP5) | 4 | U71321 | | + | + | + | | + | |
| Flightless I (Drosophila) homolog (FLII) | 3 | U80184 | | + | | | | | |
| Flightless I (Drosophila) homolog (FLII) (low match) | 1 | U80184 | | | | | | | |
| FLN29 (FLN29) | 2 | AB007447 | | + | | + | <u> </u> | + | |
| flotillin 2 (FLOT2) | 5 | M60922 | + | + | + | + | + | + | |
| folate receptor 2 (fetal) (FOLR2) | 1 | AF000380 | | + | + | + | | + | |
| forkhead (Drosophila) homolog (rhabdomyosarcoma) like 1 (FKHRL1) | 1 | AF032886 | + | + | | + | | + | |
| Formyl peptide receptor 1 (FPR1) | 9 | M60627 | + | + | + | + | | + | |
| formyl peptide receptor-like 1 (FPRL1) | 1 | M84562 | | | | | | | Found only in libraries from placenta |
| formyl peptide receptor-like 1 (FPRL1) (low score) | 1 | M84562 | | | | | | | |
| fragile X mental retardation 1 (FMR1) | 1 | L29074 | + | + | | + | | + | |
| fragile X mental retardation, autosomal homolog 1 (FXR1) | 1 | U25165 | + | + | + | + | | | |
| Friend leukemia virus integration 1 (FLI1) | 3 | M93255 | + | + | | | | | |
| fructose-bisphosphatase 1 (FBP1) | 1 | D26054 | | | | + | | + | |
| FSHD-associated repeat DNA, proximal region | 1 | U85056 | | | | | | | |
| fucose-1-phosphate guanylyltransferase (FPGT) | 1 | AF017445 | | + | + | + | | | · |
| full length insert cDNA clone ZA78A09 | 1 | AF086122 | | | | | | | |
| full length insert cDNA YP07G10 | 1 | AF075061 | | | | | | | |
| fumarate hydratase (FH) | 1 | U59309 | | + | + | + | | + | |
| FUS (low match) | 1 | X99006 | | | | | | | |
| FYN-binding protein (FYB-120/130) (FYB) | 16 | U93049 | | + | | + | | | |
| | | | | | | | | | - |

| WO 00/40/49 | | | | | | | | | |
|---|---|----------|-------------|----|--------------|----------|---|---|---------------------|
| G alpha interacting protein (GAIP) (low score) | 1 | X91809 | | | | | | | |
| G protein beta subunit-like protein 12.3 | 2 | D28398 | | | | | | | |
| G protein-coupled receptor 64 (HE6) (non-exact 59%) | 1 | X81892 | | | | + | | | |
| G protein-coupled receptor kinase 6 (GPRK6) | 2 | L16862 | + | + | + | | | + | |
| G1 to S phase transition 1 | 2 | X17644 | | + | + | + | + | + | , |
| (GSPT1) GA-binding protein transcription factor, beta | 1 | D13316 | | + | + | + | + | + | |
| subunit 2 (47kD) (GABPB2) galactose-1-phosphate | 2 | M60091 | | - | | | | | |
| uridylyltransferase (GALT) galactosidase, beta 1 | 3 | M27508 | | +- | | | + | + | |
| (GLB1) | 7 | M13701 | | | | | | | |
| (=X13223 N- acetylglucosamide-(beta 1- 4)-galactosyltransferase) | | | | | | | | | |
| galectin-9 isoform | 1 | AB006782 | + | 1 | | + | | + | |
| gamma2-adaptin (G2AD) | 1 | AF068706 | + | + | | + | | + | |
| gamma-actin | 2 | M37130 | | 1 | | <u> </u> | | + | |
| gamma-aminobutyric acid (GABA) B receptor 1 (GABBR1) | 2 | AJ012187 | _ | + | + | | | | |
| GATA-binding protein 2 (GATA2) | 1 | M68891 | | | | + | | + | |
| GATA-binding protein 3 (GATA3) | 1 | M69106 | | | + | + | | + | |
| GCN5 (general control of amino-acid synthesis, yeast, homolog)-like 1 (GCN5L1) | 3 | D64007 | + | + | + | + | | + | |
| GDP dissociation inhibitor 1 (GDI1) | 1 | D45021 | + | + | + | + | | + | high in adult brain |
| GDP dissociation inhibitor 2 (GCl2) | 4 | Y13286 | | | | | | | |
| GDS-related protein (HKE1.5) | 4 | U68142 | + | + | + | + | | + | |
| gelsolin (amyloidosis, Finnish type) (GSN) | 3 | X04412 | | + | + | + | + | + | |
| general transcription factor | 4 | Y14946 | + | + | + | + | + | + | |
| general transcription factor II, i, pseudogene 1 | 1 | AF038968 | + | + | + | + | + | + | high in fetal brain |
| (GTF2IP1) general transcription factor IIF, polypeptide 1 (74kD | 4 | X64037 | + | + | + | + | | + | |
| subunit) (GTF2F1) general transcription factor IIH, polypeptide 3 (34kD | 2 | Z30093 | В, Т | | | | | | |
| general transcription factor (IIH, polypeptide 4 (52kD | 3 | Y07595 | | + | | + | | + | |
| subunit) (GTF2H4) general transcription factor | 1 | U14134 | + | + | | + | - | + | |
| IIIA (GTF3A) general transcription factor | 1 | U02619 | - | + | | + | - | _ | |
| illC, polypeptide 1 (alpha subunit, 220kD) (GTF3C1) | | D13636 | + | + | + | + | + | + | |
| general transcription factor IIIC, polypeptide 2 (beta subunit, 110kD) (GTF3C2) | 3 | | T | | | | Ĺ | | |
| germline immunoglobulin heavy chain (IGHV@) | 1 | L06612 | | | | | | | |
| germline immunoglobulin heavy chain, variabl region | 1 | X92236 | | | | | | | |
| germline immunoglobulin heavy chain, variable | 1 | X92343 | | | | | | | |
| region, (21-2) | L | J | | | | — | | | |

| WO 00/40749 | | | | | | | | | . I/CA00/00003 |
|--|-------------|----------|--------------|--|--|---|----------|---|----------------|
| GLE1 (yeast homolog)-like, RNA export mediator (GLE1L) | 1 | AF058922 | | + | + | | | | |
| glia maturation factor, beta (GMFB) | 1 | AB001106 | + | + | | + | | + | |
| glioma-associated oncogene homolog (zinc finger protein) (GLI) | 1 | X07384 | | | | | | | |
| glioma-associated oncogene homolog (zinc finger protein) (GLI) (low | 1 - | X07384 | | | | | | ļ | |
| score) globin, alpha 2 | | V00516 | | 1 | | | | | |
| glucocorticoid receptor | 1 | M32284 | | 1 1 | | | \dashv | | |
| (i=M69104) l | | | ! | | + | + | | + | |
| glucocorticoid receptor (GRL) | 2 | U80947 | <u> </u> | + | | | | _ | |
| glucos phosphate isomerase (CONTAINS LARGE REPEAT) | 1 | L09105 | | | | | | | |
| glucosamine (N-acetyl)-6- sulfatase (Sanfilippo disease IIID) (GNS) | 1 | 212173 | + | | | | | | |
| glucosamine (N-acetyl)-6- sulfatase (Sanfilippo disease IIID) (GNS) (non- exact 56%) | 1 | Z12173 | | | | | | | |
| glucose transporter-like protein-III (GLUT3) | 1 | M20681 | | + | + | + | + | + | |
| glucose transporter-like protein-III (GLUT3) (low match) | 1 | M20681 | | | | | | | |
| glucosidase, alpha; acid (Pompe disease, glycogen storage disease type II) (GAA) | 1 | Y00839 | + | + | | + | | + | |
| glucosidase, beta; acid (includes glucosylceramidase) (GBA) | 1 | K02920 | + | + | + | + | | + | |
| glutamate dehydrogenase | 1 | M20867 | | + | + | + | + | + | |
| glutamate-ammonia ligase (glutamine synthase) (GLUL) | 12 | X59834 | + | + | + | + | | + | |
| glutamate-ammonia ligase (glutamine synthase) (GLUL) (low score) | 1 | Y00387 | | | | | | | |
| glutamate-cysteine ligase (gamma-glutamylcysteine synthetase), catalytic (72.8kD) (GLCLC) | 1 | M90656 | | | | + | | | |
| glutamine cyclotransferase | 1 | X71125 | | + | + | | | | |
| glutamine-fructose-6- phosphate transaminase 1 (GFPT1) | 1 | M90516 | | + | | + | | | |
| glutaminyl-tRNA synthetase | 1 | X72396 | | | | | | | |
| glutaminyl-tRNA synthetase (QARS) | 6 | X76013 | + | + | + | + | | + | |
| glutamyl-prolyl-tRNA synthetase (EPRS) | 1 | X54326 | | | | | | | |
| glutathione peroxidase 1 (GPX1) | 2 | M21304 | + | + | + | + | + | + | |
| glutathione peroxidase 4 (phospholipid hydroperoxidase) (GPX4) | 1 | X71973 | + | + | + | + | | + | |
| glutathione S-transferase pi | 1 | U30897 | | + | + | + | + | + | |
| (GSTP1) glutathione S-transferase | 1 | AF070657 | | 1 | | | | | |
| subunit 13 homolog glyceraldehyde-3- phosphate dehydrogenase (GAPD) | 12 | J02642 | | <u> </u> | | | + | | |
| (GAPU) | <u> </u> | 1 | 45 | | ــــــــــــــــــــــــــــــــــــــ | | | | |

| glycogenin (GYG) | 1 | U31525 | | + | + | + | | +1 | - |
|--|----------------|----------|-------|--------------|---|--------------|---|----|------------------|
| alycophorin C (Gerbich | 1 | X12496 | | + | + | + | | + | |
| blood group) (GYPC) glycoprotein M68 (GPM68) | 1 | U45955 | | + | + | | _ | | |
| glycyl-tRNA synthetase | 1 | U09587 | | + | + | + | | + | |
| (GARS) glyoxalase I (lactoyl | - 1 | L07837 | + | + | + | + | _ | + | |
| glutathione lyase) (GLYI) golgi autoantigen, golgin | | U51587 | | + | | + | _ | | |
| subfamily a, 1 (GOLGA1) | _ | | | | | | | | |
| golgi autoantigen, golgin subfamily a, 2 (GOLGA2) (non-exact, 70%) | 1 | L06147 | | | | | | | |
| golgi autoantigen, golgin subfamily a, 4 (GOLGA4) | 1 | U31906 | | | | | | | |
| golgi autoantigen, golgin subfamily b, macrogolgin (with transmembrane signal), 1 (GOLGB1) | 1 | X75304 | | + | + | + | | + | |
| gp25L2 protein | 4 | X90872 | | † | | | | | |
| grancalcin | 8 | M81637 | * | + | + | + | | | |
| granulin (GRN) | 16 | X62320 | + | + | + | + | | + | |
| granulin (GRN) (low match) | 1 | X62320 | | | | | | | |
| Granulysin (NKG5) | 5 | M85276 | + | | | | | + | |
| granzyme A (granzyme 1, cytotoxic T-lymphocyte-associated serine esterase 3) (GZMA) | 1 | M18737 | + | + | + | + | | + | |
| GRB2-related adaptor protein (GRAP) | 1 | U52518 | Tonly | | | | | | |
| Grb2-related adaptor protein 2 (GRAP2) | 7 | AF090456 | 1 | | | | + | | |
| GRO1 oncogene (melanoma growth stimulating activity, alpha) (GRO1) | 1 | X54489 | | | | + | | + | |
| growth arrest and DNA- damage-inducible gene (GADD153) | 1 | S40706 | | | | | | | |
| growth arrest-specific 7 (GAS7) | 4 | AB007854 | 1 | + | + | | | | |
| growth factor receptor- bound protein 2 (GRB2) | 1 | X62852 | В | + | | | + | + | |
| GS1 (protein of unknown function) | 1 | M86934 | | + | + | + | | | |
| GS3955 | 4 | D87119 | | + | + | + | | + | |
| GTP binding protein 1 | 1 | U87964 | | + | + | + | | | |
| (GTPBP1) GTP binding protein similar to S. cerevisiae HBS1 | 1 | U87791 | | + | + | + | | + | |
| (HBS1) GTPase activating protein- | 1 | AB011110 | | + | + | + | - | + | high fetal brain |
| like (GAPL) GTP-binding protein (low | 1 | Z49068 | | <u> </u> | ļ | - | - | | |
| match) | | P08754 | | | | | - | | |
| GTP-binding protein G(K), alpha subunit (=G(I) ALPHA-3)(=GTP-binding regulatory protein Gi alpha- 3 chain) | 1 | FU0/54 | | | | | | | |
| Gu protein (GURDB) | 2 | U41387 | + | 1 | + | + | | + | |
| guanine nucleotide binding protein | 1 | | | | | | | | |
| guanine nucleotide binding protein (G protein), alpha inhibiting activity polypeptide 2 (GNAI2) | 4 | J03004 | + | + | + | + | | + | |

| WO 00/40/49 | | | | | | | | | |
|---|----|----------|------|---|---|---|----|---|--------------------------------|
| guanine nucleotide binding protein (G protein), alpha inhibiting activity | 7 | M20597 | + | + | + | + | | + | |
| polypeptide 3 (GNAI3) | | | | | | | | | |
| guanine nucleotide binding protein (G protein), alpha stimulating activity polypeptide 1 (GNAS1) | 2 | X04409 | В, Т | + | | | + | + | |
| guanine nucleotide binding protein (G protein), alpha transducing activity polypeptide 2 (GNAT2) | 1 | Z18859 | | | | | | | |
| guanine nucleotide binding protein (G protein), beta 5 (GNB5) | 2 | AF017656 | | + | + | + | | + | |
| guanine nucleotide binding protein (G protein), beta polypeptide 1 (GNB1) | 5 | M36430 | + | + | + | + | + | + | |
| guanine nucleotide binding protein (G protein), q polypeptide (GNAQ) | 2 | AF011496 | | + | + | + | | | |
| guanine nucleotide binding protein-like 1 (GNL1) | 1 | L25665 | + | + | + | + | | + | |
| guanine nucleotide exchange factor | 1 | L13857 | + | + | + | + | | | |
| guanine nucleotide regulatory factor (LFP40) | | X15610 | + | + | + | + | | + | |
| guanine nucleotide regulatory factor (LFP40) | 1 | U72206 | + | + | + | + | | + | |
| GUANINÉ NUCLEOTIDE- BINDING PROTEIN BETA SUBUNIT-LIKE PROTEIN 12.3 (P205) (RECEPTOR OF ACTIVATED PROTEIN KINASE C 1) (RACK1) | 1 | P25388 | | | | | | | |
| GUANINE- MONOPHOSPHATE SYNTHETASE (GMPS) | 1 | U10860 | | | + | | | | |
| guanosine monophosphate reductase (GMPR) (non- exact, 72%) | 1 | M24470 | | | | | | | |
| guanosine-diphosphatase like protein | 1 | AF016032 | | | | | | | |
| guanylate binding protein 1, interferon-inducible, 67kD (GBP1) | 2 | M55542 | | + | + | + | + | + | |
| guanylate binding protein 2, interferon-inducible (GBP2) | 6 | M55543 | + | + | + | + | | + | |
| H2A histone family, member C (H2AFC) | 1 | Z83742 | | | | | | | |
| H2A histone family, member Y (H2AY) | 2 | AF041483 | + | + | + | + | | + | |
| H2B histone family, member L (H2BFL) | 2 | Z80783 | + | + | + | + | + | + | high in adrenal gland tumor |
| h2-calponin | 1 | D86059 | | | | | | | |
| H-2K binding factor-2 | 1 | L08904 | | + | + | + | | + | |
| H3 histone family, member K (H3FK) | 1 | Z83735 | | | | | | | |
| H3 histone, family 3A (H3F3A) | 7 | M11353 | + | + | + | + | | + | high in ovary |
| H3 histone, family 3B (H3.3B) (H3F3B) | 15 | Z48950 | + | + | + | + | | + | high in endothelial cells |
| hbc647 | 1 | U68494 | | + | + | + | + | | |
| heat shock 27kD protein 1 (HSPB1) | 1 | U12404 | | + | + | | +: | + | |
| heat shock 40kD protein 1 (HSPF1) | 4 | D85429 | + | + | + | + | + | + | high in testis |
| heat shock 60kD protein 1 (chaperonin) (HSPD1) | 3 | M22382 | + | + | + | + | + | + | |
| heat shock 70kD protein 1 (HSPA1A) | 7 | M59828 | + | + | + | + | + | + | high in activated T cells |
| | | | | | | | | | |

| W O 00/40/49 | | | | | | | | | |
|---|-----|----------|--------------|---|---|----------|----|----------|------------------------|
| heat shock 70kD protein 5 (glucose-regulated protein, 178kD) (HSPA5) | 13 | X87949 | | + | + | | + | | |
| heat shock 70kD protein 6 (HSP70B') (HSPA6) | 4 | X51757 | + | + | + | | | | |
| heat shock 70kD protein 9B (mortalin-2) (HSPA9B) | 2 | L15189 | | + | + | + | + | + | |
| HEAT SHOCK COGNATE 71 KD PROTEIN | 1 | P11142 | | | | | | | |
| heat shock factor binding protein 1 (HSBP1) | 2 | AF068754 | | | | | - | | |
| heat shock protein 90 | 13 | M27024 | + | + | + | + | + | + | high in many libraries |
| heat shock protein, DNAJ- like 2 (HSJ2) | 1 | D13388 | | + | + | | + | + | |
| Hect (homologous to the E6-AP (UBE3A) carboxyl terminus) domain and RCC1 (CHC1)-like domain (RLD) 1 (HERC1) | 1 | U50078 | | + | + | + | | | |
| hect domain and RLD 2 (HERC2) | 1 | AB002391 | + | + | + | + | | + | |
| helicase-like protein (HLP) | 1 | X98378 | + | + | | + | | + | |
| helix-loop-helix protein HE47 (E2A) | 1 | M65214 | | | | | | + | |
| hematopoietic cell-specific Lvn substrate 1 (HCLS1) | 18 | X16663 | + | | + | + | | + | |
| heme oxygenase (decycling) 1 (HMOX1) | 1 | X06985 | | + | | + | + | + | |
| HEMOGLOBIN ALPHA | 1 | P19015 | | | | | | | |
| hemoglobin beta (beta globin) | 5 | AF117710 | | | | | | | |
| hemoglobin, alpha 1 (HBA1) | 301 | √00491 | | | + | | + | + | |
| hemoglobin, alpha 1 (HBA1) (low match) | 1 | V00491 | | | | | | | |
| hemoglobin, alpha 1 (low match) | 1 | V00493 | | | | | | _ | |
| hemoglobin, alpha 1 (non- exact, 76%) | 1 | J00153 | · | | | <u> </u> | | | |
| hemoglobin, alpha 1 (non- exact, 82%) | 1 | ∨00493 | | | | <u> </u> | | <u> </u> | high in many libraries |
| hemoglobin, beta (HBB) | 129 | V00497 | + | + | + | + | + | + | nigh in many libraries |
| hemoglobin, beta (HBB) (low match) | 1 | V00497 | | | | | | | |
| hemoglobin, beta (HBB) (low match) | 1 | L48220 | | | | L | L. | <u></u> | · |
| hemokine (C-X-C motif), receptor 4 (fusin) (CXCR4) | 1 | D10924 | + | + | + | + | | + | |
| hemopoietic cell kinase (HCK) | 5 | M16591 | | | | + | | _ | |
| hepatitis C-associated microtubular aggregate protein p44 | 2 | D28908 | | | | | | | |
| hepatoma-derived growth factor | 1 | D16431 | + | + | + | + | | + | |
| Hermansky-Pudlak syndrome (HPS) | 2 | U65676 | | | | | | | |
| HERV-E integrase (non- exact 76%aa) | 1 | AF026246 | | | | | | | |
| heterogeneous nuclear protein similar to rat helix destabilizing protein (FBRNP) | 2 | S63912 | | + | + | + | | + | |
| heterogeneous nuclear ribonucleoprotein (C1/C2) (HNRPC) | 4 | M16342 | | | | | | | |
| heterogeneous nuclear ribonucleoprotein A/B (HNRPAB) | 1 | M65028 | + | + | + | + | + | + | |

| WO 00/40/49 | | | | | | | | | Illiah in aksalas |
|---|----|------------|---------------------------------------|----------------|---|-------|---|---|--|
| heterogeneous nuclear ribonucleoprotein A1 (HNRPA1) | 20 | X12671 | + | + | + | + | + | + | High in alveolar rhabdomyosarcoma |
| heterogeneous nuclear ribonucleoprotein A2/B1 (HNRPA2B1) | 3 | M29064 | + | + | + | + | + | + | High in activated T cell, fetal brain |
| heterogeneous nuclear ribonucleoprotein D (hnRNP D) | 2 | D55673 | + | + | + | + | + | + | |
| heterogeneous nuclear ribonucleoprotein D-like (HNRPDL) | 5 | D89092 | + | + | + | + | + | + | |
| heterogeneous nuclear ribonucleoprotein F (HNRPF) | 1 | L28010 | + | + | + | + | | + | |
| heterogeneous nuclear ribonucleoprotein F (HNRPF) (83%) | 1 | L28010 | | | | | | | |
| heterogeneous nuclear ribonucleoprotein G (HNRPG) | 2 | Z23064 | | + | + | + | | + | |
| heterogeneous nuclear ribonucleoprotein H (HNRPH) (FTP-3) | 3 | P55795 | | | | _ | | | |
| heterogeneous nuclear ribonucleoprotein H (HNRPH) (low match) | 1 | P31943 | · · · · · · · · · · · · · · · · · · · | | | | | | |
| heterogeneous nuclear ribonucleoprotein H1 (H) (HNRPH1) | 2 | L22009 | * | + | + | + | | + | |
| heterogeneous nuclear ribonucleoprotein K (HNRPK) | 21 | S74678 | + | + | + | + | + | + | |
| heterogeneous nuclear ribonucleoprotein R (HNRPR) | 1 | AF000364 | | + | + | + | + | + | |
| heterogeneous nuclear ribonucleoprotein U (scaffold attachment factor A) (HNRPU) | 3 | X65488 | + | + | + | + | + | + | |
| hexokinase 1 (HK1) | 2 | X66957 | | + | + | + | | + | |
| hexokinase 2 (HK2) | 3 | Z46376 | + | + | + | + | | + | |
| hexokinase 3 (HK3) | 2 | U51333 | | - | 1 | t | | | |
| hexosaminidase A (alpha polypeptide) (HEXA | 1 | S62047 | | | | | | | |
| HGMP07I gene for olfactory receptor | 2 | U76377 | | | | | | | |
| High density lipoprotein binding protein (HDLBP) | 2 | M64098 | + | + | + | + | + | + | |
| high-mobility group (nonhistone chromosomal) | 5 | X12597 | + | + | + | + | + | + | |
| protein 1 (HMG1) high-mobility group (nonhistone chromosomal) protein 1 (HMG1) (non- exact 60%) | 1 | D63874 | | | | | | | |
| High-mobility group (nonhistone chromosomal) protein 17 (HMG17) | 2 | M12623 | + | + | + | + | | + | |
| high-mobility group (nonhistone chromosomal) protein 2 (HMG2) | 2 | M83665 | + | + | + | + | + | + | |
| high-mobility group (nonhistone chromosomal) protein isoforms I and Y | 2 | L17131 | + | + | + | | + | + | |
| high-risk humanpapilloma viruses E6 oncoproteins targeted protein E6TP1 beta (=A8007900 KIAA0440) | 1 | AF090990.1 | | | | | | | |
| histidine ammonia-lyase (HAL) | 1 | D16626 | | | + | , onl | / | | |

| histidyl-tRNA synthetase | 2 | Z11518 | + | + | + | + | + | + | |
|---|--------------|------------|-------------|---------------|--|--|--------------|--|--|
| histocompatibility antigen (HLA-Cw3), class I | 1 | U31372 | | | | | | | |
| histone deacetylase 1 | 4 | U50079 | + | + | + | + | | + | |
| (HDAC) | | | | | | | | | |
| histone deacetylase 1 (HDAC1) | 2 | D50405 | + | + | + | + | | + | |
| histone deacetylase 5 (NY- CO-9) | 1 | AF039691 | | + | + | | | | |
| HK2 gene for hexokinase II | 1 | Z46362 | | | | | | | |
| HL9 monocyte inhibitory receptor precursor | 2 | U91928 | | | | + | | | |
| HLA class Theavy chain (HLA-Cw*1701) | 1 | | | | | | | | |
| HLA class I locus C heavy chain | 1 | X58536 | | | | | | | |
| HLA class II SB 4-beta | 1 | X03022 | | | | | | | • |
| Chain HLA class III region | 1 | U89335 | + | + | + | + | | + | |
| containing NOTCH4 gene | | Z72423 | | + | | \vdash | \vdash | | |
| | 2 | AJ006020 | | | | | | ├ | |
| HLA-A | | 1 | | 1 | | <u> </u> | | | <u> </u> |
| HLA-A*7402 | 1 | AJ223060 | | | | | <u> </u> | | |
| HLA-A11 | 1 | U02934 | | L | | L | L. | <u></u> | |
| HLA-B | 2 | X75953 | | | | | | | |
| HLA-B | 1 | X83401 | | 1 | | Г | | | |
| HLA-B | | X78426 | | + | | \vdash | | | |
| | ' | Z37166 | + | + | + | + | + | + | |
| HLA-B associated transcript-1 (D6S81E) | <u>2</u> | M33509 | | + | + | + | Ļ | <u> </u> | |
| HLA-B associated transcript-2 (D6S51E) | 4 | D44501 | | <u> </u> | | <u> </u> | | | |
| HLA-B*1529 | · · | | | | ļ | ١ | + | + | high in many libraries |
| HLA-Bw72 antigen | 119 | L09736 | + | + | + | + | <u> </u> | ļ <u>,</u> | ingit in many ilbranes |
| HLA-C gene (HLA- Cw*0701 allele) | 1 | D83957 | | | | <u> </u> | | | |
| HLA-CW*0701 | 9 | Z46810 | | | | | | | |
| HLA-Cw*0801 | 1 | D64151 | | 1 | 1 | 1 | 1 | | |
| HLA-CW*1203 | 1 | D64146 | | + | | | † | | |
| HLA-DC dassii | 2 | X00370 | | ┼┈─ | - | - | ╁─ | | |
| histocompatibility antigens alpha-chain (=K01160) | | 700070 | | | | | | | |
| HLA-DR alpha-chain | 17 | M60333 | + | + | + | + | + | + | high in spleen |
| HLA-F (leukocyte antigen | 3 | X17093 | | 1 | + | + | | + | |
| HMG box containing | 3 | AF019214 | | 1 | | \vdash | | | |
| protein 1 hMLH1 (=U83845) | | AB017806.1 | | + | | + | \vdash | ├ | ļ |
| | i ' | Y14155 | | | | ┼ | | | |
| Hmob33 | 3 | | | <u> </u> | <u> </u> | | | - | |
| HMT1 (hnRNP methyltransferase, S. cerevisiae)-like 1 (HRMT1L1) | 2 | U80213 | + | + | + | + | | + | |
| hnRNP C1/C2 | 2 | D28382 | | | | Γ | | | |
| homeobox (=X58250 Mouse homeo box protein, | 1 | M60721 | | 1 | | \vdash | | | |
| put. transcription factor involved in embryogenesis and hematopoiesis) | | | | | | | | | |
| homeobox protein (HLX1) (=M60721) | 1 | U14326 | | | | | | | |
| homeodomain-interacting protein kinase 3 (HIPK3) | 1 | AF004849 | + | | + | + | | + | |
| homolog of Drosophila past (PAST) | 2 | AF001434 | + | + | + | + | | + | |
| homolog of yeast (S. | 3 | D50916 | | +- | + | + | | + | |
| cerevisiae) ufd2 (UFD2) | <u> </u> | <u> </u> | 0 | | Ь | | Ь— | ٠ | <u> </u> |

| WO 00/40749 | | | | | | | | PC | I/CA | JU/UU | 003 | |
|--|---|------------|--|--------------|---|--|----------|--|------|-------|-----|---|
| HPV16 E1 protein binding protein | 1 | U96131 | | + | + | \Box | | + | | | | |
| HRIHFB2157 | 1 | AB015344 | | + | + | | | + | | | | |
| HRX-like protein (=AF010403 ALR) | 1 | Y08836 | | | | | <u> </u> | | | | | |
| hsc70 gene for 71 kd heat shock cognate protein | 3 | Y00371 | | | | | | | | | | |
| HSPC012 | 1 | AF077036.1 | | \vdash | | | \vdash | | | | | |
| HSPC021 | 1 | AF077207.1 | <u> </u> | | - | | _ | H | | | | |
| HsPex13p | 1 | U71374 | | + | _ | | | | | | _ | _ |
| htra2-beta-2 | - | U87836 | + | + | + | + | | + | | | | |
| HU-K4 | 1 | U60644 | | | | ┼── | \vdash | | | | | |
| hunc18b2 | 1 | U63533 | | ╁ | + | + | \vdash | + | | | | _ |
| HUNKI | 1 | Y12059 | + | + | | + | + | + | | | | |
| huntingtin-interacting protein HYPA/FBP11 (HYPA) | 1 | AF049528 | | | | - | | | | | | |
| hVps41p (HVPS41) | 1 | U87309 | · · · · · · · · · · · · · · · · · · · | † | | † | | | | | | |
| hydroxyacyl-Coenzyme A dehydrogenase/3-ketoacyl- Coenzyme A thiolase/enoyl-Coenzyme A hydratase (trifunctional protein), alpha subunit (HADHA) | 1 | U04627 | | + | + | | + | | | | | |
| (HADHA) hydroxyacyl-Coenzyme A dehydrogenase/3-ketoacyl- Coenzyme A thiolase/enoyl-Coenzyme A hydratase (trifunctional protein), beta subunit (HADHB) | 1 | D16481 | + | + | + | + | | + | | • | | |
| hydroxysteroid (17-beta) dehydrogenase 1 (HSD17B1) | 1 | U34879 | | + | | | + | | | | | |
| hypothetical protein | 1 | | | | | | | | | | | |
| hypothetical protein (AL008729) (dJ257A7.2) | 1 | İ | : | | | | | | | | | |
| hypothetical protein (CIT987SK_2A8_1 chromosome 8) | 1 | U96629 | | | | | | | - | | | |
| hypothetical protein (clone 24640) | 1 | AF055004 | | | | | | | | | | |
| hypothetical protein (clone ICRFp507G2490). | 1 | Z70222 | | 1 | | | | | | | | |
| hypothetical protein (dJ1042K10.4) (non-exact 76%) | 1 | AL022238 | | | | | | | | | | |
| hypothetical protein (dJ465N24.1 similar to predicted yeast and worm proteins) | 2 | AL031432 | | | | | | | | | | |
| hypothetical protein (dJ487J7.1.1) | 2 | AL008730 | | | | | | | | | | |
| hypothetical protein (dJ753P9.2) | 2 | AL023653 | | | | | | | | | | |
| hypothetical protein (DKFZp586I111) | 1 | AL050131.1 | | | | | | | | | | |
| hypothetical protein (J257A7.2) | 1 | AL008729 | | | | | • | | | | | |
| hypothetical protein (KIAA0440) (=AF026504 R.norvegicus SPA-1 like protein) | 1 | AB007900 | | | | | | | | | | |
| hypothetical protein (L1H 3' region) | 1 | | | | | | | | | | | |
| hypothetical protein (S164) | 1 | P49756 | | | | | | | | | | |
| | | | | | | | _ | | | | | |

| VI O 00/40/42 | | | | | | | | | |
|--|-------------|----------|---------------------------------------|--|--|--|---------|--|---|
| hypothetical protein (similar to thrombospondin) (non- exact 56%) | 1 | AF109907 | <u> </u> | | | | | | |
| hypothetical protein 3 | | | | | | | | | |
| hypothetical protein B | | U47926 | · · · · · · · · · · · · · · · · · · · | | | | | | |
| (HSU47926) (non-exact, 56%) | | | | | | | | | |
| hypothetical protein from BCRA2 region (CG005) | 3 | U50532 | + | + | + | + | | + | |
| hypoxia-inducible factor 1, | 1 | AF050115 | | - | | <u> </u> | | | |
| alpha subunit (basic helix- loop-helix transcription factor) (HIF1A) | | | | | | | | | _,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,, |
| la-associated invariant gamma-chain (clones lambda-y (1,2,3)) | 1 | M13555 | | | | | | | |
| iduronate 2-sulfatase | 2 | M58342 | + | + | + | + | | + | |
| (Hunter syndrome) (IDS) Ig heavy chain V region | | L20779 | | - | | | | _ | |
| (=D11016) | 2 | M34024 | | | | ļ | | | |
| region | | | | | | | | | |
| Ig heavy chain variable region (VH4DJ) (clone T14.4) | 1 | 275378 | | | | | | | |
| Ig heavy chain variable region (VH4DJ) (clone T22.18) | 1 | Z75392 | | | | | | | |
| lg J chain | 1 | M12378 | | + | <u> </u> | | | | |
| lg kappa | 1 | 549007 | | | | _ | | | |
| IG kappa light chain | 1 | X63398 | | | | | | | |
| variable region A20 | | D90158 | | + | - | | | | |
| J-region (=X59315) Ig lambda light chain | | Z85052 | | ↓ | | <u> </u> | _ | | |
| variable region (26- 34ITIIIF120) | | | | <u> </u> | | | | | |
| Ig mu-chain VDJ4-region | 1 | M16949 | | | | | | | |
| Ig rearranged anti-myelin kappa-chain (V-J4-region, hybridoma AE6-5) | 1 | M29469 | | | | | | | |
| lg rearranged H-chain mRNA V-region | 2 | M97920 | | | | | | | |
| Ig rearranged light-chain V | 1 | M74020 | | | | | | | |
| region (=D90158) IGF-II mRNA-binding | | U97188 | + | + | + | ├ | - | - | |
| protein 3 (KOC1) (non- exact, 75%) | | | | | | | _ | | |
| IgG Fc binding protein (FC(GAMMA)BP) | 1 | D84239 | + | + | | + | | + | |
| lgG heavy chain variable region (vH26) | 1 | M83136 | | | | | | | |
| IgM heavy chain (C mu, | | X14939 | | | | | | | |
| membrane exons) IkB kinase-beta (IKK-beta) | 1 | AF029684 | | + | | - | | | |
| IL-1 receptor type II | | U14177 | | + | | - | | | |
| IL2-inducible T-cell kinase (ITK) | 2 | S65186 | | | - | | - | | |
| immediate early protein (ETR101) | 1 | M62831 | + | | + | + | | + | |
| immunogloblin light chain (lambda) | 1 | D87018 | | | | | | | |
| Immunoglobulin (CD79A) binding protein 1 (IGBP1) | 1 | Y08915 | В, Т | + | + | | + | | |
| immunoglobulin C (mu) and C (delta) heavy chain | 2 | X57331 | | | | | | | |
| (=K02878) immunoglobulin G Fc | 1 | Z46223 | | + | | - | | | |
| receptor IIIB immunoglobulin gamma 3 | 3 | Y14737 | + | | ļ <u> </u> | + | | + | high in many libraries |
| (Gm marker) (IGHG3) | | | | <u> </u> | L | <u> </u> | L | <u> </u> | |
| | | | `^ | | | | | | |

| W O 00/40/49 | | | | | | | | | | |
|--|---|----------|----------|----------|---|---|---|---|---|--|
| immunoglobulin gamma heavy chain variable region (=X61011) | 1 | Z66542 | | | | | | | | |
| immunoglobulin heavy chain (VI-3B) | 1 | X62109 | | † | | | | | | |
| immunoglobulin heavy | 1 | X86356 | | 1 | | | | | | |
| chain J region | 2 | X86355 | | | | | | | | |
| chain J region, B1 haplotype | | AF062126 | | <u> </u> | | | | | | |
| immunoglobulin heavy chain variable region (IGH) (clone 21u-48) | 1 | AF002120 | | | | | | | | |
| immunoglobulin heavy chain variable region (IGH) (clone 23u-1) | 1 | AF062212 | | | | | | | | |
| immunoglobulin heavy chain variable region V1-18 (IGHV@) (=X60503) | 2 | M99641 | | | | | | | | |
| immunoglobulin heavy chain variable region V3-43 | 2 | M99672 | | | | | | | | |
| (IGHV@) immunoglobulin heavy chain variable region V3-7 | 3 | M99649 | | | - | | | | | |
| (IGHV@) immunoglobulin IgH heavy chain Fd fragment | 1 | U07986 | | | | _ | | | | |
| immunoglobulin kappa light | 1 | X58081 | | | | | | | | |
| immunoglobulin kappa light chain V-segment A27 | 1 | X12686 | | | | | | | | |
| immunoglobulin light chain | 1 | D86990 | | | | | | | | |
| immunoglobulin light chain (low match) | 1 | D86996 | | | | | | | _ | |
| immunoglobulin light chain variable region (lambda IIIb subgroup) from IgM rheumatoid factor | 1 | L29157 | | | | | | | | |
| immunoglobulin M heavy chain V region=anti-lipid A antibody | 1 | \$50735 | | | | | | | | |
| immunoglobulin mu (IGHM) | 9 | X57086 | + | + | | + | | + | | |
| immunoglobulin mu binding protein 2 (IGHMBP2) | 1 | L24544 | T | + | | | + | | | |
| immunogiobulin superfamily, member 2 (IGSF2) | 1 | Z33642 | | | | | | | | |
| immunoglobulin VH mRNA (487 bp) (=M99652 immunoglobulin heavy chain variable region V3-11 ((GHV@)) | 1 | X61013 | | | | 1 | | | | |
| imogen 38 (IMOGEN38) | 1 | Z68747 | | + | + | + | | + | | |
| IMP (inosine monophosphate) dehydrogenase 1 (IMPDH1) | 1 | J05272 | + | + | + | + | | | | |
| IMP (inosine monophosphate) dehydrogenase 2 (IMPDH2) | 2 | L39210 | * | + | + | + | | + | | |
| inc finger protein 151 (pHZ-67) (ZNF151) | 1 | Y09723 | + | + | + | + | | + | | |
| inc finger protein, C2H2, rapidly turned over (ZNF20) | 1 | AF011573 | | + | + | | | | | |
| inducible poly(A)-binding protein (IPABP) | 1 | U33818 | + | + | + | + | | + | | |
| inducible poly(A)-binding protein (IPABP) (low match) | 1 | U33818 | | | | | | | | |

WO 00/40749

| inducible protein (Hs.80313) inhibitor of DNA binding 2, 4 M97796 + + + + + + + + + + + + + + + + + + + | |
|---|--|
| dominant negative helix- loop-helix protein (ID2) Inhibitor of kappa light 2 AF044195 polypeptide gene enhancer in B-cells, kinase complex- associated protein (IKBKAP) Inositol 1,3,4-trisphosphate 1 U51336 + + + + + + | |
| inhibitor of kappa light 2 AF044195 polypeptide gene enhancer in B-cells, kinase complex-associated protein (IKBKAP) inositol 1,3,4-trisphosphate 1 U51336 + + + + + + | |
| Inositol 1,3,4-tnsphosphate 1 U51336 + + + + + + + | |
| 5/6-kinase | |
| inositol 1,4,5 trisphosphate 1 U23850 + + + + receptor type 1 (ITPR1) | |
| inositol 1,4,5-trisphosphate 2 X57206 B + + + + 3-kinase B (ITPKB) | |
| inositol monophosphatase 1 S38980 | |
| inositol polyphosphate-5- 2 U84400 + + + + + + + hosphatase, 145kD (INPP5D) | |
| Ins(1,3,4,5)P4-binding 1 X89399 + + + + + protein + | |
| insulin-like growth factor 2 5 Y00285 + + + + + + receptor (IGF2R) | |
| integral membrane protein 1 L38961 + + + + 1 (ITM1) | |
| integral membrane protein 1 AF038953 T + + + + + 2C (iTM2C) | |
| integral membrane protein 3 U61734 + + + + + + + + Tmp21-I (p23) | |
| integrin beta 4 binding 2 AF047433 + + + protein (ITGB4BP) | |
| integrin, alpha 2b (platelet 3 M34480 + + + glycoprotein lib of lib/illa complex, antigen CD41B) (ITGA2B) | |
| integrin, alpha 5 4 X06256 + + + + + + + + + + + + + + + + + + + | |
| intégrin, alpha L (antigen 6 Y00796 CD11A (p180), lymphocyte function-associated antigen 1; alpha polypeptide) (ITGAL) | |
| integrin, alpha M 1 M18044 (complement componentreceptor 3, alpha; also known as (CD11b (p170), macrophage antigen alpha polypeptide) (ITGAM) | |
| | |
| integrin, alpha X (antigen 1 M81695 + + + + + CD11C (p150), alpha polypeptide) (ITGAX) | |
| CD11C (p150), alpha polypeptide) (ITGAX) Integrin, beta 1 (fibronectin 2 X07979 receptor, beta polypeptide, antigen CD29 includes MDF2 MSK12) (ITGB1) | |
| CD11C (p150), alpha polypeptide) (ITGAX) Integrin, beta 1 (fibronectin 2 X07979 receptor, beta polypeptide, antigen CD29 includes | |
| CD11C (p150), alpha polypeptide) (ITGAX) Integrin, beta 1 (fibronectin 2 X07979 receptor, beta polypeptide, antigen CD29 includes MDF2 MSK12) (ITGB1) Integrin, beta 2 (antigen 32 M15395 + + + + + + + + + + + + + + + + + + + | |
| CD11C (p150), alpha polypeptide) (ITGAX) Integrin, beta 1 (fibronectin 2 X07979 receptor, beta polypeptide, antigen CD29 includes MDF2 MSK12) (ITGB1) Integrin, beta 2 (antigen 32 M15395 + + + + + + + + + + + + + + + + + + + | |
| CD11C (p150), alpha polypeptide) (ITGAX) integrin, beta 1 (fibronectin 2 X07979 receptor, beta polypeptide, antigen CD29 includes MDF2 MSK12) (ITGB1) integrin, beta 2 (antigen 32 M15395 + + + + + + + + + + + + + + + + + + + | |

PCT/CA00/00005

| WO 00/40/49 | | | | | | | | | |
|---|----|----------|-----------|------|----------|---|---|----------|---|
| intercellular adhesion molecule 3 (ICAM3) | 6 | X69819 | + | | | | | + | |
| intercellular adhesion molecule 4, Landsteiner- Wiener blood group (ICAM4) | 1 | L27670 | | | | | | + | |
| Interferon consensus sequence binding protein 1 (ICSBP1) | 1 | M91196 | W, T | lymp | homa | | | | |
| Interferon consensus sequence binding protein 1 (ICSBP1) (low match) | 1 | M91196 | | | | | | | |
| interferon regulatory factor 2 (IRF2) | 4 | X15949 | + | + | + | + | | | |
| interferon regulatory factor1 (IRF1) | 4 | L05072 | + | + | + | + | | + | |
| interferon regulatory factor5 (IRF5) | 1 | U51127 | + | + | | + | | | |
| interferon, gamma- inducible protein 16 (IFI16) | 2 | M63838 | + | + | + | + | | + | |
| interferon, gamma- inducible protein 30 (IFI30) | 9 | J03909 | + | + | | + | | + | |
| INTERFERON-INDUCED GUANYLATE-BINDING PROTEIN 1 (GUANINE NUCLEOTIDE-BINDING PROTEIN 1) (non-exact 62%) | | P32455 | | | | | | | |
| interferon-induced protein 17 (IFI17) | 3 | X84958 | | + | + | + | | + | |
| interferon-induced protein 54 (IFI54) | 5 | M14660 | | | | | | | |
| interferon-inducible (1-8D) | 5 | X57351 | T | | + | | + | + | |
| interferon-inducible (1-8U) | 1 | X57352 | | + | + | | - | + | |
| interferon-related developmental regulator 1 (IFRD1) | 5 | Y10313 | | | Ĺ | | | | |
| interferon-stimulated transcription factor 3, gamma (48kD) (ISGF3G) | 2 | M87503 | | + | | + | | + | |
| interleukin 1 receptor, type II (IL1R2) | 1 | U64094 | | | | + | | | |
| Interleukin 10 receptor, beta (I.10RB) | 1 | U08988 | Tactivate | ed | + | | | + | |
| interleukin 12 receptor, beta 1 (IL12RB1) | 2 | U03187 | + | | <u> </u> | _ | + | + | only found in T cell |
| interleukin 13 receptor, alpha 1 (IL13RA1) | 2 | Y09328 | | + | + | + | _ | _ | |
| interleukin 16 (lymphocyte chemoattractant factor) (IL16) | 6 | U82972 | | _ | | | | | |
| interleukin 18 receptor 1 (IL18R1) | 1 | U43672 | | | | | | | |
| interleukin 2 receptor, beta (IL2RB) | 9 | M26062 | | | ļ | | | + | |
| interleukin 2 receptor, gamma (severe combined immunodeficiency) (IL2RG) | 6 | D11086 | + | | + | | | | |
| interleukin 4 receptor (IL4R) | 3 | X52425 | + | + | | + | | + | |
| interleukin 6 receptor (IL6R) | 5 | X12830 | | + | | | Ŀ | + | |
| interleukin 6 signal transducer (gp130, oncostatin M receptor) (IL6ST) | 1 | M57230 | | | | | | | |
| interleukin 7 receptor (IL7R) | 14 | M29696 | + · | | | | | + | |
| interleukin 7 receptor (IL7R) (low match) | 1 | AF043123 | | | | | | | |
| interleukin 8 (IL8) | 8 | Y00787 | + | | + | | + | | High in activated T cells, bone and pancreatic islets |
| | | <u> </u> | ζ | | | l | | <u> </u> | pancreatic islets |

35

| WO 00/40749 | | | | | | | | r | C1/CA00/00005 |
|--|----|--------------------|-----------|-----|---|----------|----------|---|--------------------|
| interleukin 8 receptor alpha (ILBRA) | 11 | L19591 | | | | | | | |
| interleukin 8 receptor, beta (IL8RB) | 14 | M94582 | | | | | | | |
| interleukin enhancer binding factor 2, 45kD (ILF2) | 3 | U10323 | + | + | + | + | + | + | high in uterus |
| interleukin enhancer binding factor 3, 90kD (ILF3) | 2 | U10324 | | | | | | | |
| interleukin-1 receptor- associated kinase 1 (IRAK1) | 2 | L76191 | | + | + | + | | + | |
| interleukin-1 receptor- associated kinase 1 (low match) | 1 | U52112 | | | | | | | |
| interleukin-10 receptor, alpha (IL10RA) | 5 | U00672 | + | + | + | + | | | |
| interleukin-11 receptor, alpha (IL11RA) | 7 | Z38102 | | + | + | | | | |
| INTERLEUKIN-14 PRECURSOR (IL-14) (HIGH MOLECULAR WEIGHT B-CELL GROWTH FACTOR) (HMW-BCGF) (non-exact 46%) | 1 | P40222 | | | | | | | |
| intestinal carboxylesterase; liver carboxylesterase-2 (ICE) | 1 | U60553 | | + | | | + | | |
| inversin protein (non-exact 52%) | 1 | AF084367 | | | | | | | |
| IQ motif containing GTPase activating protein 1 (IQGAP1) | 6 | L33075 | | | | | | | |
| IQ motif containing GTPase activating protein 2 (IQGAP2) | 1 | U51903 | | + | | + | | | |
| isocitrate dehydrogenase 1 (NADP+), soluble (IDH1) | 1 | AF020038 | + | + | + | + | + | + | |
| isocitrate dehydrogenase 2 (NADP+), mitochondrial (IDH2) | 2 | X69433 | + | + | + | + | + | + | |
| isocitrate dehydrogenase 3 (NAD+) alpha (IDH3A) | 2 | U07681 | | | + | | | | |
| isocitrate dehydrogenase 3 (NAD+) gamma (IDH3G) | 1 | Z68907 | + | + | + | + | | + | |
| isolate Aus3 cytochrome b (CYTB) | 1 | AF042516 | | | | | | | |
| receptor (CCR5) | 1 | AF011524 | | | | _ | <u> </u> | | |
| isopentenyl-diphosphate delta isomerase (IDI1) | 5 | X17025 | + | + | + | + | | + | |
| Janus kinase 1 (a protein tyrosine kinase) (JAK1) | 4 | M64174 | + | + | + | + | | + | |
| Janus kinase 2 (a protein tyrosine kinase) (JAK2) | 1 | AF005216 | · <u></u> | | | | | | |
| Jk-recombination signal binding protein (RBPJK) | 2 | L07876 | | | | <u> </u> | | | |
| JM1 protein jumonji (mouse) homolog | 1 | AJ005890 U57592 | | + + | + | + | | + | |
| (JMJ) jun D proto-oncogene | 1 | X51346 | + | + | + | + | <u> </u> | + | |
| (JUND) jun dimerization protein | 1 | AF111167 | | - | | - | _ | | only found in germ |
| junction plakoglobin (JUP) | 1 | M23410 | | + | + | + | | + | |

| WO 00/40/49 | | | | | | | | | |
|---|----------------|----------|--|--|----------|--|--|----------|---------------------------------------|
| kangai 1 (suppression of tumorigenicity 6, prostate; CD82 antigen (R2 leukocyte antigen, antigen detected by monoclonal and antibody | 1 | U20770 | + | + | + | + | + | + | |
| lA4)) (KAI1) karyophenn (importin) beta 1 (KPNB1) | 2 | L39793 | + | + | + | + | + | + | |
| karyopherin (importin) beta 2 (KPNB2) | 1 | U72395 | + | + | + | + | | | <u> </u> |
| karyopherin alpha 1 | 1 | S75295 | + | + | + | | + | 一 | |
| (importin alpha 5) (KPNA1) karyopherin alpha 2 (RAG | 1 | U09559 | | | _ | - | | \vdash | |
| cohort 1, importin alpha 1) (DPNA2) | | | | | | | | | |
| karyopherin alpha 3 (importin alpha 4) (KPNA3) | 1 | D89618 | | + | | | + | | |
| karyopherin alpha 4 (KPNA4) | 1 | M17887 | | + | + | | | | |
| Katanin (80 kDa) (KAT) | 1 | AF052432 | | + | + | + | | + | · · · · · · · · · · · · · · · · · · · |
| KE03 protein | 2 | AF064604 | | t | | 1 | | ╁ | |
| Kelch-like ECH-associated protein 1 (KIAA0132) (66%aa) | 1 | D50922 | | | | | | | |
| Keratin 8 (KRT8) | 1 | X74929 | | + | + | + | + | + | |
| ketohexokinase (fructokinase) (KHK) | 7 | X78678 | | + | | + | + | <u> </u> | |
| KIAA0001 (KIAA0001) | | Q15391 | | | | - | | - | |
| (72% aa) KIAA0001 (KIAA0001) | 1 | Q15391 | | | | ┢ | - | | |
| (76% aa) KIAA0001 (KIAA0001) | 1 | Q15391 | | | \vdash | ├- | ļ | | |
| (non-exact 72%) | 5 | D13627 | | + | + | + | <u> </u> | + | ļ |
| KIAA0002 (KIAA0002) | 4 | D13627 | ļ <u>.</u> | + | + | + | <u> </u> | + | |
| KIAA0005 (KIAA0005) | - 1 | D13635 | | + | <u> </u> | <u> </u> | <u> </u> | + | |
| KIAA0010 (KIAA0010) KIAA0016 (KIAA0016) | | D13633 | + | + | + | + | ├ | + | |
| KIAA0017 (KIAA0017) | | D13641 | ļ | T | <u> </u> | <u> </u> | <u> </u> | Ľ | |
| KIAA0022 (KIAA0022) | 2 | D14664 | _ | + | + | + | <u> </u> | | |
| , , , | 1 | D14689 | _ | + | | <u> </u> | ↓ | <u> </u> | |
| KIAA0023 (KIAA0023) | | D14689 | + | + | - | + | <u> </u> | + | ļ |
| KIAA0024 (KIAA0024) | 1 | | + | | <u></u> | | <u> </u> | <u> </u> | |
| KIAA0025 (KIAA0025) | 1 | D14695 | | + | + | + | + | + | |
| KIAA0026 (KIAA0026) | 2 | D14812 | | + | + | + | | + | |
| KIAA0027 | 1 | D25217 | | + | <u> </u> | <u> </u> | ļ | | |
| KIAA0032 (KIAA0032) | 2 | D25215 | | + | + | + | | | |
| KIAA0040 (KIAA0040) | 1 | D25539 | + | + | + | + | | + | |
| KIAA0050 (KIAA0050) | 4 | D26069 | | | | <u> </u> | | L | |
| KIAA0053 (KIAA0053) | 17 | D29642 | + | | + | + | | <u> </u> | |
| KIAA0057 (KIAA0057) | 1 | D31762 | + | | + | + | + | + | high in fetal lung |
| KIAA0058 (KIAA0058) | 11 | D31767 | + | <u> </u> | + | + | | + | |
| KIAA0063 (KIAA0063) | 3 | D31884 | + | + | + | + | <u> </u> | + | ļ |
| KIAA0064 (KIAA0064) | | D31764 | + | + | + | + | | + | |
| KIAA0066 | 1 | D31886 | + | + | + | + | L. | + | |
| KIAA0068 | 1 | D38549 | | + | + | + | + | + | |
| KIAA0073 | 3 | D38552 | | + | + | + | <u> </u> | + | |
| KIAA0081 | 2 | D42039 | | + | | + | L_ | + | |
| KIAA0084 | 2 | D42043 | + | + | * | + | L | + | |
| KIAA0085 | 26 | U30498 | + | + | + | + | + | + | |
| KIAA0088 | 3 | D42041 | + | + | + | + | + | + | |
| KIAA0090 | 2 | D42044 | + | + | + | + | + | + | |
| KIAA0092 (KIAA0092) | 1 | D42054 | L | + | + | + | | + | |
| | | | | | | | | | |

| WO 00/40749 | | | | | | | | | C1/CX00/00005 |
|------------------------------------|--------------|----------|---------------|------------------|--|--|--|--------------|---------------|
| KIAA0094 | 3 | D42084 | | | + | + | | | |
| KIAA0095 (KIAA0095) | 1 | D42085 | | | | | | | |
| KIAA0096 | 1 | D43636 | + | + | + | + | | + | |
| KIAA0097 (KIAA0097) | 1 | X92474 | T | + | + | | + | | |
| KIAA0099 (KIAA0099) | 3 | D43951 | + | + | + | + | + | + | |
| KIAA0102 (KIAA0102) | 2 | D14658 | | + | | + | + | + | |
| KIAA0105 | 1 | D14661 | В | + | | | + | + | |
| KIAA0120 | 2 | P37802 | | | | | | | |
| KIAA0120 (non-exact, 65%) | 1 | M83106 | | | | | | | |
| KIAA0121 (KIAA0121) | 1 | D50911 | + | + | + | + | | + | |
| KIAA0123 | 1 | D21064 | | + | + | + | | + | |
| KIAA0128 | 1 | D50918 | + | + | + | + | | + | _ |
| KIAA0129 (KIAA0129) | 1 | D50919 | + | + | + | + | | Π | |
| KIAA0130 (KIAA0130) | 1 | AF055995 | | + | + | + | | | |
| KIAA0136 | 2 | D50926 | | | | | | | |
| KIAA0137 (KIAA0137) | 1 | AB004885 | | + | + | + | | + | |
| KIAA0140 (KIAA0140) | 1 | D50930 | + | + | | + | | + | |
| KIAA0141 (KIAA0141) | 3 | D50931 | - | 1 | | T | 1 | | |
| KIAA0144 (KIAA0144) | 3 | D63478 | + | + | + | + | | + | |
| KIAA0144 (KIAA0144) (Iow | 1 | D63478 | | | | | | | |
| match) KIAA0144 (non-exact 61%) | | Q14157 | | ╁ | | | | ╁ | |
| KIAA0144 (non-exact 65%) | i | Q14157 | | | | ├─ | ├ | | |
| KIAA0146 | | D63480 | | +- | + | + | - | + | |
| KIAA0148 (KIAA0148) | - | D63482 | ļ | + | | \vdash | - | + | |
| KIAA0154 | 2 | D63876 | + | + | + | + | - | + | |
| KIAA0156 | 1 | D63879 | | + | + | + | ₩ | + | |
| KIAA0160 | 2 | D63881 | | + | | ┼ | - | - | |
| | 1 | D79983 | + | + | | + | - | | |
| KIAA0161 (KIAA0161) | 3 | D79986 | | ļ <u> </u> | - | <u> </u> | | - | |
| KIAA0164 (KIAA0164) | 1 | D79989 | | + | | | ┼ | ┼ | <u> </u> |
| KIAA0167 (KIAA0167) | | D79990 | <u> </u> | ┿ | + | + | - - | + | |
| KIAA0168 (KIAA0168) | 3 | D79991 | | ' - | <u> </u> | ↓ : | ╂ | | |
| KIAA0169 | 3 | _ | | + | + | + | ↓ | + | |
| KIAA0171 (KIAA0171) | 3 | D79993 | ļ | | + | + | ـــــ | + | ļ |
| KIAA0174 (KIAA0174) | 7 | D79996 | + | + | 1 | + | | + | , |
| KIAA0179 | 2 | D80001 | | + | + | | | | <u> </u> |
| KIAA0181 | 1 | D80003 | | + | + | + | | + | ļ |
| KIAA0183 | 4 | D80005 | + | + | + | + | + | + | |
| KIAA0184 | 1 | D80006 | + | + | + | + | ↓ | + | |
| KIAA0191 (72% aa) | 1 | D83776 | | | ļ | | $oldsymbol{ol}}}}}}}}}}}}}}}}}}$ | <u> </u> | |
| KIAA0191 (non-exact 77%) | 1 | | | <u> </u> | | | <u> </u> | | |
| KIAA0193 (KIAA0193) | 1 | D83777 | + | + | + | + | $oldsymbol{ol}}}}}}}}}}}}}}}}}$ | + | |
| KIAA0200 (KIAA0200) | | D83785 | | T + | + | + | | + | |
| KIAA0210 (KIAA0210) | 3 | D86965 | | | | | | | |
| KIAA0217 | 2 | D86971 | + | + | + | + | | + | |
| KIAA0219 | 2 | U77700 | | + | + | + | | + | |
| KIAA0222 (KIAA0222) | 1 | D86975 | | | | | | | |
| KIAA0223 | 2 | D86976 | | | | | | T | |
| KIAA0229 | 1 | D86982 | + | + | T | | 1 | | |
| KIAA0232 (KIAA0232) | 1 | D86985 | | + | + | + | | + | |
| KIAA0233 (KIAA0233) | 1 | D87071 | | 1 | 1 | 1 | 1 | T | |
| KIAA0235 | 2 | D87078 | + | + | + | + | \top | † | |
| KIAA0239 | 1 | D87076 | + | + | | T | + | \top | |
| | | 1 | | | | | | | |

| KIAA0239 (non-exact 80%) | 1 | D87076 | | | | | | | |
|--|--------------|----------|-------------|-----|---|----------|--|----------|-------------|
| KIAA0240 | 1 | D87077 | | | | | | | |
| KIAA0242 | 4 | D87684 | + | + | + | + | + | + | |
| KIAA0248 | 2 | D87435 | | + | + | + | | + | |
| KIAA0249 (KIAA0249) | 3 | D87436 | + | + | + | + | | + | |
| KIAA0253 | 5 | D87442 | + | + | + | + | + | + | |
| KIAA0254 (KIAA0254) | 1 | D87443 | · | + | + | + | | | |
| KIAA0255(KIAA0255) | 4 | D87444 | | + | + | + | | + | |
| KIAA0262 (KIAA0262) | 3 | D87451 | + | + | + | + | | + | |
| KIAA0263 (KIAA0263) | 1 | D87452 | + | + | + | + | | + | |
| KIAA0264 | 3 | D87453 | | + | + | + | | + | |
| KIAA0268 | 1 | D87742 | + | ++ | | + | _ | + | |
| KIAA0269 | 1 | Q92558 | | | | | _ | | |
| KIAA0275 (KIAA0275) | 13 | D87465 | + | + | | + | - | + | |
| KIAA0304 (KIAA0304) | 2 | AB002302 | + | ++ | + | + | + | + | |
| KIAA0308 | 2 | AB002306 | | + | + | - | | + | |
| KIAA0310 (KIAA0310) | | AB002308 | | + + | + | + | | + | |
| KIAA0314 (=U96635 M.musculus ubiquitin protein ligase Nedd-4) | 3 | AB002312 | | | | | | | |
| KIAA0315 (KIAA0315) | 4 | AB002313 | | + | + | + | + | + | |
| KIAA0325 (=L08505 R.norvegicus cytoplasmic dynein heavy chain (MAP 1C)) | 2 | AB002323 | | | | | | | |
| KIÁA0329 (KIAA0329) | 1 | AB002327 | | + | + | + | | + | |
| KIAA0330 | 1 | AB002328 | + | + | + | | | + | |
| KIAA0332 | 1 | AB002330 | | + | + | + | | + | |
| KIAA0333 | 2 | AB002331 | | + | + | + | + | + | |
| KIAA0336 (KIAA0336) | 3 | AB002334 | + | + | + | + | | + | |
| KIAA0336 (KIAA0336) (low match) | 1 | AB002334 | | | | | | | |
| KIAA0342 (KIAA0342) | 1 | AB002340 | | + | + | | | + | |
| KIAA0344 (KIAA0344) | 2 | AB002342 | | | | + | | + | |
| KIAA0354 (KIAA0354) | 1 | AB002352 | + | + | + | + | | + | |
| KIAA0365 (KIAA0365) | 3 | AB002363 | + | + | + | + | + | + | |
| KIAA0370 | 6 | AB002368 | | + | + | + | + | + | |
| KIAA0372 (KIAA0372) | 1 | AB002370 | | | | | Ι | | |
| KIAA0373 (KIAA0373) | 1 | AB002371 | | + | | + | | | |
| KIAA0375 (KIAA0375) | 1 | AB002373 | | + | | + | | | |
| KIAA0377 (KIAA0377) | 1 | AB002375 | | + | | + | + | | |
| KIAA0379 | 1 | AB002377 | | | | + | | | |
| KIAA0379 (non-exact, 65%) | 1 | AB002377 | | | | | | | |
| KIAA0380 (KIAA0380) | 1 | AB002378 | + | + | | + | | + | |
| KIAA0380 (KIAA0380) (60%aa) | 1 | AB002378 | | | | | | | |
| KIAA0382 (KIAA0382) | 2 | AB002380 | | + | + | + | <u> </u> | + | |
| KIAA0383 | 1 | AB002381 | | ļ | | | | <u> </u> | |
| KIAA0386 (KIAA0386) | 5 | AB002384 | | | | | | <u> </u> | |
| KIAA0392 | 1 | AB002390 | | | | L_ | | <u> </u> | |
| KIAA0397 (KIAA0397) | 4 | AB007857 | | + | + | + | + | + | |
| KIAA0403 | 3 | AB007863 | | | | | | | |
| KIAA0404 | 1 | AB007864 | | + | | + | | | |
| KIAA0409 | 1 | AB007869 | | + | | + | | | |
| KIAA0421 | 1 | AB007881 | + | + | + | | | + | |
| KIAA0424 (non-exact 82%) | 1 | AB007884 | | | | | | | |

| KIAA0428 (KIAA0428) | 9 | Y13829 | | | | | | | |
|---------------------------------------|--|------------|-----|----------|----------|--|--|----------|--|
| KIAA0429 (KIAA0429) | 2 | AB007889 | + | + | + | + | | + | |
| KIAA0430 (KIAA0430) | 2 | AB007890 | | | | | | | only in ovary |
| KIAA0432 (KIAA0432) | 2 | U86753 | | + | + | | | | |
| KIAA0435 (KIAA0435) | | AB007895 | | | | | | | |
| KIAA0438 (KIAA0438) | 1 | AB007898 | | + | + | + | | + | |
| KIAA0447 (KIAA0447) | 3 | AB007916 | + | + 1 | + | + | | + | |
| KIAA0449 | 1 | AB007918 | | + | | | | + | |
| KIAA0456 | 1 | AB007925 | | + | + | + | | + | |
| KIAA0458 (KIAA0458) | 1 | AB007927 | | | | | | | |
| KIAA0462 | 1 | AB007931 | + | + | + | + | | + | |
| KIAA0465 | | AB007934 | | + | + | + | + | + | |
| KIAA0476 (KIAA0476) | 1 | AB007945 | | + | + | + | | | |
| KIAA0489 | 1 | AB007958 | | + | | | | - | |
| KIAA0494 (KIAA0494) | 1 | AB007963 | + | + | + | + | | + | |
| KIAA0515 | | AB011087 | + | + | + | + | | + | |
| KIAA0521 | 3 - | AB011093 | - + | ++ | | | \vdash | + | |
| KIAA0525 | 1 | AB011097 | | +- | | + | | | |
| KIAA0530 | | AB011102 | | + | + | + | | - | |
| KIAA0532 | ' | AB011104 | + | + | + | + | - | + | |
| KIAA0537 (KIAA0537) | | AB011109 | · | | | | ├ | - | |
| KIAA0540 | | AB011103 | + | + | + | + | | + | |
| · ··· · · · · · · · · · · · · · · · · | | AB011115 | | <u> </u> | + | +- | | + | <u> </u> |
| KIAA0543 | l ' | AB011116 | | + | + | · · | | + | |
| KIAA0544 | 1 | AB011110 | | + | + | + | | + | |
| KIAA0549 | 2 | 1 | | + | | <u> </u> | - | + | |
| KIAA0551 | 2 | AB011123 | | + | + | + | ļ | + | |
| KIAA0554 | 8 | AB011126 | | <u> </u> | | + | | Ι, | |
| KIAA0561 | 1 | AB011133 | | + | | <u> </u> | ļ | <u> </u> | |
| KIAA0562 (KIAA0562) | 1 | AB011134 | | ļ | | L | ļ | <u> </u> | |
| KIAA0563 (KIAA0563) | 1 | AB011135 | | <u> </u> | <u></u> | <u> </u> | | Ļ | ļ |
| KIAA0569 (KIAA0569) | 2 | AB011141 | | + | + | + | | + | ļ |
| KIAA0571 (KIAA0571) | 2 | AB011143 | | + | + | + | <u> </u> | <u> </u> | |
| KIAA0573 | 1 | AB011145 | | + | | + | <u> </u> | + | |
| KIAA0576 | 1 | AB011148 | | <u> </u> | | 1 | <u> </u> | | |
| KIAA0580 | 1 | AB011152 | | | | l | | | · · · · · · · · · · · · · · · · · · · |
| KIAA0584 | 1 | AB011156 | | + | | <u> </u> | | | |
| KIAA0592 | 3 | AB011164 | + | + | + | + | | + | |
| KIAA0596 | 1 | AB011168 | | + | + | | | | |
| KIAA0598 (KIAA0598) | 1 | AB011170 | | + | + | + | 1 | | |
| KIAA0608 | 1 | AB011180 | | 1 | + | + | | | |
| KIAA0614 | 2 | AB014514 | + | + | + | + | | + | |
| KIAA0615 (KIAA0615) | 1 1 | AB014515 | | | | | Ì | | |
| KIAA0621 | 1 1 | AB014521 | | + | + | \Box | T | + | |
| KIAA0648 | 1 | AB014548 | | + | + | + | | + | |
| KIAA0652 (KIAA0652) | 1 1 | AB014552 | + | + | + | + | | + | |
| KIAA0668 | 1 1 | AB014568 | | +- | <u> </u> | | <u> </u> | T | |
| KIAA0669 | + + - | AB014569 | | + | t | \vdash | | 1 | |
| KIAA0671 (KIAA0671) | 1 | AB014571 | | + | + | + | - | + | |
| KIAA0675 (KIAA0675) | | AB014575 | | + | + | + | + | +- | |
| KIAA0676 | 1 1 | AB014576 | | + | + | + | \vdash | + | |
| KIAA0677 (KIAA0677) | 2 | AB014577 | | ++ | + | + | + | + | |
| ואוראסטוו (ואוראסטטוו) | | AB014578 | + | | <u> </u> | <u> </u> | ₩- | + | |
| KIAA0678 | | L ARIDAS/A | + | + | + | + | | 1 7 | |

| WU 00/40/49 | | | | | | | | | |
|--|------------------|------------|------|----|--|--|--|--------------|----------------------------------|
| KIAA0680 (KIAA0680) | 1 | AB014580 | | | | | | | |
| KIAA0692 | 1 1 | AB014592 | + | + | + | + | | + | |
| KIAA0697 | 1 | AB014597 | | | | | | | |
| KIAA0699 | 1 | AB014599 | + | + | + | + | | + | |
| KIAA0700 | 1 | AB014600 | | + | + | + | | + | |
| KIAA0737 (KIAA0737) | 3 | AF014837 | + | + | + | + | | + | |
| KIAA0748 (KIAA0748) | 2 | AB018291 | | + | | | | | |
| KIAA0763 (KIAA0763) | 2 | AB018306 | + | + | + | + | | + | |
| KIAA0769 (KIAA0769) | 2 | AB018312 | | + | + | + | | + | |
| KIAA0782 | 1 | AB018325 | + | + | | + | | | high in BPH stroma |
| KIAA0796 | 1 | AB018339 | | + | + | + | | + | |
| KIAA0798 (KIAA0798) | 1 | AB018341 | | | | | | | |
| KIAA0823 | 1 | AB020630 | | | | | | | |
| KIAA0854 | 1 | AB020661 | + | + | + | + | | + | |
| KIAA0856 | 1 | AB020663 | | + | + | + | | + | |
| KIAA0860 | 1 | AB020667 | | + | \vdash | + | | 一 | |
| KIAA0862 | 1 | AF054828 | | + | + | + | | | |
| KIAA0871 (non-exact 88%) | | AB020678 | | + | | | | \vdash | |
| KIAA0873 | - i - | AB020680 | | ++ | + | + | | + | |
| KIAA0892 | | AB020699 | + | + | + | + | | + | |
| KIAA0906 | - | AB020713 | + | + | + | + | - | + | |
| KIAA0991 | <u> </u> | AB023208.1 | | + | | \vdash | | | |
| killer cell lectin-like | ···· | U11276 | | | + | + | - | + | |
| receptor subfamily B, member 1 (KLRB1) | ' | 011270 | | | | | | | |
| killer cell lectin-like receptor subfamily C, member 4 (KLRC4) | 1 | U96846 | | | | | l | | |
| kinectin 1 (kinesin receptor) (KTN1) | 7 | D13629 | | | | | | | |
| kinesin family member 5B (KIF5B) | 2 | X65873 | | + | + | + | ļ | L. | |
| kinesin-like DNA binding protein | 1 | AB017430 | + | + | + | + | | + | |
| Krueppel-related DNA- binding protein (TF6) (low match) | 1 | M61869 | ···- | | | | | | |
| Kruppel related gene (clone pHKR1RS) | 1 | M20675 | | | | | | | |
| Kruppel-like zinc finger protein Zf9 | 3 | U51869 | + | + | + | + | + | + | |
| Kruppel-like zinc finger protein Zf9 (non-exact 76%) | 1 | U44975 | | + | + | | + | + | |
| kruppel-type zinc finger protein, ZK1 | 1 | AB011414.1 | | | | | | | |
| L apofernitin | 3 | X03742 | | | | | | | |
| lactate dehydrogenase A (LDHA) | 3 | X02152 | | + | + | + | + | + | |
| lactate dehydrogenase A (LDHA) (non-exact, 81%) | 1 | X02152 | | | | | | | high in fatal lung |
| lactate dehydrogenase B (LDHB) | 6 | X13794 | + | + | + | + | + | + | high in fetal lung fibrablast |
| lactotransferrin (LTF) | 1 | U07643 | + | ļ | <u> </u> | + | <u> </u> | + | high in bone marrow |
| laminin binding protein (low score) | 1 | D28372 | | ļ | | | <u> </u> | | |
| laminin receptor 1 (67kD); Ribosomal protein SA (LAMR1) | 20 | X15005 | + | + | + | + | + | + | high in many libraries |
| laminin receptor homolog {3' region} | - | \$35960 | | | | | | | |
| laminin, gamma 1 (formerly LAMB2) (LAMC1) | 2 | J03202 | + | + | + | | | + | |

| WO 00/40749 | | | | | | | _ | | |
|---|---|----------|------|----|---|---|----|---|-----------------------|
| latent transforming growth factor beta binding protein 1 (LTBP1) | 2 | M34057 | | + | + | + | | + | |
| LAZ3/BCL6 (=Z79582;D28522/4) | 1 | Z79581 | | | | | | | |
| LDLC | 2 | Z34975 | + | + | + | + | | + | |
| lecithin-cholesterol acyltransferase (LCAT) (non-exact, 66%) | 1 | M17959 | | | | | | | |
| lectin, galactoside-binding, soluble, 2 (galectin 2) (LGALS2) | 1 | M87842 | | | | + | | | |
| Tectin, galactoside-binding, soluble, 3 binding protein (galectin 6 binding protein) (LGALS3BP) | 1 | L13210 | + | + | + | + | | + | |
| leucine rich repeat (in FLII) interacting protein 1 (LRRFIP1) | 5 | AJ223075 | + | + | + | + | + | + | |
| leucocyte immunoglobulin- like receptor-5 (LIR-5) | 2 | AF072099 | | | | + | | | |
| leucocyte immunoglobulin- like receptor-6a (LIR-6) | 7 | AF025530 | | | | | | | |
| leucocyte immunoglobulin- like receptor-7 (LIR-7) | 2 | U82275 | | + | | | | | only found in CNS |
| leukemia virus receptor 1 (GLVR1) | 1 | L20859 | + | + | + | + | | + | |
| leukocyte adhesion protein p150,95 alpha subunit | 1 | M29484 | | | | | | | |
| leukocyte antigen, HLA-A2 | 3 | Y13267 | | 1 | | | | | |
| leukocyte immunoglobulin- like receptor (MIR-10) | 3 | AF025528 | | + | | | | | |
| leukocyte tyrosine kinase (LTK) | 1 | X60702 | + | | | | | | found only in blood |
| leukocyte-associated Ig- like receptor 1 (LIAR1) | 3 | AF013249 | | | | + | | | |
| leukotriene A4 hydrolase (LTA4H) | 6 | J03459 | + | + | + | + | + | + | |
| leupaxin (LDPL) | 2 | AF062075 | + | | | + | | + | |
| ligase I, DNA, ATP- dependent (LIG1) | 1 | M36067 | В, Т | + | + | | + | + | |
| LIM and SH3 protein 1 (LASP1) | 2 | X82456 | + | + | + | + | + | + | t |
| LIM domain kinase 2 (LIMK2) | 2 | AC002073 | + | + | + | + | | + | |
| line-1 protein | 1 | | | | | | | | 75 |
| Line-1 repeat mRNA with 2 open reading frames | 1 | U93566 | + | + | + | + | + | + | |
| Line-1 repeat with 2 open reading frames | 1 | M22332 | + | +_ | + | + | + | + | high in gastric tumor |
| LINE-T REVERSE TRANSCRIPTASE HOMOLOG | 1 | P08547 | | | | | | | |
| lipase A, lysosomal acid, cholesterol esterase (Wolman disease) (LIPA) | 4 | X76488 | + | + | + | + | | + | |
| lipase, hormone-sensitive (LIPE) | 1 | L11706 | + | + | | | | + | |
| ĽMP7 | 1 | L11045 | | L | L | | L. | | |
| Lon protease-like protein (LONP) | 2 | X74215 | + | + | + | + | ŀ | + | |
| low density lipoprotein- related protein 1 (alpha-2- macroglobulin receptor) (LRP1) | 2 | AF058414 | | | | | + | | only in liver |
| low density lipoprotein- related protein-associated protein 1 (alpha-2- macroglobulin receptor- associated protein 1) (LRPAP1) | | M63959 | | + | + | | + | + | |

| Invariant Inva | WO 00/40/49 | | | | | | | | | |
|--|--|----|----------|---|----------|---------|-----|------|------|------|
| macroplobulin receptor associated protein 1) (LRPAP1) (non-exact, 75%) Gow-afmily F-c-gamma | related protein-associated | 1 | M63959 | | | | | | | |
| (LRPAP1) (non-exact, 75%) low-elfinity Fc-gamma | macroglobulin receptor- | | | | | | | | | |
| Image: Continue Image: Con | (LRPAP1) (non-exact, | | | | | | | | | |
| LPS-induced TNF-alpha | low-affinity Fc-gamma | | L08107 | | | | | | | |
| LSH-T | LPS-induced TNF-alpha | 9 | AF010312 | + | | | | + | | |
| transporter subunit LAT1 [tung resistance-related protein (I.RP) AF011333 B CLY75) [LY75] TAF011333 B CLY75] TAF011333 TAF01 | | 1 | U00921 | + | + | + | + | | + | |
| International Content Inte | transporter subunit LAT1 | | | | | | | | | |
| (LY75) | protein (LRP) | | | | + | + | + | | + | |
| Vimphocyte antigen HLA- 8"4402 and HLA-B"5101 Yimphocyte (yotsolic protein 1 (1-plastin) (LCP1) Vimphocyte (yotsolic protein 2 (SH2 domain-containing leukocyte protein 6" (5R) (LCP2) Vimphocyte (yotsolic protein 6" (5R) (LCP2) Vimphocyte (yotsolic protein 6" (5R) (LCP2) Vimphocyte (yotsolic protein 6" (5R) (LCP2) Vimphocyte (yotsolic protein 6" (5R) (LCP2) Vimphocyte (yotsolic protein 7" (5R) (LCP2) Vimphocyte (yotsolic protein 7" (5R) (LCP2) Vimphocyte (yotsolic protein 7" (5R) (LCP2) Vimphocyte (yotsolic protein 7" (5R) (1.5P1) Vimphocyte (yotsolic protein 7" (1.5P1) Vimphocyte (yotsolic protein 7" (1.5P1) Vimphodi (yotsolic protein 7" (1.5P1) V | (ĽY75) | | | В | | | | | | |
| ### ### ### ### ### ### ### ### ### ## | | | | | | | | | | |
| | B*4402 and HLA-B*5101 | | | | | | | | | |
| profein 2 (SH2 domain-containing leukocy) protein of 76kD) (LCP2) lymphocyte glycoprofein 2 X04391 + | orotein 1 (L-plastin) (LCP1) | | | | | li casa | | | | olod |
| Imphocyte glycoprotein | protein 2 (SH2 domain- containing leukocyte | 4 | | | , | iynip | mom | а, і | acuv | aleu |
| 1 (LSP1) | lymphocyte glycoprotein T1/Leu-1 | 2 | X04391 | + | | + | | | | |
| Synophoid Syno | | 16 | M33552 | + | + | + | + | | + | |
| LyP1 | lymphocyte-specific protein tyrosine kinase (LCK) | 7 | M36881 | | + | | | | + | |
| Image: Content | LyP1 | | | | | | | | | |
| Imphoid-specific SP100 | membrane protein (LRMP) | , | | + | | + | + | | | |
| Convertase (LPC) | lymphoid-specific SP100 homolog (LYSP100-A) | , | | | | | | | | |
| PROTECTIVE PROTEIN PRECURSOR (CATHEPSIN A) (CARBOXYPEPTIDASE C) Iysosomal-associated membrane protein 1 (LAMP1) Lysosomal-associated membrane protein 2 (LAMP2) Iysozyme (renal amyloidosis) (LYZ) Iysyl-IRNA synthetase (KARS) M phase phosphoprotein 10 (U3 small nucleolar ribonucleoprotein) (MPP- 10) M1-type and M2-type pyruvate kinase m6A methyltransferase (MT-A70) mab-21 (C. elegans)-like 1 (MAB21L1) MacMarcks 1 | convertase (LPC) | | | + | + | + | + | | + | |
| Tysosomal-associated | PROTECTIVE PROTEIN PRECURSOR (CATHEPSIN A) (CARBOXYPEPTIDASE C) | 1 | P10619 | | | | | | | |
| Lysosomal-associated membrane protein 2 ((LAMP2) | lysosomal-associated membrane protein 1 | 1 | J04182 | + | + | + | + | + | + | |
| Iysozyme (renal as 39 M19045 + | Lysosomal-associated membrane protein 2 | 1 | J04183 | | + | + | + | + | + | |
| Iysyl-tRNA synthetase | lysozyme (renal amyloidosis) (LYZ) | | | | + | + | + | | | |
| M phase phosphoprotein 1 X98494 10 (U3 small nucleolar ribonucleoprotein) (MPP-10) 2 X56494 M1-type and M2-type 2 X56494 pyruvate kinase 7 AF014837 + + + (MT-A70) mab-21 (C. elegans)-like 1 U38810 + + + + (MAB21L1) MacMarcks 1 X70326 + + + + + macrophage-associated 1 Z22968 + + + + + | lysyl-tRNA synthetase | 2 | | + | + | + | + | | + | |
| M1-type and M2-type 2 X56494 pyruvate kinase 7 AF014837 + + + m6A methyltransferase 7 AF014837 + + + (MT-A70) 1 U38810 + + + + (MAB21L1) 1 X70326 + + + + MacMarcks 1 X70326 + + + + macrophage-associated 1 Z22968 + + + | M phase phosphoprotein 10 (U3 small nucleolar ribonucleoprotein) (MPP- | 1 | | | | | | | | |
| MacMarcks | M1-type and M2-type | 2 | X56494 | | | | | | | |
| mab-21 (C. elegans)-like 1 1 U38810 + <td>m6A methyltransferase</td> <td>7</td> <td></td> <td>+</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> | m6A methyltransferase | 7 | | + | | | | | | |
| MacMarcks 1 X70326 + + + + + + + + + + + + + + + + + + + | mab-21 (C. elegans)-like 1 (MAB21L1) | | | | <u> </u> | | | | | |
| | | 1 | 1 | + | | + | | + | L | |
| | macrophage-associated antigen (MM130) | 1 | Z22968 | | + | + | + | | + | |

| W O 00/40/49 | | | | | | | | | | |
|--|----|--------------|----------|--------------|--|----------------|-------------|--|---------------------------------------|-----------|
| MADS box transcription | 1 | U49020 | | + | + | + | | + | | |
| enhancer factor 2, polypeptide A (myocyte | | | | | | | | | | |
| enhancer factor 2A) | | | | 1 | | | | | | 1 |
| (MEF2A) | | | | | | | l | | |] |
| MADS box transcription | 1 | L08895 | | + | + | + | | + | | |
| enhancer factor 2, | | | | 1 1 | | | | | | |
| polypeptide C (myocyte | | | | | | | | | | 1. |
| enhancer factor 2C) (MEF2C) | | | | | | | | | | 1 |
| major cytoplasmic tRNA- | | X17516 | | | | \vdash | | | | 1 |
| Val(IAC) (=M33940) | | | | | | | | | |] |
| major histocompatibility | 1 | M95531 | | | | | | | |]; |
| complex class I beta chain | | | | | | | | | | 1 |
| (HLA-B) major histocompatibility | 41 | Z93949 | + | + | + | + | | + | high in villous | ┨ |
| complex, class I, A (HLA-A) | 41 | 255545 | • | ' | · | 1 | | | adenoma | L |
| major histocompatibility | 1 | Z72422 | | | | | | | · · · · · · · · · · · · · · · · · · · | 1 |
| complex, class I, A (HLA-A) | | | | | | | | | | H |
| (low match) | | | | ļ | | | | | | ┧. |
| major histocompatibility | 82 | M24097 | + | + | + | + | + | + | | |
| complex, class I, C (HAL- | | ļ 1 | | | | | | | | 1 |
| major histocompatibility | 77 | M20022 | + | + | + | + | | + | | 1 |
| complex, class I, E (HLA-E) | ,, | 11123022 | • | ` | | | | | | _ |
| major histocompatibility | 2 | U15085 | + | + | + | + | | + | | 1 |
| complex, class II, DM | - | | | | | | | | | 1 |
| BETA (HLA-DMB) | | 175-188 | | <u> </u> | <u> </u> | L., | | | | 4 |
| major histocompatibility | 10 | M57466 | + | + | + | + | | + | | 1 |
| complex, class II, DP beta 1 (HLA-DPB1) | | | | | | | | | | |
| major histocompatibility | 9 | V00522 | + | + | + | + | _ | + | | 1 |
| complex, class II, DR beta | J | | | | | 1 | | | | |
| 1 (HLA-DRB1) | | | | | | <u> </u> | <u> </u> | ļ | | 1 |
| Major histocompatibility | 2 | M24070 | | + | + | 1 | + | + | İ | 1 |
| complex, class II, Y box- | | | | | | ĺ | i | | | |
| binding protein I; DNA- binding protein B (YB1) | | <u> </u> | | | l | | | | | |
| malate dehydrogenase 1, | 1 | D55654 | + | + | + | + | + | + | | 1 |
| NAD (soluble) (mdh1) | • | | | <u></u> | | L | | L | | |
| malate dehydrogenase 1, | 3 | D55654 | | + | + | | + | + | | 7 |
| NAD (soluble) (MDH1) | | | | ↓ | <u> </u> | | <u> </u> | | | 4 |
| malonyl-CoA | 2 | AF097832 | | 1 | | | | | ļ | |
| decarboxylase precursor maltase-glucoamylase | 1 | AF016833 | | + | | + | - | | | \forall |
| (mg) | ' | 71 010033 | | | | ` | | | | |
| manic fringe (Drosophila) | 1 | U94352 | + | + | + | + | | + | | 1 |
| homolog (MFNG) | | | | <u>L</u> | <u></u> | L | | | | ╛ |
| mannose phosphate | 1 | X76057 | | + | + | + | | + | | |
| isomerase (MPI) | | Vanne | - | | <u> </u> | | <u> </u> | <u> </u> | | 4 |
| mannose phosphate | 2 | X76057 | | + | + | + | | + | | |
| isomerase (mpi) | 7 | X56253 | | ++ | + | \vdash | + | + | | \dashv |
| mannose-6-phosphate receptor (cation | 3 | 750233 | | 1 | ` | 1 | ' | <u> </u> | | |
| dependent) (M6PR) | | | | | L | L | | L | | J |
| mannose-P-dolichol | 1 | AF038961 | <u> </u> | + | + | + | | + | | 7 |
| utilitzation defect 1 | | | | 1 | |] | | | | |
| (MPDU1) | | TENDOE | | + | | ├ ┯ | + | + | | 4 |
| mannosidase, alpha B, | 1 | U60885 | | * | | ¯ | ~ | , T | | 1 |
| lysosomal (MANB) mannosyl (alpha-1,3-)- | 1 | M55621 | + | ++ | + | + | + | + | | \dashv |
| glycoprotein beta-1,2-N- | ' | | - | | ĺ | 1 | | l | | |
| acetylglucosaminyltransfer | | 1 | | | l | | l | 1 | | |
| ase (MGAT1) | | | | 1 | | | <u> </u> | L | | 1 |
| map 4q35 repeat region | 1 | AF064849 | | <u> </u> | <u></u> | | <u> </u> | L | | |
| MAP kinase-interacting | 2 | AB000409 | | + | + | + | + | + | | 1 |
| serine/threonine kinase 1 | | | | | | 1 | | <u> </u> | | |
| (MKNK1) | | 1170070 | | + | <u> </u> | ₩ | | | | 4 |
| MAP/ERK kinase kinase 3 | 5 | U78876 | | * | | | | 1 | | - |
| (MEKK3) MAP/ERK kinase kinase 5 | 1 | D84476 | | + | + | ╁ | + | | | 4 |
| (MEKK5) | ' | 204410 | | 1 | ` | 1 | l . | | | 1 |
| W | | 1 | | | | | | | · | _ |

| | | | | | | | | | | |
|--|---|----------|---------------------------------------|-----|---|----------|---|---|-------|---|
| MAP/microtubule affinity- regulating kinase 3 (MARK3) | 4 | M80359 | | + | + | | | + | | |
| Marenostrin protein | | Y14441 | | 1 | | | | | • | |
| MASL1 | 1 | AB016816 | | ╁─┤ | | <u> </u> | | | - | |
| MAX dimerization protein | 3 | L06895 | | 1 | | | | + | | |
| (MAD) MaxiK potassium channel | 1 | AF035046 | | - | | | | | | _ |
| beta subunit | · | | | | | | | | | |
| MBP-2 for MHC binding protein 2 | 1 | X65644 | | + | + | + | | + | | |
| Meis (mouse) homolog 3 (MEIS3) | 1 | U68385 | _ | + | + | + | | + | | |
| melanoma-associated antigen p97 (melanotransferrin) | 1 | M12154 | | | | | | | | |
| membrane cofactor protein (CD46, trophoblast- lymphocyte cross-reactive antigen) (MCP) | 4 | X59405 | | + | + | + | | + | | |
| membrane component, chromosome 17, surface marker 2 (ovarian carcinoma antigen CA125) (M17S2) | 4 | D14696 | | + | + | + | + | + | | · |
| membrane metallo- endopeptidase (neutral endopeptidase, enkephalinase, CALLA, CD10) (MME) | 2 | J03779 | В | | + | + | + | + | | |
| membrane protein, palmitoylated 1 (55kD) (MPP1) | 2 | M64925 | | + | + | + | + | + | | |
| meningioma expressed antigen (MGEA) | 1 | U94780 | | | | + | | | | |
| meningioma-expressed antigen 11 (MEA11) | 1 | U73682 | + | + | | + | + | | | |
| Menkes Disease (ATP7A) putative Cu++-transporting P-type ATPase | 1 | L06133 | | + | | | | | | |
| metallothionein 2A (MT2A) | 1 | V00594 | | + | + | + | + | + | | |
| metaxin 1 (MTX1) | 1 | U46920 | | + | | + | | + | | |
| methionine | 2 | X68836 | + | + | + | + | | + | | |
| adenosyltransferase II, alpha (MAT2A) | | | | | | | | | | |
| methyl-CpG binding domain protein 1 (MBD1) (non-exact 59%aa) | 1 | Y10746 | | | | | | | | |
| methylene tetrahydrofolate dehydrogenase (NAD+ dependent), methenyltetrahydrofolate cyclohydrolase (MTHFD2) | 2 | X16396 | + | + | + | + | | + | | |
| methylenetetrahydrofolate dehydrogenase (NADP+ dependent), methenyltetrahydrofolate cyclohydrolase, formyltetrahydrofolate synthetase (MTHFD1) | 1 | J04031 | | + | + | + | + | + | | |
| methyltransferase, putative | 2 | AJ224442 | | | | | | | | |
| MHC antigen (HLA-B) (=L42024) | 1 | U14943 | | | | | | | | |
| MHC class 1 region | 2 | AF055066 | · _ , | | | | | | | |
| MHC class I antigen (HLA-A2) | 1 | U70863 | · · · · · · · · · · · · · · · · · · · | | | | | | | |
| MHC class I antigen (HLA-A33) | 1 | U19736 | | | | | | | | |
| MHC class I antigen (HLA-C) | 1 | U38975 | | | | | | | | |

| | | | | | | | | | _ |
|--|-------------|----------|---|--|---------------|---|---|--|---|
| MHC class I antigen B*5801 (HLA-B) | 1 | U52813 | | | | | | | |
| MHC class I antigen HLA-A (HLA-A) | 2 | AF015930 | | | | | | | |
| MHC class I antigen HLA-A | | U36687 | | ┝╌┤ | | - | _ | | l |
| (HLA-A-2402 allele) | | | | | | | | | |
| MHC class I antigen HLA- A11K | 2 | X13112 | | | | | | | _ |
| MHC class I antigen HLA-B | 1 | U67331 | | | | | | | |
| (8*0801 variant) (=AF028596) | | | | | | | | | |
| MHC class I antigen HLA-B (8*0801 variant) (=U88254) | 1 | U67330 | | | | | | | |
| MHC class I antigen HLA-B (B*48 allele) | 1 | AF017328 | | | | | | | |
| MHC class I antigen HLA-B (HLA-B*1502 allele) | 1 | AF014770 | | | | | | | |
| MHC class I antigen HLA-B (HLA-B*40MD) | 1 | U58643 | | | | | | | |
| MHC class I antigen HLA-B (HLA-B*4103 allele) | 1 | AF028596 | - | | | | | | |
| MHC class I antigen HLA-B | 1 | AF035648 | | | | | - | | |
| gene (HLA-B*4402 variant allele) | | | | | | | | | |
| MHC class I antigen HLA-B GN00110-B*3910 | 1 | U52175 | | | | | | | |
| MHC class I antigen HLA- Cw*04011 | 1 | D83030 | | | | | | | |
| MHC class I antigen R69772 HLA-A (A*0302) | 1 | U56434 | | | | | | | |
| MHC class I antigen | 1 | U58469 | | | | | • | | |
| SHCHA (HLA-B*4403 variant) | | | | | | | | | |
| MHC class I | 1 | U06697 | | | | | | | |
| histocompatibility antigen (HLA-B) (clone C21/14) | |] | | | | | | ŀ | |
| MHC class THLA B71 | 2 | L07950 | | | | _ | | \vdash | |
| MHC class I HLA-A (Aw33.1) | 1 | Flp | | | | _ | | \vdash | |
| MHC class I HLA-B | 1 | U18660 | | - | | | _ | - | |
| MHC class I HLA-B (HLA- | 1 | U18661 | | | | | | | |
| B-07ZEL allele) (=X86704) | • | | | | | | | | |
| MHC class I HLA-B (HLA- B-08NR allele) | 1 | U28759 | | | | | | | |
| MHC class I HLA-B*3512 | 1 | L76094 | | | | | | | |
| MHC class I HLA-B41 variant (=U17572) | 3 | U17572 | | | | | | | |
| MHC class I HLA-B44.2 | 1 | M24038 | | | | | | | |
| chain MHC class I HLA-B51- | | L41086 | | | | | | <u> </u> | |
| cd3.3 | | | | <u> </u> | | | | | |
| MHC class I HLA-C allele | 2 | Z33459 | | | | | | | |
| MHC class I HLA-Cw*0304 (=M84172; M99389) | 1 | D64150 | | | | | | | |
| MHC class I HLA-Cw*0803 | 3 | Z15144 | | | | | | | |
| MHC class I HLA-Cw6 | 1 | M28206 | | | | | | | |
| MHC class I HLA-J antigen | 1 | L56139 | | | | | | | |
| MHC class I lymphocyte antigen A2 (A2.1) variant DK1 | 1 | M19670 | | | | | | | |
| MHC class I mic-B antigen | 1 | X91625 | | | | | | | |
| MHC class I polypeptide- | 1 | L14848 | | | | + | · | | |
| related sequence A (MICA) | 1 | U61274 | | | | | · | | |
| heavy chain (C*0701new allele) (=AF017331) | • | 001214 | | | | | | | |
| MHC class II DNA | 1 | L18885 | | | | - | - | | |
| Sequence (clone A37G7-1C11) | | | | | _ | | | | |
| | | • | | | | | | | |

| WO 00/40749 | | | | | | | | | CI/CA00/00003 |
|--|----|-----------|---|--|--|--------------|----------|----------|---------------|
| MHC class II DQ-alpha associated with DRw6, | 1 | M16995 | + | | + | + | | + | |
| DQw1 protein | | | _ | | | | | | |
| MHC class II DQ-beta | 2 | M17564 | | + | | + | | + | |
| associated with DR2, DQw1 protein | | | | | | | | | |
| MHC class II HAL-DQ- | 1 | M33842 | | | | | | | |
| LTR5 (DQ,w8) DNA fragment, long terminal | | | | 1 | | | | | |
| repeat region | | | | | <u> </u> | | | | |
| MHC class II hla-dr alpha- | 1 | J00195 | | | | | | | |
| chain (=J00197;M60334;K01117 | | | | | | | | | |
| 1;J00194;M60333;X00274) | | | | | | | | | |
| MHC class II HLA-DRB1 | 1 | AF007883 | | <u> </u> | | L | | | |
| MHC class II HLA-DRw11- beta-I chain (DRw11.3) | 1 | M21966 | | | | | | | |
| MHC class II lymphocyte | 1 | M23907 | | | | | | | |
| antigen (DPw4-beta-1) | 1 | P33076 | | ₩ | | | | | |
| MHC CLASS II TRANSACTIVATOR CIITA | ' | P33076 | | | | | | | |
| (non-exact 57%) | | | | ļ | | | | | |
| MHC HLA-E2.1 (=X87679) | 1 | M32507 | | | ļ | | | | |
| MHC HLA-E2.1 (alpha-2 domain) (low match) | 1 | M32507 | | 1 | | | | | |
| Mi-2 autoantigen 240 kDa | 1 | U08379 | | | | | | | |
| protein (non-exact 84%) microsomal stress 70 | 7 | U04735 | | - | ļ | _ | <u> </u> | | |
| protein ATPase core (stch) | | 004735 | | |] | • | | | |
| microtubule-associated protein 4 (MAP4) | 1 | U19727 | + | + | + | + | | + | |
| microtubule-associated | 1 | X73882 | | + | | <u> </u> | | | |
| protein 7 (MAP7) | 2 | M16801 | | + | <u> </u> | + | | + | |
| mineralocorticoid receptor (aldosterone receptor) | 2 | W 10801 | | * | | 1 | | • | |
| (MLR) | | | | <u> </u> | | | | | |
| minichromosome maintenance deficient (S. | 1 | X62153 | İ | + | + | + | | + | |
| cerevisiae) 3 (MCM31) | | | | | <u> </u> | | | <u> </u> | |
| minichromosome | 1 | AB011144 | | + | + | + | | + | |
| maintenance deficient (S. cerevisiae) 3-associated | | | | | | | ŀ | | |
| protein (MCM3AP) | | V-7.1-70F | ļ | | <u> </u> | | + | + | |
| minichromosome maintenance deficient (S. | 2 | X74795 | + | + | + | + | 1 | * | |
| cerevisiae) 5 (cell division | | | | | | | | | |
| cycle 46) (MCM5) mitochondiral cytochrome b | 1 | AF042517 | | ┼ | <u> </u> | | | | |
| (CYTB) | • | 7, 0,2517 | | 1 | | | | | |
| mitochondrial 16S rRNA | 11 | Z70759 | | | | | | | |
| mitochondrial ATP | 2 | X59066 | | | | | | | |
| synthase (F1-ATPase) | | i | | | | | | | |
| mitochondrial ATP | 1 | X69907 | | | | | | | |
| synthase c subunit (P1 form) | | | | | | | | | |
| mitochondnal cytochrome b | 6 | AF042508 | | | | | | | |
| (CYTB) mitochondrial cytochrome b | | AB006202 | | + | | _ | | | |
| small subunit of complex II | · | | | | <u> </u> | | | | |
| mitochondrial CYTOCHROME C | 1 | P00395 | | | | | | | |
| OXIDASE POLYPEPTIDE I | | 1 | | | | • | | | |
| mitochondrial | 1 | P00403 | | Ī | | | | | |
| CYTOCHROME C OXIDASE POLYPEPTIDE | | 1 | | | | | | | |
| 11 | | | | <u> </u> | | | | | |
| mitochondrial cytochrome C oxidase subunit II | 2 | P00403 | | | 1 | | | | |
| C oxinase annuilt ii | | I | L | | Ь— | | | Щ. | <u> </u> |

| WU 00/40/49 | | | | | | | | | | _ |
|---|-----|----------|--|--|----------------|--------------|--|-------------|--------------|----------|
| mitochondrial cytochrome | 5 | U12691 | | | | | | | | |
| oxidase subunit II (COII) | | | | 1 1 | | i | l | | | ١ |
| (=U12692 Hsa4 | | | | | | ļ | i | | | ı |
| mitochondrion cytochrome | | | | ۱ ۱ | | | | | | |
| oxidase subunit II) | | X89763 | | - | | | | | | ᅱ |
| mitochondrial DNA loop | 1 | V03/02 | | | | | - 1 | | | -1 |
| attachment sequences (clone LAS34) | | | | | | | İ | | | |
| mitochondrial DNA | 1 | U94703 | | + | | | | $\neg \neg$ | | ٦ |
| polymerase accessory | • | 001.00 | | 1 | | | | l | | |
| subunit precursor (MtPolB) | | | | 1 1 | | | | | | |
| nuclear gene encoding | | | | | | 1 | | 1 | | |
| mitochondrial protein, | | | | | | | | | | 4 |
| mitochondrial DNA, | 1 | X93334 | | | | | | . 1 | | - |
| complete genome | | | | | | | | | | \dashv |
| mitochondrial genes for | 8 | V00710 | | 1 | | | | | | |
| several tRNAs (Phe, Val, | | 1 | | ļ | | | | | | |
| Leu) and 12S and 16S | | Ī | | 1 1 | 1 | | | . | | |
| ribosomal RNAs. | 3 | V00660 | | | - | | | | | ┥. |
| mitochondrial genes for tRNA (Phe) and 12S rRNA | | "00000 | | | | | | | | |
| (fragment) | | | | | | | | | | _ |
| mitochondrial inner | 1 | AF106622 | | | | | | | | ٦ |
| membrane preprotein | | | l | | | | | | | |
| translocase Tim17a | L | L | L | | L | | | | | _ |
| mitochondrial isolate Afr7 | 1 | AF042503 | | | | | | | | |
| cytochrome b(CYTB) | | Vasaria | | <u> </u> | | <u> </u> | | | | \dashv |
| mitochondrial loop | 1 | X89843 | | | 1 | | | | | |
| attachment sequence | | 1 | | | | | | | | - |
| (clone LAS88) | 14 | AF014893 | | + | | | | \vdash | | \dashv |
| mitochondrial NADH dehydrogenase subunit 2 | 14 | AFU14093 | | | | | | | | |
| | ĺ | 1 | | | | | | | | - |
| (ND2) mitochondrial translational | | L34600 | | + | + | + | | + | | コ |
| initiation factor 2 (MTIF2) | | | | | 1 | l | | | | |
| mitochondrion cytochrome | 1 | U09500 | | 1 | | | | | | コ |
| b | , | | <u>i</u> | <u> </u> | <u></u> | | | L | | |
| mitogen inducible gene | 1 | Z24725 | | + | + | + | | + | | - [|
| mig-Ž | ļ | | | <u> </u> | | | <u> </u> | | | 4 |
| mitogen inducible gene | 1 | Z24725 | | | | | ĺ | | | |
| mig-Ž (non-exact, 71%) | | <u> </u> | | | ↓ | ⊢ | | | | 4 |
| mitogen-activated protein | 2 | U43784 | 1 | + | + | + | | + | | - |
| kinase-activated protein | | | ļ | ı | ľ | | | | | - 1 |
| kinase 3 (MAPKAPK3) | | X80199 | | + | + | + | + | + | | ㅓ |
| MLN51 | 2 | | | 1 | L | | | L. | | 4 |
| MLN64 (=D38255 CAB1) | 1 | X80198 | + | + | + | + | | l | | ╛ |
| moesin (MSN) | 14 | M69066 | + | + | + | + | | + | | \neg |
| monocytic leukaemia zinc | 2 | U47742 | | + | + | + | | + | | ᅱ |
| finger protein (MOZ) | 1 | 04/142 | 1 | 1 | 1 | | | | | |
| MOP1 () | 2 | U29165 | | | † | | | | | \neg |
| 1 4 | 2 | D21094 | | + | + | + | \vdash | + | | \dashv |
| motor protein (Hs.78504) | | | <u> </u> | <u> </u> | <u> </u> | | - | ١ | | \dashv |
| mouse double minute 2, | 1 | U39736 | 1 | | + | + | | | | |
| human homolog of; p53- | 1 | 1 | 1 | 1 | 1 | 1 | 1 | | | |
| binding protein (MDM2) | 1 | X98263 | | + | - | + | | + | | \dashv |
| M-phase phosphoprotein 6 | i ' | Va0502 | 1 | 1 | ' | ` | | ` | | |
| (MPP-6) M-phase phosphoprotein, | 1 | X98260 | | + | +- | | | | | ヿ |
| m-phase phosphoprotein, | 1 ' | 7.00200 | | 1 | | 1 | | | | [|
| MPS1 | 1 1 | L20314 | | 1 | † | | - | | | ヿ |
| | 2 | D64154 | | + | | + | + | + | <u> </u> | ᅱ |
| Mr 110,000 antigen | | 1 | | + | ↓ | Ļ | <u> </u> | <u> </u> | | -1 |
| MRC OX-2, V-like region | 1 | X05324 | 1 | 1 | | | 1 | | | - |
| (=M17227) | | V/00000- | ļ | ↓ | | ├ | | ├ | | \dashv |
| mu-adaptin-related protein- | 1 | Y08387 | | 1 | | | 1 | | | 1 |
| 2; mu subunit of AP-4 (MU- | } | | | 1 | | 1 | |] | | |
| ARP2) | 1 | X53793 | + | + | + | + | ╁ | + | | ᅥ |
| multifunctional polypeptide similar to SAICAR | ' | 703/93 | , | ' | ' | Ι΄. | | 1 | | |
| synthetase and AIR | 1 | | | | | | 1 | ł | 1 | - 1 |
| carboxylase (ADE2H1) | 1 | [| | | | 1 | ł | <u></u> | | ╝ |
| SELECTION (NOTELLITY) | 1 | | | | | | | | | _ |

| WO 00/40/49 | | | | | | | | | | |
|---|----|----------|---|---|---|---|---|---|------|---|
| murine leukemia viral (bmi- 1) oncogene homolog (BMI1) | 1 | L13689 | | 1 | | + | | + | | |
| mutant (Daudi) beta2 - microglobulin | 44 | X07621 | | | | | | | | |
| mutated in colorectal cancers (MCC) | 1 | M62397 | | + | + | | | + | | |
| myeloid cell leukemia sequence 1 (BCL2-related) (MCL1) | 9 | L08246 | + | + | + | + | + | - | | |
| myeloid cell nuclear differentiation antigeN (MNDA) | 11 | M81750 | + | | | | | + | | |
| myeloid differentiation primary response gene (88) (MYD88) | 4 | U70451 | | + | + | + | | + | | |
| myeloid leukemia factor 2 (MLF2) | 3 | U57342 | | + | | + | | + | | |
| myeloid/lymphoid or mixed- lineage leukemia (trithorax (Drosophila) homolog); translocated to, 7 (MLLT7) | 8 | U89867 | | + | + | + | | + | | |
| MYH9 (cellular myosin heavy chain) | 1 | M81105 | | | | | | | - | |
| myomesin (M-protein) 2 (165kD) (MYOM2) | 1 | X69089 | | | | | | | | |
| myosin IE (MYO1E) | 11 | X98411 | | + | _ | + | | | | |
| myosin light chain kinase (MLCK) | 1 | U48959 | + | | + | + | | + | | |
| myosin phosphatase, target subunit 1 (MYPT1) | 2 | D87930 | | + | + | + | | + | | |
| myosin regulatory light chain (=U26162) | 2 | D50372 | | | | | | | | |
| myosin VIIa (low match 71) | 1 | U55208 | | | | | | | ···· | |
| myosin, heavy polypeptide 9, non-muscle (MYH9) | 3 | M81105 | + | + | + | + | | + | | |
| myosin, light polypeptide, regulatory, non-sarcomeric (20kD) (MLCB) | 6 | X54304 | + | + | + | + | + | + | | : |
| myosin-l beta | 1 | X98507 | + | + | + | + | | + | | |
| myristoylated alanine-rich protein kinase C substrate (MARCKS, 80K-L) (MACS) | 1 | D10522 | | + | + | | | | | |
| myxovirus (influenza) resistance 1, homolog of murine (interferon-inducible protein p78) (MX1) | 1 | M30817 | + | + | + | + | | + | | |
| myxovirus (influenza) resistance 2, homolog of murine (MX2) | 3 | M30818 | | | + | | | | | |
| N-acetylgalactosaminidase, alpha- (NAGA) | 2 | M62783 | | + | + | | + | + | | |
| N-acetylglucosamine receptor 1 (thyroid) (NAGR1) | 1 | L03532 | | + | + | + | | + | | |
| NACP/alpha-synuclein | 2 | U46896 | | | | | | | | |
| N-acylaminoacyl-peptide hydrolase (APEH) | 1 | D38441 | | + | + | | + | + | | |
| N-acylsphingosine amidohydrolase (acid ceramidase) (ASAH) | 11 | U47674 | + | + | + | + | | + | | |
| NAD+-specific isocitrate dehydrogenase beta subunit precursor (encoding mitochondrial protein) | 1 | U49283 | + | + | + | + | + | + | | |
| NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 5 (13kD, B13) (NDUFA5) | 1 | U53468.1 | + | + | + | + | + | + | | |

| WO 00/40749 | | | | | | | | • | C1/CA00/00003 |
|---|--------------|-----------|--|--|--------------|--------------|--------------|--|----------------|
| NADH dehydrogenase | 1 | AF047181 | | + | + | + | + | + | |
| (ubiquinone) 1 beta subcomplex, 5 (16kD, | | 1 | | | | | | | |
| SGDH) (NDUFB5) | | _i | | | | | | | |
| NADH dehydrogenase | 1 | AF050640 | | + | + | + | + | + | |
| (ubiquinone) Fe-S protein 2 (49kD) (NADH-coenzyme | | | | | | | | | |
| Q eductase) (NDUFS2) | | | | l | | | | | |
| NADH dehydrogenase | 1 | M22538 | | | + | + | + | + | |
| (ubiquinoné) flavoprotein 2 | | | | 1 | | 1 | | | |
| (24kD) (NDÚFV2) NADH:ubiquinone | 2 | AF053070 | | + | + | + | + | + | |
| dehydrogenase 51 kDa | _ | | | | | | | | |
| subunit (NDUFV1) | | | | | | | | | |
| NADH-CYTOCHROME B5 REDUCTASE (B5R) | 1 | P00387 | | 1 | | | | İ | |
| (50%aa) | | | | | | | | l | |
| NADH-UBIQUINONE | 1 | P03886 | | | | | | | |
| OXIDOREDUCTASE ICHAIN 1 | | } | | | | | | | [|
| Nardilysin (N-arginine | 2 | U64898 | + | + | + | + | _ | + | |
| dibasic convertase) | _ | | | | | 1 | | | |
| (NRD1) | | X80909 | | +- | + | | + | + | |
| nascent-polypeptide- associated complex alpha | 5 | Vonana | | 1 | | 1 | | ` | |
| polypeptide (NACA) | İ | | | | | l | | | |
| natural killer cell group 7 | - 8 | 569115 | | T | | + | | + | |
| sequence (NKG7) natural killer cell transcript | 19 | M32011 | + | | | \vdash | <u> </u> | | |
| 4 (NK4) | '5 | | | | | <u> </u> | | | |
| natural killer-associated | 1 | U30274 | + | | | | | | blood only |
| transcript 3 (NKAT3) natural killer-associated | | AF022045 | + | + | ├ | | | - | blood only |
| transcript 5 (NKAT5) | · ' | 74 OZZOVO | | | | | | | |
| natural killer-tumor | 1 | L04288 | В | | + | | + | + | |
| recognition sequence (NKTR) | | | | | İ | | | | |
| N-deacetylase/N- | 2 | AF042084 | + | + | | + | | + | |
| sulfotransferase (heparan | | | | | | | | | |
| glucosaminyl) 2 (NDST2) | 3 | Z35102 | | + | - | ╁ | | - | |
| Ndr protein kinase | 1 - | U96113 | | | | ╁ | | ├ | |
| Nedd-4-like ubiquitin- protein ligase WWP1 | ' | 090113 | <u> </u> | | | | | | |
| nel (chicken)-like 2 | 3 | D83018 | | + | + | 1 | | | |
| (NELL2) | | 1100440 | | ++ | | | + | ļ | |
| N-ethylmaleimide-sensitive factor attachment protein. | 1 | U39412 | | • | 1 | 1 | • | | |
| alpha (NAPA) | | | j | | | | | | |
| N-ethylmaleimide-sensitive | 1 | U78107 | | + | + | + | | | |
| factor attachment protein, | i | | | | | |] | | |
| gamma (NAPG) neural precursor cell | 3 | X92544 | + | ++ | + | + | - | + | high in testis |
| expressed, | | | | | İ | | 1 | į . | |
| developmentally down- | • | | | | | | | ı | |
| regulated 5 (NEDD5) neural precursor cell | 1 | D23662 | | + | + | + | + | + | |
| expressed. | , | | | | | | | | |
| developmentally down- | | | | | | 1 | | | |
| regulated 8 (NEDD8) neuregulin 1 (NRG1) | 1 1 | U02330 | | + | + | + | + | + | |
| neuroblastoma RAS viral | 4- | AB020692 | + | + | + | + | | + | |
| (v-ras) oncogene homolog | _ | | | | | | | | |
| (NRAS) | <u> </u> | Venene | L | | | <u> </u> | | | |
| Neuroblastoma RAS viral (v-ras) oncogene homolog | 1 | X68286 | | | | | Ι΄ | | |
| (NRAS) (low match) | | | 1 | | L | | Ŀ | L | |
| Neurofibromin 2 (bilateral | 1 | S73853 | Ī | + | | | | + | |
| acoustic neuroma) (NF2) | 2 | U19251 | + | ++ | ╀ | + | \vdash | + | |
| neuronal apoptosis inhibitory protein (NAIP) | 2 | 019231 | ļ , | ' | | \perp | L | Ĺ | |
| neuronal cell adhesion | 1 | AB002341 | | + | + | + | | + | |
| molecule (NRCAM) | <u> </u> | J | <u> </u> | 1 | 1 | <u> </u> | <u> </u> | Ц | <u> </u> |

| neuropathy target esterase (NTE) | 1 | AJ004832 | | + | | + | | + | |
|---|-----------|------------|---------------|---------------|--------------|--------------|--------------|--|--|
| neuropeptide Y3 receptor, 5'UTR (low score) | 1 | D28433 | | | | | | | |
| neurotrophic tyrosine kinase, receptor, type 1 (NTRK1) | 14 | X03541 | + | + | + | + | + | + | |
| neutrophil cytosolic factor 4 (40kD) | 2 | U50720 | | | | | | | |
| NG31 | 1 | AF129756 | | | | | | | |
| NGAL (=X83006) | 1 | X99133 | | | | | | | |
| nibrin (NBS) | 1 | AF051334 | | | | | | | |
| NIK | 1 | AB014587 | | + | + | + | | + | |
| Ninjurin 1; nerve injury- | 1 | U72661 | | + | + | + | | + | |
| induced protein-1 nitrilase 1 (NIT1) | | AF069987 | | | | | | | |
| (=AF069984) | | | | 1 | | | | | |
| NKG2-D (low match) (non- exact, 58%) | 1 | X54870 | | | | | | | |
| Nmi | 1 | U32849 | | | | | L | | |
| N-myristoyltransferase 1 (NMT1) | 1 | AF043324 | - | + | + | + | + | + | |
| No arches-like (zebrafish) zinc finger protein (NAR) | 1 | U79569 | | + | + | + | | + | |
| non-histone chromosome protein 2 (S. cerevisiae)- like 1 (NHP2L1) | 1 | D50420 | + | + | + | + | + | + | |
| non-muscle (fibroblast) tropomyosin | 1 | | | | | | | | |
| non-muscle alpha-actinin | | U48734 | | | | | | | |
| non-muscle myosin alkali light chain (Hs.77385) | 3 | M22918 | + | + | + | + | + | + | High in fetal adrenal gland and BPH stroma |
| non-neuronal enolase (EC 4.2.1.11) | 1 | X16289 | | | | | | | |
| non-receptor tyrosine phosphatase 1 | 1 | M33689 | | | | | | | |
| normal keratinocyte substraction library mRNA, iclone H22a | 3 | X53778 | + | + | + | + | + | + | high in many libraries |
| notch group protein (N) | 3 | M99437 | ····· | | | | | | |
| novel protein | 1 | X99961 | | | <u> </u> | | _ | - | |
| novel T-cell activation | <u> </u> | X94232 | | + | + | + | | + | |
| protein | · · · · · | A60196 | | - | | <u> </u> | <u> </u> | | |
| N-ras protein NRU | | U60111 | | + | <u> </u> | ļ | | + | |
| N-sulfoglucosamine sulfohydrolase (sulfamidase) (SGSH) | 1 | 000111 | | | | | | ľ | |
| Insulin induced gene 1 (INSIG1) | 1 | U96876 | + | + | + | + | + | + | |
| ntegrin, alpha 4 (antigen CD49D, alpha 4 subunit of | 3 | L12002 | + | | | + | _ | | |
| VLA-4 receptor) (ITGA14) Interferon, gamma-inducible | 1 | M63838 | + | + | + | + | | + | |
| protein 16 (IFI16) Interleukin 1, beta (IL1RB) | 1 | M15330 | | | | 1 | \vdash | - | |
| nuclear antigen H731-like | 2 | U83908 | | + | + | + | | + | |
| protein nuclear antigen Sp100 | 4 | U36501 | + | | | + | + | + | |
| (SP100) Nuclear antigen Sp100 | 1 | P23497 | | | | ┼- | - | - | |
| (SP100) (85%aa) Nuclear antigen Sp100 | 1 | P23497 | | + | ├ | - | | - | ļ |
| (SP100) (89%aa) | <u>'</u> | M97856 | + | | - | <u> </u> | <u> </u> | <u> </u> | |
| nuclear autoantigenic sperm protein (histone- binding) (NASP) | | 1419 / 650 | ···· | | | | | L | |

| WO 00/40/49 | | | | | | | | | |
|---|-----|----------|---|---|---|---|---|----------|---|
| nuclear corepressor KAP-1 (KAP-1) (=U95040; X97548 TIF1beta zinc finger | 1 | U78773 | | | | | | | |
| protein) | 4 | U22897 | + | + | + | + | + | + | |
| Nuclear domain 10 protein (NDP52) | 4 | 022097 | • | * | • | 1 | | | _ |
| Nuclear factor (erythroid- derived 2)-like 2 (NFE2L2) | 1 . | S74017 | | + | + | + | + | + | |
| Nuclear factor of kappa light polypeptide gene enhancer in B-cells 1 | 2 | M58603 | | + | + | | + | + | - |
| (p105) (NFKB1) | | | | | | | | <u> </u> | |
| nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, alpha (NFKBIA) | 3 | M69043 | | + | + | + | | + | |
| nuclear factor related to kappa B binding protein (NFRKB) | 1 | U08191 | | + | + | + | | + | |
| nuclear mitotic apparatus protein 1 (NUMA1) | 3 | Z11583 | + | + | + | + | + | + | |
| nuclear receptor coactivator 2 (GRIP1) | 1 | X97674 | | | | | | | |
| nuclear receptor coactivator 3 (AIB3) | 2 | AF010227 | + | + | + | | | + | |
| nuclear receptor coactivator 4 (ELE1) | 22 | X77548 | | + | + | + | + | + | |
| nuclear receptor interacting protein 1 (NRIP1) | f | X84373 | | + | | + | | + | |
| nuclear respiratory factor 1 (NRF1) | 1 | U02683 | В | + | + | | | | _ |
| nuclear RNA helicase, DECD variant of DEAD box family (DDXL) | 4 | U90426 | + | + | + | + | | + | |
| nuclear transcription factor Y, alpha (NFYA) | 1 | X59711 | В | | | | | | |
| nuclear transcription factor, X-box binding 1 (NFX1) | 3 | U15306 | | + | + | | + | | |
| nuclear transport factor 2 (placental protein 15) (PP15) | 1 | X07315 | + | + | + | + | | + | |
| nucleobindin (=M96824) | 1 | U31336 | | | | | | | |
| nucleobindin 1 (NUCB1) | 2 | M96824 | + | + | + | + | | + | |
| nucleolar phosphoprotein p130 (P130) | 1 | Z34289 | | + | + | | | | |
| nucleolar protein (KKE/D repeat) (NOP56) | 1 | Y12065 | + | + | + | + | | + | |
| nucleolar protein (MSP58) | 1 | AF015308 | | | | | | | |
| nucleolar protein 1 (120kD) (NOL1) | 1 | M32110 | + | + | | | | | |
| nucleolar protein p40 | 1 | U86602 | + | + | + | + | | + | |
| nucleolin (NCL) | 2 | M60858 | + | + | + | + | | + | |
| nucleophosmin (nucleolar phosphoprotein B23, numatrin) (NPM1) | 14 | M28699 | + | + | + | + | | + | |
| nucleophosmin-retinoic acid receptor alpha fusion protein NPM-RAR long form | 1 | U41742 | | | | | | | |
| nucleoporin (NUP358) (=D42063 RanBP2 (Ran- binding protein 2)) | 2 | L41840 | | | | | | | |
| nucleoporin 153kD (NUP153) | 1 | Z25535 | | | | | | | |
| nucleoponn 98kD (NUP98) | 1 | U41815 | | | | | | | |
| nucleosome assembly protein | 1 | D28430 | | | | | | | |
| nucleosome assembly protein 1-like 1 (NAP1L1) | | M86667 | | + | + | + | | + | |
| nucleosome assembly protein 1-like 4 (NAP1L4) | 2 | U77456 | + | + | + | + | | + | |
| | | | | | | | | | |

| nucleosome assembly protein, 5'UTR | 1 | D28430 | | | | 7 | | | |
|---|---|--|---|--|-------------|-----------------|-----|-------|-----------------------|
| olfactory receptor (OR7- | 1 | U86281 | | | | | | | |
| 141) | | 024002 | | | | \dashv | | | |
| LIKE PROTEIN HGMP07E | 1 | P34982 | | | ļ | | | | |
| (OR17-4) (non-exact 65%) oligodendrocyte myelin | | L05367 | | + | | | - | | |
| glycoprotein (OMG) | , | 200007 | | 1 | | | | | |
| O-linked N- | 1 | U77413 | + | + | | + | + | + | |
| acetylglucosamine (GlcNAc) transferase | | | | | | | | | |
| acetylglucosamine:polypep tide-N-acetylglucosaminyl transferase) (OGT) | | | | | | | | | |
| oncofetal trophoblast glycoprotein 5T4 precursor (non-exact 55%) | 1 | A53531 | | | | | | | |
| Oncogene TIM (TIM) (non- exact 84%) | 1 | U02082 | | | | | | | |
| ORF (Hs.77868) | 1 | M68864 | + | + | + | + | + | + | |
| ORF1; MER37; putative | 1 | U49973 | | | | $\vdash \vdash$ | | | |
| transposase similar to pogo element Length = 454 | · | | | | | | | | |
| origin recognition complex, subunit 2 (yeast homolog)- like (ORC2L) | 2 | U27459 | | | | + | | | |
| origin recognition complex, subunit 4 (yeast homolog)- like (ORC4L) (low match) | 1 | AF022108 | | + | + | + | | | |
| ornithine aminotransferase (gyrate atrophy) (OAT) | 2 | M23204 | | | Ψ. | • | | | |
| ornithine decarboxylase (ODC) | 1 | M20372 | | | | | | | |
| ornithine decarboxylase | 11 | D78361 | + | 1 + 1 | + | + | + | + | High in pancreas, |
| antizyme, ORF 1 and ORF | | | | | | | | | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor | 2 | U07132 | + | + | + | + | | + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) | | | + | + | + | + | + | + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM- | 2 | U07132 | | | | | + | | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM-40) ovel centrosomal protein | 2 | U07132 AB002806 | | | | | + | | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM-40) | 6 | U07132 AB002806 D28381 | | + | + | + | + | + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM- 40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast tumor protein | 2 6 1 1 | U07132 AB002806 D28381 AB008515 L34839 | | + | + | + | | + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM- 40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast tumor protein oviductal glycoprotein 1, 120kD (OVGP1) | 2 6 1 1 | U07132 AB002806 D28381 AB008515 L34839 U09550 | | + | + | + + | + | + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM-40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast tumor protein oviductal glycoprotein 1, 120kD (OVGP1) oxidase (cytochrome c) | 2 6 1 1 | U07132 AB002806 D28381 AB008515 L34839 U09550 X80695 | | + | + | + | | + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM-40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast tumor protein oviductal glycoprotein 1, 120kD (OVGP1) oxidase (cytochrome c) assembly 1-like (OXAIL) oxoglutarate dehydrogenase (lipoamide) | 2 6 1 1 | U07132 AB002806 D28381 AB008515 L34839 U09550 | | + | + | + + | + | + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM-40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast tumor protein oviductal glycoprotein 1, 120kD (OVGP1) oxidase (cytochrome c) assembly 1-like (OXAIL) oxoglutarate dehydrogenase (lipoamide) (OGDH) oxysterol binding protein | 2 6 1 1 1 1 1 1 1 | U07132 AB002806 D28381 AB008515 L34839 U09550 X80695 | | + + | + + + + + | + + | + | + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM-40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast tumor protein oviductal glycoprotein 1, 120kD (OVGP1) oxidase (cytochrome c) assembly 1-like (OXAIL) oxoglutarate dehydrogenase (lipoamide) (OGDH) | 2 6 1 1 1 1 1 1 1 4 | U07132 AB002806 D28381 AB008515 L34839 U09550 X80695 D10523 | + | + + + | + + + + + | + + | + + | + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM-40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast tumor protein oviductal glycoprotein 1, 120kD (OVGP1) oxidase (cytochrome c) assembly 1-like (OXAIL) oxoglutarate dehydrogenase (lipoamide) (OGDH) oxysterol binding protein (OSBP) OZF | 2 6 1 1 1 1 1 1 1 4 | U07132 AB002806 D28381 AB008515 L34839 U09550 X80695 D10523 M86917 | + | + + + + + + | + + + + + + | + + + | + + | + + + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM-40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast tumor protein oviductal glycoprotein 1, 120kD (OVGP1) oxidase (cytochrome c) assembly 1-like (OXAIL) oxoglutarate dehydrogenase (lipoamide) (OGDH) oxysterol binding protein (OSBP) OZF OZF (non-exact zinc finger) p21/Cdc42/Rac1-activated kinase 1 (yeast Ste20- related) (PAK1) | 2 6 1 1 1 1 1 1 1 1 1 2 | U07132 AB002806 D28381 AB008515 L34839 U09550 X80695 D10523 M86917 X70394 X70394 U51120 | + | + + + + + + | + + + + + + | + + + | + + | + + + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM-40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast tumor protein oviductal glycoprotein 1, 120kD (OVGP1) oxidase (cytochrome c) assembly 1-like (OXAIL) oxoglutarate dehydrogenase (lipoamide) (OGDH) oxysterol binding protein (OSBP) OZF OZF (non-exact zinc finger) p21/Cdc42/Rac1-activated kinase 1 (yeast Ste20- | 2 6 1 1 1 1 4 1 1 1 2 | U07132 AB002806 D28381 AB008515 L34839 U09550 X80695 D10523 M86917 X70394 X70394 U51120 D63392 | + | + + + + + + | + + + + + + | + + + | + + | + + + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM-40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast tumor protein oviductal glycoprotein 1, 120kD (OVGP1) oxidase (cytochrome c) assembly 1-like (OXAIL) oxoglutarate dehydrogenase (lipoamide) (OGDH) oxysterol binding protein (OSBP) OZF (non-exact zinc tinger) p21/Cdc42/Rac1-activated kinase 1 (yeast Ste20- related) (PAK1) P35-related protein (= | 2 6 1 1 1 1 1 1 1 1 1 2 | U07132 AB002806 D28381 AB008515 L34839 U09550 X80695 D10523 M86917 X70394 X70394 U51120 | + | + + + + + + | + + + + + + | + + + | + + | + + + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM-40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast tumor protein oviductal glycoprotein 1, 120kD (OVGP1) oxidase (cytochrome c) assembly 1-like (OXAIL) oxoglutarate dehydrogenase (lipoamide) (OGDH) oxysterol binding protein (OSBP) OZF OZF (non-exact zinc finger) p21/Cdc42/Rac1-activated kinase 1 (yeast Ste20- related) (PAK1) P35-related protein (= S80990 ficolin) | 2 6 1 1 1 1 4 1 1 1 2 | U07132 AB002806 D28381 AB008515 L34839 U09550 X80695 D10523 M86917 X70394 X70394 U51120 D63392 | + | + + + + + + | + + + + + + | + + + | + + | + + + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM-40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast tumor protein oviductal glycoprotein 1, 120kD (OVGP1) oxidase (cytochrome c) assembly 1-like (OXAIL) oxoglutarate dehydrogenase (lipoamide) (OGDH) oxysterol binding protein (OSBP) OZF OZF (non-exact zinc finger) p21/Cdc42/Rac1-activated kinase 1 (yeast Ste20-related) (PAK1) P35-related protein (= S80990 ficolin) p40 p40phox (=U50720) | 2 6 1 1 1 1 4 1 1 2 | U07132 AB002806 D28381 AB008515 L34839 U09550 X80695 D10523 M86917 X70394 X70394 X70394 U51120 D63392 U93569 | + | + + + + + + | + + + + + + | + + + | + + | + + + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM-40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast tumor protein oviductal glycoprotein 1, 120kD (OVGP1) oxidase (cytochrome c) assembly 1-like (OXAIL) oxoglutarate dehydrogenase (lipoamide) (OGDH) oxysterol binding protein (OSBP) OZF OZF (non-exact zinc finger) p21/Cdc42/Rac1-activated kinase 1 (yeast Ste20-related) (PAK1) P35-related protein (= S80990 ficolin) p40 p40phox (=U50720) P47 LBC oncogene | 2 6 1 1 1 1 4 1 1 2 | U07132 AB002806 D28381 AB008515 L34839 U09550 X80695 D10523 M86917 X70394 X70394 U51120 D63392 U93569 X77094 | + | + + + + + + | + + + + + + | + + + | + + | + + + | and activated T cells |
| antizyme, ORF 1 and ORF 2 orphan receptor (Hs.100221) OS-9 precurosor osteonectin (=X82259 BM-40) ovel centrosomal protein RanBPM (RANBPM) over-expressed breast tumor protein oviductal glycoprotein 1, 120kD (OVGP1) oxidase (cytochrome c) assembly 1-like (OXAIL) oxoglutarate dehydrogenase (lipoamide) (OGDH) oxysterol binding protein (OSBP) OZF OZF (non-exact zinc finger) p21/Cdc42/Rac1-activated kinase 1 (yeast Ste20-related) (PAK1) P35-related protein (= \$80990 ficolin) p40 p40phox (=U50720) | 2 6 1 1 1 4 1 1 2 | U07132 AB002806 D28381 AB008515 L34839 U09550 X80695 D10523 M86917 X70394 X70394 U51120 D63392 U93569 X77094 U03634 | + | + + + + + + | + + + + + | + + + + | + + | + + + | and activated T cells |

| VVO 00/40/42 | | VERENZ | | , , | | | | | |
|--|------------------|-----------------|------------------|----------------|--|--|--|--|------------------------|
| p62 nucleoporin | | X58521 | | | | | | | |
| p63 mRNA for | 1 | X69910 | + | + | + | + | | + | |
| transmembrane protein PAC clone DJ0701016 | 1 | Q07108 | | + | | | | | |
| from 7g33-g36 (non-exact | ' | 207100 | | | | <u> </u> | | | |
| 54%) | | 1 | | | | | | | |
| palmitoyl-protein | 10 | U44772 | | + | + | + | | + | |
| thioesterase (ceroid- | | | | | | | | | |
| lipofuscinosis, neuronal 1, infantile: Haltia-Santavuori | | 1 | | | | | İ | | |
| disease) (PPT) | | | | | | | | | |
| papillary renal cell | 1 | X99720 | + | + | + | + | + | + | |
| carcinoma (translocation- | | 1 | | | | 1 | l | | |
| associated) (PRCC) PAR protein | 1 | AF115850 | | + | - | + | ┢ | <u> </u> | |
| partial EST (clone c-1gh04) | - i - | Z43627 | | | | - | ! | - | |
| , | | 1 | | + | | | | | |
| PAX3/forkhead transcription factor gene | 1 | U02368 | | | | 1 | 1 | ŀ | |
| Ifusion | | | | | | | | 1 | |
| paxillin (PXN) | 4 | D86862 | | + | + | + | | + | |
| PBK1 protein | 2 | AJ007398 | + | + | + | + | | + | |
| PBS-EST (nz92e01.s1 | 1 | AA732534 | | + | | | | | |
| NCI CGAP GCB1 clone | · | | | | | ł | | 1 | |
| IMAGE:1302936) (low | | | | | | | | l | ł |
| Score) | 1 | AJ224747 | + | + | - | + | | + | |
| PDZ domain protein (Drosophila inaD-like) | ' | 70227171 | • | | | 1 | 1 | l ['] | |
| (INALD) | | | | | | | | L | |
| PEBP2aC Runt domain | 1 | Z38108 | | | | | | | |
| encoding gene (=Z35728) | 1 | J04605 | | | | ├ | - | - | |
| peptidase D (PEPD) | · | | | + | + | + | + | + | high in many libraries |
| peptidylprolyl isomerase A (cyclophilin A) (PPIA) | 3 | Y00052 | | • | * | ľ | • | • | mgn ni many iivianes |
| peptidylprolyl isomerase D | 2 | L11667 | - T - | + | + | | + | + | |
| (cyclophilin D) (PPID) | | | | <u> </u> | | | | | |
| peptidylprolyl isomerase E | 1 | AF042386 | | + | + | | + | + | |
| (cyclophilin E) (PPIE) PERB11.1 (=U56942 MHC | 1 | U69630 | | + | | | ├ | | |
| class I chain-related protein | , | 009030 | | | | | | ļ | |
| (A) | | | | | | | | <u> </u> | |
| perforin 1 (preforming | 14 | M28393 | | | | | | | |
| protein) (PRF1) | | V96027 | | - | | - | ▙ | | ļ |
| peroxisomal acyl-CoA thioesterase (PTE1) | 2 | X86032 | | - | | 1 | | | : |
| Peroxisomal acyl- | 1 | X71440 | | + | + | + | + | + | |
| coenzyme A oxidase | | | | | 1 | | <u> </u> | | |
| peroxisomal farnesylated | 1 | X75535 | | + | + | + | + | + | , |
| protein (PXF) phorbol-12-myristate-13- | | D90070 | B, W | + | \vdash | + | + | | |
| acetate-induced protein | ' | 555070 | ٥, ٠٠ | | | 1 | 1 | 1 | 1 |
| (PMAIP1) | | | | | | L | <u> </u> | L_ | |
| phosphate carrier | 1 | X77337 | | |] | | | 1 | |
| (mitochondrial gene?) | 3 | X60036 | | + | + | + | \vdash | + | |
| Phosphate carrier, mitochondrial (PHC) | 3 | \ ______\ | • | 1 | | ' | | ` | |
| phosphate | 1 | L28957 | T | † | + | 1 | + | 1 | |
| cytidylyltransferase 1, | |] | | 1 | | | | | 1 |
| choline, alpha isoform | | | | | | | l | ŀ | |
| (PCYT1A) | 1 | Q92903 | | + | - | +- | ┼ | | |
| CYTIDYLYLTRANSFERAS | ' | 402000 | | | | | [| 1 | |
| E (CDP-DIGLYCERIDE) | | | | | <u> </u> | | | <u> </u> | |
| phosphatidylinositol 3- | 2 | U57843 | | | | | | 1 | 1 |
| kinase delta catalytic | | <u> </u> | | 1 | | | 1 | 1 | 1 |
| subunit phosphatidylinositol 4- | 3 | AB005910 | + | + | + | + | | + | |
| kinase, catalytic, beta | | | • | | | 1 | 1 | | |
| polypeptide (PIK4CB) | | 1 | | <u> </u> | L | <u> </u> | | _ | |
| phosphatidylinositol glycan, | 1 | L19783 | | + | + | + | + | + | |
| class H (PIGH) | | .1 | | | L | ــــــــــــــــــــــــــــــــــــــ | 1 | L | <u> </u> |

PCT/CA00/00005

WO 00/40749

| WO 00/40/49 | | | | | | | | | | |
|--|---------------|----------|--|--|--|----------|---|----------|-------------|--|
| phosphatidylinositol transfer protein (PI-TPbeta) | 2 | D30037 | | | • | | | | | |
| phosphatidylinositol transfer protein, | 2 | X98654 | B, T lymphoma | + | | | | | | |
| membrane-associated (PITPNM) | | | | | | | | | | |
| phosphatidylinositol transfer protein, | 7 | X98654 | | | | | | | | |
| membrane-associated (PITPNM) (non-exact 64%) phosphatidylinositol-4- | 1 | U14957 | | | + | _ | + | | | |
| phosphatidylinositor== phosphate 5-kinase, type , alpha (PIP5K2A) | • | 014337 | | | ŕ | | | | | |
| phosphatidylinositol-4- phosphate 5-kinase, type II, beta (PIP5K2B) | 1 | U85245 | | + | + | + | | + | | |
| phosphodiesterase 7A (PDE7A) | 1 | L12052 | B, W | + | + | | + | | | |
| phosphodiesterase IB (PDES1B) | 1 | U56976 | | ON | LY | | | | | |
| phosphoglucomutase 1 (PGM1) | 2 | M83088 | | + | + | + | | + | | |
| phosphogluconate dehydrogenase (PGD) | 1 | U30255 | | | + | | | | | |
| phosphoglycerate kinase 1 (PGK1) | 12 | V00572 | | | + | + | + | + | | |
| phosphoglycerate mutase 1 (brain) (PGAM1) | 3 | J04173 | + | + | + | | + | + | | |
| phosphoglycerate mutase 2 (muscle) (PGAM2) | 1 | M55673 | | | + | + | | T | | |
| phosphoinositide-3-kinase, catalytic, alpha polypeptide (PIK3CA) | 1 | Z29090 | | + | • | • | | | | |
| phosphoinositide-3-kinase, catalytic, delta polypeptide (PIK3CD) | 4 | U86453 | | + | + | + | | + | | |
| phosphoinositide-3-kinase, catalytic, gamma polypeptide (PIK3CG) | 1 | X83368 | | | | | | | | |
| phospholipase C | 1 | X14034 | | | | | | | | |
| phospholipase C, delta 1 (PLCD1) | 2 | U09117 | | + | + | + | | + | | |
| phospholipase C, gamma 1 (formerly subtype 148) (PLCG1) | 1 | M34667 | + . | + | + | + | | + | | |
| phospholipid scramblase | 1 | AF008445 | | | | | | | | |
| phosphoribosyl pyrophosphate synthetase- associated protein 1 (PRPSAP1) | 1 | D61391 | | + | + | | | + | | |
| phosphonbosylglycinamide formyltransferase, phosphoribosylglycinamide synthetase, phosphoribosylaminoimida zole synthetase (GART) | 3 | X54199 | | + | + | + | + | + | | |
| phosphorylase kinase, alpha 2 (liver), glycogen storage disease IX (PHKA2) | 3 | D38616 | | + | + | + | + | + | | |
| phosphorylase, glycogen; brain (PYGB) | 1 | U47025 | + | + | + | | | + | | |
| phosphorylase, glycogen; brain (PYGB) (low match, non-exact, 75%) | 1 | U47025 | | | | | | | | |
| phosphorylase, glycogen; liver (Hers disease, lycogen storage disease | 1 | Y15233 | | | + | + | | + | | |
| type VI) (PYGL) phosphorylation regulatory protein HP-10 | 2 | | | | | \vdash | | | | |
| phosphotidylinositol transfer protein (PITPN) | 1 | D30036 | + | + | + | + | | + | | |
| transfer protein (FITEN) | | | 75 | ــــــــــــــــــــــــــــــــــــــ | Ь | Ц_ | Ц | Щ. | | |

| factor (PEDF) | | | | | | | | | | |
|--|--|---------------|---------|----------|--------|-----|-------|-----------|-----|------|
| Direction Dire | | 1 | U29953 | + | | + | + | + | | |
| associated protein (PNN) placental (Diff33) placental (Diff33) placental (Diff33) placental (Diff48) placent | l' | | | + | | | | | | |
| Diacental (Chiff33) 100 | associated protein (PNN) | 1 | | | B, | mon | ocyte | e, fl | ymp | noma |
| Exact. 99% Diacenta (Diff48) 18 | placenta (Diff33) | | | | + | + | + | | + | |
| placenta (Diff48) 18 | placenta (Diff33) (non- exact, 69%) | 1 | U49188 | |] | | | | | |
| Tratich | placenta (Diff48) | | | + | | | | | | |
| Diacental(Diff48) (low match) Diasminogen activator, unokinase receptor (PLAUR) Diasminogen activator, unokinase receptor (PLAUR) Diatelet factor 4 (PF4) 1 M25897 + | | - | U49187 | | | | | | | |
| Diasminogen activator, | placenta(Diff48) (low | 1 | U49187 | | | | | | | |
| Diatelet/endothetial cell adhesion molecule (CD31 intigen) (PECAM1) Diatelet-activating factor acetylhydrolase 2 (40kD) (PAFAH2) Diatelet-activating factor acetylhydrolase 2 (40kD) (PAFAH2) Diatelet-activating factor acetylhydrolase, isoform ib, alpha subunit (45kD) (PAFAH181) Diatelet-activating factor acetylhydrolase, isoform ib, alpha subunit (45kD) (PAFAH181) Diatelet-activating factor receptor (PTAFR) Diatelet-activating factor receptor (PTAFR) Diatelet-activating factor receptor (PTAFR) Dieckstrin (PLEK) (10w Match) M85169 + + + + + + + + + + + + + + + + + + | urokinase receptor | 1 | X74039 | | + | | + | | + | |
| Adhesion molecule (CD31 | platelet factor 4 (PF4) | | 1 | | | | | | ŀ | |
| acetylhydrolase 2 (40kD) (PAFAH2) | adhesion molecule (CD31 Intigen) (PECAM1) | | | | | | | + | + | |
| Scetylhydrolase, isoform lb, alpha subunit (45kD) | acetylhydrolase Ž (40kD) (PAFAH2) | | | | | | | | | |
| receptor (PTAFR) | acetylhydrolase, isoform lb, alpha subunit (45kD) (PAFAH1B1) | | | + | | + | + | + | | |
| Dieckstrin (PLEK) 10 | | 1 | D10202 | | + | | | | + | |
| match pleckstrin hornology, Sec7 and coiled/coil domains 1(cytohesin 1) (PSCD1) pleckstrin hornology, Sec7 and coiled/coil domains 1(cytohesin 1) (PSCD1) pleckstrin hornology, Sec7 and coiled/coil domains, binding protein (PSCDBP) pM5 protein 1 | pleckstrin (PLEK) | | _ | | | + | + | | + | |
| Discription | | 1 | X07743 | | | | | | | |
| Dieckstrin homology, Sec7 and coiled/coil domains, binding protein (PSCDBP) Dipoly (PSCDBP) Dipo | pleckstrin homology, Sec7 and coiled/coil domains 1(cytohesin 1) (PSCD1) | 4 | M85169 | + | + | | + | | + | |
| DMS protein | pleckstrin homology, Sec7 and coiled/coil domains. | 4 | | + | | | + | | | |
| Doly (ADP-ribose) 1 | pM5 protein | | | + | + | + | + | | + | |
| polymerase (NAD (+) ADP- nbosyltransferase) (=X16674) poly(A) polymerase (PAP) | · · · · · · · · | | | | | | | | | |
| Doly(A)-binding protein-like | polymerase (NAD (+) ADP- ribosyltransferase) (=X16674) | | | | | | | | | |
| 1 (PABPL1) | 1 | | | <u> </u> | | | | | | |
| (PCBP1) | 11 (PABPL1) | | | | | | | | | |
| Description | poly(rC)-binding protein 1 (PCBP1) | 3 | 1112121 | + | + | + | + | + | + | |
| (autosomal dominant) (PKD1) polymerase (DNA | protein | 1 | | | | | | | | |
| directed), beta (POLB) polymerase (DNA directed), gamma (POLG) polymerase (RNA) II (DNA 1 X63564 + + + + + + + + + + + + + + + + + + + | (autosomal dominant) (PKD1) | - - | | | | | | | | |
| polymerase (DNA directed), gamma (POLG) polymerase (RNA) II (DNA 1 X63564 + + + + + + + + + + + + + + + + + + + | | 1 | D29013 | | + | | | + | + | |
| Dolymerase (RNA) II (DNA | polymerase (DNA | 6 | D84103 | | | | | | | |
| polymyositis/scleroderma 1 L01457 + + + + + + + autoantigen 2 (100kD) (PMSCL2) | polymerase (RNA) II (DNA directed) polypeptide A | 1 | X63564 | + | + | + | + | + | + | |
| | polymyositis/scleroderma autoantigen 2 (100kD) | 1 | | + | + | + | + | + | + | |
| protein (heterogeneous nuclear ribonucleoprotein I) (PTB) | polypyrimidine tract binding protein (heterogeneous nuclear ribonucleoprotein I) | 1 | X65372 | + | + | + | + | + | + | |

| WO 00/40749 | | | | | | | | 10 | .1/CA00/00003 |
|---|---|----------|---|---|---|----------|---|----|------------------|
| positive regulator of programmed cell death iCH-1L (Ich-1) | 3 | U13021 | | | + | | | | |
| postmeiotic segregation increased 2-like 12 (PMS2L12) | 1 | M16514 | + | + | + | + | | + | |
| postmeiotic segregation increased 2-like 8 | 1 | U38964 | + | + | + | + | | + | |
| (PMS2L8) potassium inwardly- rectifying channel, subfamily J, member 15 | 1 | D87291 | | | | + | | + | |
| (KCNJ15) potassium voltage-gated channel, KQT-like subfamily, member 1 | 1 | AF051426 | *************************************** | + | + | + | | + | |
| (KCNQ1) POU domain, class 2, associating factor 1 | 1 | Z49194 | | | | + | | | |
| (POU2AF1) POU domain, class 2, transcription factor 1 | 2 | X13403 | | + | | + | | | |
| (POU2F1) PPAR binding protein (PPARBP) | 1 | Y13467 | + | + | + | + | | + | |
| PPAR gamma2 pre-B-cell colony- | 8 | U02020 | | + | - | \vdash | | | |
| enhancing factor (PBEF) prefoldin 1 (PFDN1) | 1 | Y17392 | + | + | + | + | + | + | |
| prefoldin 5 (PRFLD5) | 3 | D89667 | В | + | + | - | + | | |
| prefoldin subunit 3 (=U96759 von Hippel- Lindau binding protein (VBP-1)) | 1 | Y17394 | | | | | | | |
| pregnancy-associated plasma protein A (PAPPA) | 1 | U28727 | | + | | + | | | high in placenta |
| pre-mRNA splicing factor SF3a (60kD), similar to S. cerevisiae PRP9 (spliceosome-associated protein 61) (SF3A60) | 1 | U08815 | + | + | + | + | | + | |
| pre-mRNA splicing factor SF3a (60kD), similar to S. cerevisiae PRP9 (spliceosome-associated protein 61) (SF3A60) (low score) | 1 | U08815 | | | | | | | , |
| pre-mRNA splicing factor SRp20, 5'UTR | 2 | D28423 | | | | | | | |
| preprotein translocase (TIM17) | 3 | X97544 | + | + | + | + | | + | |
| prion protein | 1 | X82545 | | | | | | | |
| prion protein (p27-30) (Creutzfeld-Jakob disease, Gerstmann-Strausler- Scheinker syndrome, fatal familial insomnia) (PRNP) | | M13899 | | + | + | + | | + | |
| pristanoyl-CoA oxidase (low match) | 1 | Y11411 | | | | | | | |
| pristanoyl-CoA oxidase (low score) | 1 | Y11411 | | ļ | | Ļ | | _ | |
| procollagen-lysine, 2- oxoglutarate 5- dioxygenase (lysine hydroxylase, Ehlers-Danlos syndrome type VI) (PLOD) | 1 | M98252 | | + | + | + | | + | · |
| procollagen-proline, 2- oxoglutarate 4- dioxygenase (proline 4- hydroxylase), alpha polypeptide 1 (P4HA1) | 1 | M24486 | + | + | + | + | + | + | |

| WO 00/40749 | | | | | | | | PC | [/CA00/00005 |
|---|----------|------------------|---|---|----------|----------|---|----|--------------|
| procollagen-proline, 2- oxoglutarate 4- dioxygenase (proline 4- hydroxylase), beta polypeptide (protein disulfide isomerase; thyroid hormone binding protein p55) (P4HB) | 4 | X05130 J03191 | + | + | + | + | + | + | |
| profilin 1 (PFN1) | | | • | | <u> </u> | <u> </u> | | · | |
| progesterone receptor- associated p48 protein (P48) | 2 | U28918 | | + | | + | + | + | |
| prohibitin (PHB) | 1 | S85655 | | + | + | L | + | | |
| proliferating cell nuclear antigen (PCNA) | 3 | J04718 | + | + | + | + | + | + | |
| proliferation-associated gene A (natural iller- enhancing factor A) (PAGA) | 4 | L19184 | • | + | + | Ť | | | |
| proline-rich protein BstNI subfamily 2 (PRB2) (non- exact, 43%aa) | | S62936 | | | | | | | |
| proline-serine-threonine phosphatase interacting protein 1 (PSTPIP1) | 1 | | | + | | + | | + | |
| prolyl endopeptidase (PREP) | 2 | X74496 | | * | | * | | * | |
| prolylcarboxypeptidase (angiotensinase C) (PRCP) | 5 | L13977 | | + | + | + | + | + | |
| promyelocytic leukemia (PML) | 1 | M80185 | + | + | + | + | | + | |
| properdin P factor, complement (PFC) | 4 | X57748 | + | | | | | | |
| pro-platelet basic protein (includes platelet basic protein, beta- thromboglobulin, connective tissue-activating peptide III, neutrophil-activating peptide-2) (PPBP) | 1 | M54995 | | | + | + | + | + | |
| pro-platelet basic protein (includes platelet basic protein, beta-thromboglobulin, connective tissue-activating peptide III, neutrophil-activating peptide-2) (PPBP) | 7 | M54995 | + | | + | | | | |
| proprotein convertase subtilisin/kexin type 7 (PCSK7) | 4 | U40623 | | | | | | | |
| prosaposin (variant Gaucher disease and variant metachromatic leukodystrophy) (PSAP) | 89 | D00422 | + | + | + | + | + | + | |
| prostaglandin- endoperoxide synthase 1 (prostaglandin G/H synthase and cyclooxygenase) (PTGS1) | 1 | U63846 | В | + | | | + | + | |
| prostaglandin- endoperoxide synthase 2 (prostaglandin G/H synthase and cyclooxygenase) (PTGS2) | 2 | L15326 | | | | | | | |
| prostaglandin- endoperoxide synthase-1 (=L08404; U84208) (all promoters) | | D64068 | , | | | | | | |
| prostate carcinoma tumor antigen (pcta-1) | 2 | L78132 | | | | | | | |

PCT/CA00/00005

| WO 00/40749 | | | | | | | | PC | T/CA00/00005 |
|--|----------|----------|----------|-----|---|---|-----|----|------------------------|
| protease inhibitor 1 (anti- | 17 | K02212 | | + 1 | + | + | + 1 | + | high in many libraries |
| elastase), alpha-1- antitrypsin (PI) | •• | | | | | | | | |
| protease inhibitor 2 (anti- elastase). | 1 | M93056 | | | | + | | + | |
| monocyte/neutrophil (ELANH2) (low match) | <u>_</u> | | | + | _ | | | + | |
| proteasome (prosome, macropain) 26S subunit, ATPase, 1 (PSMC1) | 3 | L02426 | В | | + | | | • | |
| proteasome (prosome, macropain) 26S subunit, ATPase, 3 (PSMC3) | 1 | М34079 | + | + | + | + | | + | |
| proteasome (prosome, macropain) 26S subunit, ATPase, 4 (PSMC4) | 2 | AF020736 | - | | - | | | | |
| proteasome (prosome, macropain) 26S subunit, ATPase, 5 (PSMC5) | 5 | L38810 | + | + | + | + | + | + | |
| proteasome (prosome, macropain) 26S subunit, ATPase, 6 (PMSC6) | 2 | D78275 | + | + | + | + | | + | |
| proteasome (prosome, macropain) 26S subunit, non-ATPase, 11 (PSMD11) | 7 | AF001212 | 7 " | + | | | + | | |
| proteasome (prosome, macropain) 26S subunit, non-ATPase, 2 (PSMD2) | 2 | D78151 | | + | + | | | + | |
| proteasome (prosome, macropain) 26S subunit, non-ATPase, 5 (PSMD5) | 1 | S79862 | T | + | + | | + | | |
| proteasome (prosome, macropain) 26S subunit, non-ATPase, 7 (Mov34 | 1 | D50063 | - | + | + | + | | + | high in many libraries |
| homolog) (PMSD7) proteasome (prosome, macropain) 26S subunit, on-ATPase, 12 (PMSD12) | 1 | AB003103 | | + | + | + | | + | |
| proteasome (prosome, macropain) activator subunit 1 (PA28 alpha) | š | L07633 | + | + | + | + | | + | |
| (PSME1) proteasome (prosome, macropain) subunit, alpha type, 3 (PSMA3) | 2 | D00762 | <u> </u> | + | + | + | | + | , |
| proteasome (prosome, macropain) subunit, alpha type, 5 (PSMA5) | 3 | X61970 | + | + | + | + | | + | |
| proteasome (prosome, macropain) subunit, alpha type, 7 (PSMA7) | 3 | AF054185 | | + | + | + | + | + | |
| proteasome (prosome, macropain) subunit, alpha type, 7 (PSMA7) (low match) | 1 | AF022815 | | | | | | | |
| proteasome (prosome, macropain) subunit, beta type, 1 (PSMB1) | 1 | D00761 | + | + | + | + | + | + | |
| proteasome (prosome, macropain) subunit, beta type, 10 (PSMB10) | 1 | X71874 | + | + | | + | + | + | |
| proteasome (prosome, macropain) subunit, beta type, 6 (PMSB6) | 1 | D29012 | | + | + | + | | + | |
| proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional protease 7) | 1 | U17497 | + | + | + | + | | + | |
| (PSMB8) proteasome (prosome, macropain) subunit, beta type, 9 (large multifunctional protease 2) (PSMB9) | 3 | Z14977 | + | | | + | | + | |

| WO 00/40749 | | | | | | | | PC | 1/CA00/00005 |
|---|--------------|----------|------------------|----------|----------|----------|---|----------|------------------------|
| proteasome (prosome, macropain) subunit, beta ype, 7 (PSMB7) | 1 | D38048 | + | + | + | + | + | + | |
| protective protein for beta- galactosidase | 3 | M22960 | + | + | + | + | + | + | |
| (galactosialidosis) (PPGB) protein A alternatively spliced form 2 (A-2) | 1 | U47925 | | + | | <u> </u> | | | |
| protein activator of the interferon-induced protein kinase (PACT) | 1 | AF072860 | | + | + | + | | + | high in testis |
| protein disulfide isomerase- related protein (P5) | 2 | D49489 | + | + | + | + | + | + | |
| protein geranylgeranyltransferase type I, beta subunit (PGGT18) | 1 | L25441 | + | + | + | | | | |
| protein homologous to chicken B complex protein, guanine nucleotide binding (H12.3) | 20 | M24194 | + | + | + | + | + | + | high in many libraries |
| protein kinase A anchoring protein | 1 | AF037439 | | + | | | | | |
| protein kinase C substrate 80K-H (PRKCSH) | 2 | U50317 | + | + | + | + | | + | |
| protein kinase C, beta 1 (PRKCB1) | 6 | X06318 | + | + | + | + | | + | |
| protein kinase C, delta (PRKCD) | | D10495 | + | + | + | + | | + | |
| protein kinase C, eta (PRKCH) | 1 | M55284 | | | + | ļ | | + | |
| protein kinase C, mu (PRKCM) (non-exact 78%) | 1 | X75756 | | <u> </u> | ļ | - | _ | <u>_</u> | |
| Protein kinase C-like 1 (PRKCL1) | 2 | D26181 | + | + | + | Ļ | | Ľ | |
| protein kinase, AMP- activated, gamma 1 non- catalytic subunit (PRKAG1) | | U42412 | B, T lymphoma | • | , | | | | |
| protein kinase, cAMP- dependent, regulatory, type I, alpha (tissue specific extinguisher 1) (PRKAR1A) | 4 | M18468 | | + | + | + | + | + | |
| protein kinase, DNA- activated, catalytic polypeptide (PRKDC) | 1 | U47077 | | + | + | | + | + | |
| protein kinase, mitogen- activated 1 (MAP kinase 1; p40, p41) (PRKM1) | | Z11695 | В | + | | | + | | |
| protein kinase, mitogen- activated 6 (extracellular signal-regulated kinase, p97) (PRKM6) | 1 | L77964 | | + | | + | + | + | |
| protein kinase, mitogen- activated, kinase 3 (MAP kinase kinase 3) (PRKMK3) | 1 | U66839 | + | + | + | + | + | | |
| protein phosphatase 1, catalytic subunit, alpha isoform (PPP1CA) | 5 | M63960 | + | + | + | + | + | + | |
| protein phosphatase 1, regulatory subunit 10 (PPPR10) | 3 | Y13247 | | + | + | + | | + | |
| protein phosphatase 1, regulatory subunit 7 (PPP1R7) | 2 | Z50749 | + | + | + | + | + | + | |
| protein phosphatase 2 (formerly 2A), catalytic subunit, beta isoform (PPP2CB) | 1 | X12656 | + | + | + | + | + | + | |
| protein phosphatase 2 (formerly 2A), regulatory subunit B" (PR 72), alpha isoform and (PR 130), beta isoform (PPP2R3) | 1 | L07590 | | | + | + | | + | |
| | | | 20 | | | • | · | | |

| WO 00/40749 | | | | | | | | | 1/CA00/00003 | |
|--|----|----------|----|---|---|---|---|---|----------------|---|
| protein phosphatase 2, regulatory subunit B (856), alpha isoform (PPP2R5A) | 2 | L42373 | + | + | + | + | | + | |] |
| protein phosphatase 2, regulatory subunit B (B56), delta isoform (PPP2R5D) | 3 | D78360 | | + | + | + | | + | |] |
| protein phosphatase 2, regulatory subunit B (B56), gamma isoform (PPP2R5C) | 1 | D26445 | + | + | + | + | | + | | |
| protein phosphatase 2A regulatory subunit alpha- isotype (alpha-PR65) | 5 | J02902 | + | + | + | + | | + | | |
| protein phosphatase 4 (formerly X), catalytic subunit (PPP4C) | 2 | AF097996 | + | + | + | + | | + | | |
| protein tyrosine kinase 2 beta (PTK2B) | 4 | L49207 | | + | | + | | + | | |
| protein tyrosine phosphatase epsilon | 1 | X54134 | | | | | | | | |
| protein tyrosine phosphatase type IVA, member 2 (PTP4A2) | 2 | L48723 | + | + | + | + | | + | | |
| protein tyrosine phosphatase, non-receptor type 1 (PTPN1) | 1 | M31724 | + | + | + | + | | | | |
| protein tyrosine phosphatase, non-receptor type 12 (PTPN12) | 1 | M93425 | | + | + | + | | + | high in testis | |
| protein tyrosine phosphatase, non-receptor type 12 (PTPN12) (non- exact, 70%) | 1 | M93425 | | | | | | | | |
| protein tyrosine phosphatase, non-receptor type 2 (PTPN2) | 2 | M25393 | | + | + | + | | + | | |
| protein tyrosine phosphatase, non-receptor type 4 (megakaryocyte) (PTPN4) | 1 | M68941 | | | + | + | | + | | |
| protein tyrosine phosphatase, non-receptor type 6 (PTPN6) | 7 | M74903 | + | + | + | + | | + | | |
| protein tyrosine phosphatase, non-receptor type 7 (PTPN7) | 1 | D11327 | + | | | + | | + | | |
| protein tyrosine phosphatase, receptor type, alpha polypeptide (PTPRA) | 1 | M34668 | + | + | + | + | | + | | |
| protein tyrosine phosphatase, receptor type, c polypeptide (PTPRC) | 44 | Y00638 | + | + | | + | | + | | |
| protein tyrosine phosphatase, receptor type, M (PTPRM) | 1 | X58288 | | + | + | * | | + | | |
| protein tyrosine phosphatase, receptor type, N polypeptide 2 (PTPRN2) | 2 | U81561 | | + | | + | | + | | |
| protein with polyglutamine repeat (ERPROT213-21) | 1 | U94836 | + | + | + | + | | + | | |
| protein-kinase, interferon- inducible double stranded RNA dependent inhibitor (PRKRI) | 1 | U28424 | | + | + | + | + | + | | |
| protein-L-isoaspartate (D- aspartate) O- methyltransferase (PCMT1) | 4 | D13892 | | + | + | | | | | |
| proteoglycan 1, secretory granule (PRG1) | 7 | J03223 | | + | | + | | + | | |
| prothymosin, alpha (gene sequence 28) (PTMA) | 12 | M14483 | + | + | + | + | + | + | | |
| | | | 21 | | | | | | | _ |

| prp28, U5 snRNP 100 kd protein (U5-100K) | 7 | AF026402 | + | . + | + | + | | + | |
|--|--------------|----------|---|--------------|---|--|----|--|--|
| PRP4/STK/WD splicing factor (HPRP4P) | | AF001687 | | + | + | + | | + | |
| PTK7 protein tyrosine | 1 | U40271 | | + | + | + | ┢╾ | + | |
| kinase 7 (PTK7) purinergic receptor P2X, | | AF000234 | | + | + | + | ├ | + | |
| ligand-gated ion channel, 4 (P2RX4) | 3 | AF000254 | | | , | * | | ľ | |
| purinergic receptor P2X, ligand-gated ion channel, 7 (P2RX7) | 1 | Y12851 | + | | | | | | macrophage only |
| puromycin-sensitive aminopeptidase (PSA) | 1 | Y07701 | | + | + | | | + | |
| putative ATP(GTP)-binding protein | 2 | AJ010842 | | + | | | | + | |
| putative brain nuclearly- targeted protein (KIAA0765) | 1 | AB018308 | + | + | + | + | | + | |
| putative chemokine receptor; GTP-binding protein (HM74) | 1 | D10923 | + | | | | | | |
| putative dienoyl-CoA | 1 | AF030249 | | | | - | | | |
| putative G-binding protein | 1 | AF065393 | | + | | | | | |
| Putative human HLA class Il associated protein I (PHAP1) | 7 | U73477 | В | + | | | + | | |
| Putative L-type neutral amino acid transporter (KIAA0436) | 1 | AB007896 | | | | | | | |
| putative mitochondrial space protein 32.1 | 1 | AF050198 | | | | | | | |
| PUTATIVE MUCIN CORE PROTEIN PRECURSOR 24 (MULTI- IGLYCOSYLATED CORE | 1 | Q04900 | | | | | | | |
| PROTEIN 24) (MGC-24) (MUC-24) | | | | | | | | | |
| putative nucleic acid binding protein | 2 | X76302 | + | + | + | + | | + | |
| putative outer mitochondrial membrane 34 kDa translocase Htom34 | 1 | U58970 | | + | + | + | | + | |
| putative p150 (non-exact 88%) | 1 | U93568 | | | | | | | |
| putative translation initiation factor (SUI1) | 1 | L26247 | + | + | + | + | + | + | High in moderately differentiated colon adenocarcinoma |
| putative tumor suppressor protein (123F2) | 1 | AF061836 | | + | + | + | | + | |
| pyrroline 5-carboxylate reductase | 1 | M77836 | + | + | + | + | | + | |
| pyruvate dehydrogenase (lipoamide) alpha 1 (PDHA1) | 1 | D90084 | | + | + | + | + | + | |
| pyruvaté dehydrogenase (lipoamide) beta (PDHB) | 2 | J03576 | + | + | + | + | | + | |
| Pyruvate dehydrogenase complex, lipoyl-containing component X; E3-binding protein (PDX1) | 3 | Y13145 | | + | + | | | | |
| pyruvate kinase, muscle (PKM2) | 11 | M23725 | | | | | + | | |
| RAB, member of RAS oncogene family-like (RABL) | 1 | U18420 | | + | + | + | | + | |
| RAB1, member RAS oncogene family (RAB1) | 3 | M28209 | | + | + | + | | + | |
| RAB11A, member RAS | 2 | X56740 | + | + | + | + | | + | high in spleen |
| oncogene family (RAB11A) | | 1 | | | L | Ь | | L | L |

| V, C 00, 10, 15 | | | | | | | | | | _ |
|---|-------------|-----------|-------------|--|--|--------------|--|--|---------------|-----------|
| RAB11B, member RAS | 1 | D45418 | | + | | | | + | | - [|
| oncogene family (Rab11B) RAB27A, member RAS | 3 | U38654 | | | | + | - | | | ┥. |
| oncogene family (RAB27A) | 3 | | | | | | | | | _] ` |
| RAB5B, member RAS | 1 | X54871 | | + | + | + | | + | | |
| oncogene family (RAB5B) RAB6, member RAS | 1 | M28212 | | + | | | | + | | -1 |
| oncogene family (RAB6) | | | | | | | | | | 4 |
| RAB7, member RAS | 1 | X93499 | + | + | + | + | - 1 | + | | ı |
| oncogene family (RAB7) RAB7, member RAS | 2 | D84488 | | + | + | + | | + | | ヿ゙ |
| oncogene family-like 1 | | | | | | | Ì | | | - 1 |
| (RAB7L1) RAB9, member RAS | | U44103 | | | | | | - | | \dashv |
| oncogene family (RAB9) | | | | | | | | | | ┛ |
| RAD50 (S. cerevisiae) | 2 | U63139 | | + | + | + | | | | |
| homolog (RAD50) RAD51 (S. cerevisiae) | 1 | AF029669 | | + | + | + | | + | | ┪ |
| homolog C (RAD51C) | | | | | | L_ | | | | 4 |
| Radin blood group (RD) | 2 | L03411 | | + | + | + | | + | | 4 |
| RAE1 (RNA export 1, S.pombe) homolog (RAE1) | 3 | U84720 | + | + | + | + | | + | | |
| ralA-binding protein | 2 | L42542 | + | + | + | + | | | | \exists |
| (RLIP76) | | AF012086 | | ļ | | | | | | \dashv |
| RAN binding protein 2-like 1 (RANBP2L1) | 2 | AF012066 | | 1 | | | | | | |
| Ran GTPase activating | 3 | X82260 | + | + | + | + | | + | | |
| protein 1 (RANGAP1) RAN, member RAS | 1 | M31469 | | | | | | | | \dashv |
| oncogene family (RAN) | | 11107400 | | | | | | | | |
| (low match) | | D42063 | | ↓ | | | | | | -1 |
| RanBP2 (Ran-binding protein 2) (=U19248; | 1 | D42003 | | | | | | | | |
| L41840 sapiens | | | | | | | | | | |
| nucleoporin (NUP358)) ransforming growth factor, | 4 | D50683 | + | ╁ | + | + | | + | | \dashv |
| beta receptor II (70-80kD) | | | | | | 1 | | | | |
| (TGFBR2) RAP1A, member of RAS | 10 | M22995 | + - | + | + | + | + | + | | ᅱ |
| oncogene family (RAP1A) | '0 | | | | | | | | | _ |
| RAR-related orphan | 1 | U16997 | | 1 | | | | + | | |
| receptor C (RORC) RAS guanyl releasing | 1 | Y12336 | + | + | - | | - | | | \neg |
| protein 2 (calcium and | | | | | | | 1 | | | |
| DAG-regulated) ras homolog gene family, | 12 | X05026 | + | + | + | + | + | + | high in ovary | ᅱ |
| member A (ARHA) | | | | | | <u> </u> | | | | _ |
| ras homolog gene family, member G (rho G) (ARHG) | 1 | X61587 | + | + | + | + | | | | |
| ras homolog gene family, | 2 | Z35227 | + | + | + | 1 | | + | | \exists |
| member H (ARHH) | | M37191 | | + | ├ | - | \vdash | | | - |
| ras inhibitor (RIN1) | 2 | AF053535 | + | + + | + | + | - | + | | \dashv |
| Ras-GTPase activating protein SH3 domain- | | 71 033333 | · | | _ | | } | | | |
| binding protein 2 | | | | | ļ | | | ļ | | |
| (KIAA0660) Ras-GTPase-activating | 3 | U32519 | + | + | + | + | - | + | | \dashv |
| protein SH3-domain- | • | | | | | | | | | - |
| binding protein (G3BP) ras-related C3 botulinum | 11 | M29871 | | ┼ | + | + | - | + | | \dashv |
| toxin substrate 2 (rho | '' | | | | | | | | | |
| family, small GTP binding protein Rac2) (RAC2) | | | | | | | | | | |
| RAS-RELATED PROTEIN | 1 | P09526 | | 1 | | 1 | | | <u> </u> | \exists |
| RAP-1B (GTP-BINDING | | | | - | 1 | | | İ | | |
| PROTEIN SMG P21B) | | X85133 | | + | + | + | - | | | \dashv |
| rearranged T cell receptor | 1 | L06891 | | + | | + | | \vdash | | \dashv |
| beta variable region (TCRB) (=X58810) | | | | | | | | | | _ |
| regulator of Fas-induced apoptosis (TOSO) | 1 | AF057557 | В | | | | + | | | |
| apoptosis (1030) | l | | | | ь | | ь | Ц | | |

| VI O 00/40/42 | | | | | | | | | | |
|---|-----|----------|------|---|---|---|---|---|-------|--|
| regulator of G protein signalling 6 (RGS6) | 1 | AF073920 | - | + | | | | | | |
| regulator of G-protein | 2 | AF037195 | + | + | + | + | | 1 | _ | |
| signalling 14 (RGS14) regulator of G-protein | - 6 | L13391 | + | + | + | + | | + | | |
| signalling 2, 24kD (RGS2) | U | | · | | | | | | | |
| regulator of G-protein signalling 5 (RGS5) (49% | 1 | 015539 | | | | | | | | |
| regulatory factor X, 4 | 1 | M69297 | | - | + | + | | | | |
| (influences HLA class II expression) (RFX4) | , | | | | | | | | | |
| regulatory factor X, 5 (influences HLA class II expression (RFX5) | 2 | X85786 | Т | + | + | | | + | | |
| replication protein A1 (RPA1) | 1 | M63488 | + | + | + | + | | + | | |
| replication protein A3 (14kD) (RPA3) (low match) | 1 | L07493 | | | | | | | | |
| reproduction 8 (D8S2298E) | 1 | D83767 | | + | + | + | | | | |
| requiem, apoptosis response zinc finger gene (REQ) | 2 | U94585 | + | + | + | + | | + | | |
| requiem, apoptosis response zinc finger gene (REQ) (=AF001433) (low match) | 1 | U94585 | | | | | | | | |
| restin (Reed-Steinberg cell- expressed intermediate filament-associated | 1 | M97501 | В, Т | + | + | | | | | |
| protein) (RSN) retinoblastoma 1 (including osteosarcoma) (RB1) | 3 | L11910 | + | + | + | + | | | | |
| retinoblastoma binding protein 2 homolog 1 (RBBP2H1) | 1 | AF087481 | | | | | | | | |
| retinoblastoma-binding protein 1 (RBBP1) | 1 | S66427 | + | + | | | | | | |
| retinoblastoma-binding protein 2 (RBBP2) | 5 | S66431 | + | + | + | + | | + | | |
| retinoblastoma-binding protein 4 (RBBP4) | 1 | X71810 | | + | + | + | | + | | |
| retinoblastoma-binding protein 4 (RBBP4) | 1 | X74262 | | + | + | + | | + | | |
| retinoblastoma-binding protein 7 (RBBP7) | 1 | U35143 | | | | | | | | |
| retinoblastoma-like 2 (p130) (RBL2) | 1 | X76061 | | + | + | + | | + | | |
| retinoic acid receptor responder (tazarotene nduced) 3 (RARRES3) | 1 | AF060228 | | + | | + | + | + | | |
| retinoic acid receptor, alpha (RARA) | 1 | X06538 | + | + | | + | | | | |
| retinoic acid responsive (NN8-4AG) | 1 | U50383 | | + | | + | | + | | |
| retinoid X receptor beta (RXR-beta) | 2 | X66424 | | + | + | + | | + | | |
| REV3 (yeast homolog)-like, catalytic subunit of DNA polymerase zeta (REV3L) | 1 | AF035537 | | | | | | | | |
| Rho GDP dissociation inhibitor (GDI) beta (ARHGDIB) | 23 | L07916 | + | + | + | + | + | + | | |
| Rho GTPase activating protein 4 (ARHGAP4) | 2 | X78817 | + | + | | | | | | |
| Rho GTPase activating protein 4 (ARHGAP4) (low match) | 1 | P98171 | | | | | | | | |
| Rho-associated, coiled-coil containing protein kinase 2 (ROCK2) | 1 | AB014519 | | | | | | | | |
| ribonuclease 6 precursor (RNASE6PL) | 2 | U85625 | + | + | + | + | + | + | | |
| N | | ······ | | | | | | | | |

| nbonuclease 6 precursor | | U85625 | | T 1 | · · · | | | Γ | |
|--|-------------|--------|---------------|----------|----------|----------|----------|----------|--------------------------------------|
| (RNASE6PL) (low match) | | X55988 | | | | | + | <u> </u> | |
| family, 2 (liver, eosinophilderived neurotoxin) (RNASE2) | 1 " | 733966 | | | | | Ť | | |
| ribonuclease/angiogenin inhibitor (RNH) | 3 | M36717 | + | + | + | + | <u> </u> | + | |
| ribonucleoside diphosphate reductase M1 subunit | 1 | X65708 | | <u> </u> | | | | | |
| nbonucleotide reductase M2 polypeptide (non-exact 91%) | 1 | P31350 | | | | | | | |
| ribophorin I (RPN1) | 1 | Y00281 | + | + + | + | + | _ | + | |
| ribophorin II (RPN2) | 1 | Y00282 | + | + | + | + | + | + | |
| ribosomal 18S rRNA | 3 | M10098 | | | | | | | |
| ribosomal 285 RNA | 1 | M11167 | | | ļ | | | <u> </u> | |
| ribosomal phosphoprotein P0, 5'UTR (low match) | 1 | D28418 | - | | | _ | | | |
| Ribosomal protein | 30 | L25899 | + | + | + | + | + | + | high in many libraries |
| (RPL10) RIBOSOMAL PROTEIN | | P53025 | * | | <u> </u> | Ĺ | | Ļ | right in many incidies |
| L10A (CSA-19) | _ | | | | <u> </u> | <u> </u> | | | |
| ribosomal protein L11 (RPL11) | 4 | X79234 | + | + | + | + | + | + | Alveolar rhabdomyosarcoma |
| ribosomal protein L12 (RPL19) | 2 | L06505 | + | + | + | + | + | + | |
| ribosomal protein L13 (PRL13) | 1 | P26373 | + | + | + | + | + | + | high in many libraries |
| nbosomal protein L14 (RPL14) | 4 | D87735 | + | + | + | + | + | + | high in many libraries |
| ribosomal protein L17 (RPL17) | 4 | X53777 | + | | | | | | blood only |
| ribosomal protein L18 (RPL18) | 10 | L11566 | + | + | + | + | | + | |
| nbosomal protein L18a (RPL18A) | 5 | L05093 | | + | + | + | + | + | High in fetal adrenal gland and skin |
| ribosomal protein L18a homologue | 2 | X80821 | | | | + | | | |
| nbosomal protein L19 (RPL19) | 15 | X63527 | + | + | + | + | + | + | |
| ribosomal protein L21 (RPL21) | 6 | U14967 | + | + | + | + | + | + | |
| ribosomal protein L22 (RPL22) | 3 | D17652 | + | + | + | + | | + | |
| ribosomal protein L23 (RPL23) | 2 | X55954 | + | + | + | + | + | + | high in many libraries |
| ribosomal protein L23a (RPL23A) | 5 | U37230 | + | + | + | + | + | + | high in many libraries |
| ribosomal protein L26 (RPL26) | 8 | X69392 | + | + | * | + | <u> </u> | Ļ | |
| ribosomal protein L27 (RPL27) | 6 | L05094 | + | + | + | + | <u> </u> | + | |
| ribosomal protein L27a (RPL27A) | 10 | U14968 | + | + | + | + | + | | |
| ribosomal protein L28 (RPL28) | 6 | U14969 | + | + | + | + | <u> </u> | + | |
| ribosomal protein L29 (RPL29) | 6 | U10248 | + | + | + | + | + | + | |
| ribosomal protein L3 (RPL3) | 81 | | + | + | + | + | + | | high in many libraries |
| ribosomal protein L3 homologue | 81 | X06323 | | | | | Ŀ | | |
| ribosomal protein L30 (RPL30) | 6 | X79238 | + . | + | + | + | + | + | high in lymphoma |
| ribosomal protein L30 (RPL30) (low score) | 1 | X79238 | | | | | | | |
| ribosomal protein L31 (RPL31) | 10 | X15940 | + | + | + | + | + | + | High in alveolar rhabdomyosarcoma |

| 3 | X03342 | + | + | + | + | + | + | |
|----|---|--|---|---|------------|------------|------------|--|
| 1 | ' | | + | + | | | | |
| 5 | | | + | + | + | + | + | |
| 1 | | _ | | | | | | |
| 5 | D23661 | + | | | | | | high in barstead prostate |
| 4 | X66699 | + | + | + | + | + | | high in many libraries |
| 1 | Z26876 | + | + | + | + | + | | high in many libraries |
| 27 | | + | + | + | + | | | high in many libraries |
| | | | | | | | | high in many libraries |
| | | + | + | + | + | + | | High in alveolar rhabdomyosarcoma |
| 1 | | | | | | | | |
| 7 | | | | | | | | high in many libraries |
| 14 | | + | | | l | | ŀ | high in conorm |
| | | | | | | | | High in uterus, and seminoma |
| 5 | | + | | | | | | high in ovary |
| | | | | | | | | |
| | | | | | | | | high in many libraries |
| | | + | + | + | | | + | high in many libraries |
| | | | | | | | L | |
| | | + | | | <u> </u> | | | high in many libraries |
| | | | | | | | | |
| | | | | | | | i | |
| | | | | | l | | | Uish in acceptate |
| | | | | | <u> </u> | | | High in prostate invasive tumor |
| 2 | | + | + | + | | | Ľ | high in many libraries |
| | | | | <u> </u> | <u> </u> | | <u> </u> | |
| 7 | | + | + | + | + | + | <u> </u> | high in many libraries |
| 4 | X17206 | + | + | + | + | + | + | high in many libraries |
| | P15880 | | | | | | | |
| 7 | L06498 | + | + | + | + | ļ | | high in many libraries |
| 3 | <u>'</u> | + | + | + | | + | | high in CD34+/CD38- hematopoietic cells and skin tumor |
| 3 | D14530 | | + | + | + | | + | |
| 7 | M31520 | + | + | + | + | + | + | high in uterus |
| 3 | M64716 | + | + | + | + | + | + | high in barstead prostate |
| 2 | X69654 | | +_ | + | + | + | + | |
| 5 | U57847 | + | + | + | + | + | + | |
| | 1 5 1 5 4 1 7 14 15 5 10 5 4 1 3 2 12 2 3 2 8 7 4 2 7 3 3 7 3 2 12 12 12 12 12 12 12 12 12 12 12 12 1 | 1 AF047440 5 L38941 1 L38941 5 D23661 4 X66699 1 Z26876 27 D23660 4 AF026844 14 U14966 1 U14966 7 X69391 14 X52967 15 M36072 5 Z28407 10 U09953 5 U14972 4 X06617 1 AB007152 3 X53505 2 L01124 12 M13934 2 M32405 3 M60854 2 M13932 8 X69150 7 M81757 4 X17206 2 P15880 7 L06498 3 D14530 7 M31520 3 M64716 2 X69654 | 1 AF047440 5 L38941 1 L38941 5 D23661 + 4 X66699 + 1 Z26876 + 27 D23660 + 4 AF026844 + 14 U14966 + 1 U14966 7 X69391 + 14 X52967 + 15 M36072 + 5 Z28407 + 10 U09953 5 U14972 + 4 X06617 + 1 AB007152 3 3 X53505 + 2 L01124 12 M13934 + 2 M32405 + 3 M60854 + 2 M13932 + 4 X17206 + 2 P15880 7 L06498 + 3 L04483 + 3 D14530 7 M31520 + 3 M60716 + 2 X69654 | 1 AF047440 + + 5 L38941 + 1 L38941 + 4 X66699 + | 1 AF047440 | 1 AF047440 | 1 AF047440 | 1 |

| ribosomal protein S28 (RPS28) | 3 | U58682 | + | + | + | + | | + | |
|--|----------|----------|-------------|---|----------|----------|--|----|--|
| ribosomal protein S29 (RPS29) | 2 | U14973 | + | + | + | + | + | + | |
| ribosomal protein S3 (RPS3) | 9 | X55715 | + | + | + | + | + | + | high in many libraries |
| ribosomal protein S3 (RPS3) (low match) | 1 | U14990 | | | | | | | |
| ribosomal protein S3A (RPS3A) | 21 | Z83334 | | + | + | + | + | + | high in many libraries |
| ribosomal protein S3A (RPS3A) (low score) | 1 | M77234 | | | | | | | |
| ribosomal protein S4, X- linked (RPS4X) | 9 | M58458 | + | + | + | + | | + | high in ovary and Synovial sarcoma |
| nbosomal protein S4, Y- linked (RPS4Y) | 2 | M58459 | + | + | + | + | + | + | |
| ribosomal protein S5 (RPS5) | 4 | U14970 | + | + | + | + | + | + | high in lymphoma |
| RIBOSOMAL PROTEIN S6 (PHOSPHOPROTEIN NP33) | 1 | P10660 | | | | | | | |
| ribosomal protein S6 (RPS6) | 22 | M20020 | + | + | + | + | + | + | |
| ribosomal protein S6 (RPS6) (non-exact 86%) | 1 | M77232 | | | | | | | |
| ribosomal protein S6 kinase, 90kD, polypeptide 1 (RPS6KA1) | 3 | L07597 | + | + | + | + | | + | |
| ribosomal protein S6 kinase, 90kD, polypeptide 2 (RPS6KA2) | 1 | X85106 | | | | | | | |
| ribosomal protein S7 (RPS7) | 4 | Z25749 | | + | + | + | + | + | |
| ribosomal protein S8 (RPS8) | 6 | X67247 | | + | + | + | + | + | |
| ribosomal protein S9 (RPS9) | 8 | U14971 | | | | | | | colon tumor |
| nbosomal protein, large, P0 (RPLP0) | | M17885 | T | | + | | | + | , |
| ribosomal protein, large, P1 (RPLP1) | 12 | M17886 | T | + | + | | + | | |
| ribosomal RNA 18S (=M10098; K03432) (=polyadenylating sequence) | 11 | X03205 | | | | | | | |
| ribosomal RNA 28S | 2 | M11167 | | | | | | | |
| ribosomal RNA, 16S | 1 | U25123 | | | | | | | |
| ring finger protein (non- exact 58%) | 1 | AJ001019 | | | <u> </u> | | | | |
| ring finger protein 3 (RNF3) | 1 | AJ001019 | | | | <u> </u> | 1 | | 1 |
| ring finger protein 4 (RNF4) | 3 | AB000468 | | + | + | + | | + | |
| ring zinc-finger protein | 3 | U41315 | | + | + | + | - | + | |
| (ZNF127-Xp) RNA (guanine-7-) | 1 | AB007858 | | + | + | + | - | + | |
| methyftransferase (RNMT) RNA binding motif protein 5 (RBM5) | 4 | U23946 | + | + | + | + | - | + | |
| RNA binding motif, single stranded interacting protein 2 (RBMS2) | 1 | D28483 | | + | | + | _ | + | |
| RNA helicase (putative), (Myc-regulated DEAD box protein) (MRD8) | 1 | X98743 | + | + | + | + | | + | · |
| RNA helicase-related protein | 1 | AF083255 | | + | + | + | | + | |
| RNA pol Il largest subunit | 2 | X74872 | - | | · | Ī | | | |
| RNA polymerase I subunit (RPA40) | 1 | AF008442 | | + | + | | | + | |
| RTVP-1 protein | 2 | X91911 | + | + | + | + | t | + | |
| Tr. VI - I protoni | <u> </u> | 1,0,0,, | | | Ь | <u> </u> | Ь. | Щ. | <u> </u> |

| S100 calcium-binding protein A10 (annexin II ligand, calpactin I, light polypeptide (p11)) | 2 | M81457 | | | + | | + | + | |
|---|----|----------|---|---|---|---|----------|---|---|
| (S100A10) | | | | | | | l | | i |
| S100 calcium-binding protein A11 (calgizzarin) (S100A11) | 1 | X80201 | - | + | + | + | - | + | |
| S100 calcium-binding protein A4 (calcium protein, calvasculin, metastasin, murine placental homolog)(S100A4) | 3 | M80563 | 8 | | + | | + | | |
| S100 calcium-binding protein A8 (calgranulin A) (S100A8) | 7 | M21005 | | | + | + | | + | high in bone marrow |
| S100 calcium-binding protein A9 (calgranulin B) (S100A9) | 14 | X06233 | | | + | + | | | high in invasive larynx squamous cell carcinoma |
| S164 gene | | AF109907 | | | | 1 | | | |
| S-adenosylmethionine decarboxylase 1 (AMD1) | 3 | M88003 | + | + | + | + | | + | |
| SB classII histocompatibility antigen alpha-chain | 5 | M27487 | + | + | + | + | | + | |
| SC35-interacting protein 1 (SRRP129) | 5 | AF030234 | + | + | + | + | + | + | |
| scaffold attachment factor B (SAFB) | 1 | U72355 | + | + | + | + | | + | |
| scaffold attachment factor B (SAFB) (non-exact 78%) scRNA molecule. | 1 | U72355 | | | | _ | | | |
| transcribed from Alu repeat | ı | L13713 | | | ŀ | | | l | |
| SEC14 (S. cerevisiae)-like (SEC14L) | 4 | D67029 | | + | + | + | | + | |
| SEC23-like protein B (SEC23B) | 2 | X97065 | + | + | + | + | | + | |
| SEC63 (SEC63) | 1 | AF100141 | | + | + | | | + | |
| secreted protein, acidic, cysteine-rich (osteonectin) (SPARC) | 7 | M25746 | • | + | + | + | + | + | high in bone marrow stroma |
| secretory carrier membrane protein 1 (SCAMP1) | 1 | AF038966 | | * | | + | | | |
| secretory carrier membrane protein 2 (SCAMP2) | 1 | AF005038 | + | + | + | + | + | + | |
| secretory carrier membrane protein 3 (SCAMP3) | 1 | AF005039 | | | | | | | |
| secretory granule proteoglycan core (clones lambda-PG[6,7,8]) | 1 | M33649 | | | | | | | |
| selectin L (lymphocyte adhesion molecule 1) (SELL) | 43 | X17519 | + | | | + | | + | |
| selectin P ligand (SELPLG) | 13 | U02297 | + | + | | | <u> </u> | | |
| sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D (SEMA4D) | 2 | U60800 | | + | | + | | + | |
| Ser/Arg-related nuclear matrix protein (plenty of prolines 101-like) (SRM160) | 4 | AF048977 | | + | + | + | + | + | |
| serine palmitoyltransferase subunit I (SPTI) | 1 | Y08685 | | + | + | + | | + | |
| serine palmitoyltransferase. | 1 | AB011098 | + | + | + | + | | + | |

| WO 00/40/49 | | | _ | | | | | | |
|---|-----|------------|---------|---|----------|----|-------------|---|---------------------------------------|
| serine protease | 1 | J02907 | | | | | | | |
| serine protease inhibitor, | 1 | U78095 | + | + | + | + | | + | · · · · · · · · · · · · · · · · · · · |
| Kunitz type, 2 (SPINT2) | | | | | | | L | | |
| serine/threonine kinase 10 (STK10) | 1 | AB015718 | + | + | + | + | | + | |
| serine/threonine kinase 19 (STK19) | 1 | L26260 | + | + | + | + | | | |
| serine/threonine kinase 4 | 1 | U18297 | | + | | | | + | |
| (STK4) serine/threonine protein | 1 | X66358 | | + | + | + | | + | |
| kinase KKIALRE (KKIALRE) | | | | | | | | | |
| serine/threonine protein- kinase (NIK) | 1 | Y10256 | | + | + | + | | | |
| SERINE/THREONINE- PROTEIN KINASE RECEPTOR R3 | 1 | P37023 | | | | | | | |
| PRECURSOR (SKR3) serologically defined colon cancer antigen 16 (NY-CO- | 2 | AF039694 | | | <u> </u> | | | | |
| 16) | - 1 | AF039698 | В. Т | | + | | + | | |
| serologically defined colon cancer antigen 33 (SDCCAG33) | | | | | | | | | |
| serologically defined colon cancer antigen 33 (SDCCAG33) (low score) | 1 | AF039698 | | | | | | | |
| serologically defined colon cancer antigen 33 (SDCCAG33) (low score) | 1 | AF039698 | | | | | | | |
| serum deprivation response | 1 | AF085481.1 | | | | | | | |
| (phosphatidylserine-binding protein) (SDPR) (=S67386) | | | | | | | | | |
| serum/glucocorticoid regulated kinase (SGK) | 2 | Y10032 | + | + | + | + | | + | |
| SET domain, bifurcated 1 (SETDB1) | 2 | D31891 | + | + | + | | | + | |
| SH2 domain protein 1A, Duncan's disease lymphoproliferative syndrome) (SH2D1A) | 1 | AF073019 | Τ | | | | | + | |
| SH3 binding protein (SAB) | 2 | AB005047 | + | + | + | + | | + | |
| SH3 domain protein 1B | 4 | U61167 | + | | | + | | + | |
| (SH3D1B) SH3BGR PROTEIN (=21- GLUTAMIC ACID-RICH PROTEIN:21-GARP) (non- | 1 | P55822 | | | | | | | |
| exact 82%aa) SH3-binding domain | 1 | AF042081 | + | + | + | + | | + | |
| glutamic acid-rich protein like (SH3BGRL) | · . | | | | | Ĺ. | | | |
| SH3-domain GRB2-like 1 (SH3GL1) | 1 | U65999 | + | + | + | + | | + | |
| SHC (Src homology 2 domain-containing) transforming protein 1 (SHC1) | 2 | X68148 | | + | + | + | | + | |
| siah binding protein 1 (SiahBP1) | 2 | U51586 | | + | + | + | | + | |
| siah binding protein 1 (SiahBP1) (non-exact, 69%) | 1 | U51586 | | | | | | | |
| Sialomucin CD164 (CD164) | 9 | D14043 | | | | | | | |
| sialophorin (gpL115, leukosialin, CD43) (SNP) | 2 | J04536 | | | | | | | |
| sialyftransferase (STHM) | 1 | U14550 | | 1 | + | + | | + | |
| sialyttransferase 1 (beta- galactoside alpha-2,6- sialytransferase) (SIAT1) | 2 | X17247 | + | + | + | + | + | + | |

| sialyltransferase 4A (beta- galactosidase alpha-2,3- sialytransferase) (SIAT4A) | 1 | AF059321 | В | + | + | | + | + | |
|---|-----|------------|----------|-----|---|---|---|---|--------------|
| sialytransferase 8 (alpha- 2, 8-polysialytransferase) D (SIAT8D) | 1 | L41680 | | + | | | | | |
| signal peptidase 25kDa | 1 | L38950 | | † † | | | | | <u> </u> |
| signal recognition particle 14kD (homologous Alu RNA-binding protein) (SRP14) | 1 | X73459 | + | + | + | + | + | + | |
| signal recognition particle 54kD (SRP54) | 1 | U51920 | | | + | + | | + | |
| signal recognition particle 9kD (SRP9) | 2 | U20998 | | + | + | + | + | + | |
| signal recognition particle receptor ('docking protein') SRPR | 5 | X06272 | | | | | | | |
| signal regulatory protein, beta, 1 (SIRP-BETA-1) | 5 | Y10376 | | + | | | | + | |
| signal sequence receptor, alpha (translocon- associated protein alpha) (SSR1) | 2 | 212830 | | | | + | | + | |
| signal sequence receptor, beta (translocon- associated protein beta) (SSR2) | 2 | X74104 | + | + | + | + | | + | |
| signal fransducer and activator of transcription (STAT5A) | 4 | L41142 | + | + | + | + | + | + | |
| signal transducer and activator of transcription 2, 113KD (STAT2) | 1 | U18671 | | | | | | + | * |
| signal transducer and activator of transcription 3 (acute-phase response factor) (STAT3) | 3 | L29277 | | | | , | | | |
| signal transducer and activator of transcription 5A (STAT5A) | 2 | U48730 | + | + | + | + | + | + | |
| signal transducing adaptor molecule (SH3 domain and ITAM motif) 1 (STAM) | 1 | U43899 | | | | | | | |
| silencing mediator of retinoid and thyroid hormone action (SMRT) | 1 | U37146 | | | | | | | |
| similar to beta-transducin superfamily proteins (SAZD) | 1 | U02609 | + | + | + | | | + | |
| similar to S. cerevisiae SSM4 (TEB4) | 1 | AB011169 | | + | + | + | | + | |
| similar to yeast pre-mRNA splicing factors, Prp1/Zer1 and Prp6 | 1 | AF026031 | + | + | + | + | | + | |
| SIT protein | 1 - | AJ010059.1 | | | | | | | |
| Sjogren syndrome antigen A1 (52kD, ribonucleoprotein | 2 | M62800 | | | | | + | | |
| autoantigen SS-A/Ro) (SSA1) Sjogren syndrome antigen | 1 | M62800 | | | | | | ļ | |
| A1 (52kD, ribonucleoprotein autoantigen SS-A/Ro) (SSA1) (non-exact 63%) | • | 14102000 | | | | | | | |
| (match to zinc finger) SKAP55 homologue (SKAP-HOM) | 1 | AJ004886 | <u> </u> | + | + | + | | + | |
| skb1 (S. pombe) homolog (SKB1) | 2 | AF015913 | + | + | + | + | | + | |

| WO 00/40/49 | | | | | | | | | |
|--|----------|----------|----------------|---|--|---|---|----------|------------------------|
| skeletal muscle abundant protein | 1 | X87613 | + | + | + | + | | + | |
| SMA3 (SMA3) | 1 | X83300 | + | + | | + | | + | |
| small acidic protein | 3 | U51678 | + | + | + | + | | + | |
| small EDRK-rich factor 2 (SERF2) | 2 | Y10351 | + | + | + | + | + | + | high in fetal lung |
| small inducible cytokine A5 (RANTES) (SCYA5) | 2 | M21121 | + | + | + | + | + | + | high in many libraries |
| small inducible cytokine subfamily C, member 2 (SCYC2) | 1 | D63789 | | | | | | | |
| small nuclear ribonucleoprotein polypeptide B" (SNRPB2) | 2 | M15841 | | + | + | + | | + | |
| small nuclear ribonucleoprotein polypeptide N (SNRPN) | 4 | J04615 | + | + | + | + | + | + | |
| small nuclear ribonucleoprotein polypeptides B and B1 (SNRPB) | 2 | J04564 | + | + | + | + | | + | |
| small nuclear RNA activating complex, polypeptide 5, 19kD (SNAPC5) | 1 | AF093593 | + | + | + | + | | + | |
| smallest subunit of ubiquinol-cytochrome c ireductase | 1 | D55636 | + | + | . + | + | + | + | high in fetal lung |
| SMC (mouse) homolog, X | 1 | L25270 | + | + | + | + | | + | |
| chromosome (SMCX) SMT3B protein (2) | | X99585 | + | + | + | + | + | + | |
| SNARE protein (YKT6) | 1 | U95735 | _ | - | - | | | - | |
| (low match) | 1 | U20428 | | | - | - | ├ | \vdash | |
| SNC73 protein (SNC73) | 2 | J00220 | + | + | | + | + | + | high in many libraries |
| solute carrier family 1 | | U53347 | | + | | + | ├ | + | |
| (neutral amino acid transporter), member 5 (SLC1A5) | <u>-</u> | 030047 | | | | | | | |
| Solute carner family 11 (proton-coupled divalent metal ion transporters), member 1 (SLC11A1) | 7 | D50403 | + | | | | | | |
| solute carrier family 17 (sodium phosphate), member 3 (SLC17A3) | 1 | U90545 | | | | + | | | |
| solute carrier family 19 (folate transporter), member 1 (SLC19A1) | 1 | U17566 | B, lymphoma | + | | | + | | |
| solute carrier family 2 (facilitated glucose transporter), member 1 (SLC2A1) | 1 | K03195 | + | + | + | + | + | + | |
| solute carrier family 23 (nucleobase transporters), member 2 (SLC23A2) | 3 | D87075 | | + | + | + | | + | |
| solute carrier family 25 (mitochondrial carrier; oxoglutarate carrier), member 11 (SLC25A11) | 1 | AF070548 | В, Т | + | + | | + | + | |
| solute carrier family 31 (copper transporters), member 2 (SLC31A2) | 3 | U83461 | | + | | + | | | |
| solute carrier family 4, anion exchanger, member 2 (erythrocyte membrane protein band 3-like 1) (SLC4A2) | 1 | X62137 | | + | + | | | + | |
| solute carrier family 4, sodium bicarbonate cotransporter, member 8 (SLC4A8) | 1 | AB018282 | | + | | | | | |

| WU 00/40/49 | | | | _ | | | | | |
|--|----|------------|------|---|---|---|---|---|--|
| solute carrier family 7 (cationic amino acid transporter, y+ system), | 2 | M80244 | T, W | + | + | | + | | |
| member 5 (SLC7A5) | | | | | | | | | |
| solute carner family 7 (cationic amino acid transporter, y+ system), member 6 (SLC7A6) | 3 | D87432 | + | + | + | | | + | |
| solute carrier family 7 (cationic amino acid transporter, y+ system), member 6 (SLC7A6) (non- exact 77%) | 1 | D87432 | | | | | | | |
| solute carrier family 9 (sodium/hydrogen exchanger), isoform 6 (SLC9A6) | 1 | AF030409 | | + | + | + | | + | |
| somatic cytochrome c (HCS) | 2 | M22877 | | | | | | | |
| SON DNA binding protein (SON) | 2 | X63753 | | + | + | + | | + | |
| son of sevenless (Drosophila) homolog 1 (SOS1) | 1 | L13858 | + | + | | + | | | |
| sorcin (SRI) | 1 | M32886 | | | | | | | |
| sortilin 1 (SORT1) | 2 | X98248 | | + | | + | | + | |
| sortilin-related receptor, L(DLR class) A repeats- containing (SORL1) | 6 | Y08110 | | | | | | | |
| sorting nexin 1 (SNX1) | 3 | U53225 | + | + | + | + | | + | |
| sorting nexin 2 (SNX2) | 2 | AF043453 | | | | | | | |
| sorting nexin 6 (SNX6) (=U83194.1 TRAF4- associated factor 2) | 1 | AF121856.1 | | | | | | | |
| Sp3 transcription factor (SP3) | 1 | X68560 | + | + | + | + | | + | |
| Sp3 transcription factor (SP3) | 4 | M97191 | + | + | + | + | | + | |
| special AT-rich sequence binding protein 1 (binds to nuclear matrix/scaffold- associating DNA's) (SATB1) | 1 | M97287 | | | | | | | |
| speckle-type POZ protein (SPOP) | 4 | AJ000644 | | | | | | | |
| speckle-type POZ protein (SPOP) (non-exact) | | AJ000644 | | | | | | | |
| spectrin SH3 domain binding protein 1 (SSH3BP1) | 6 | U87166 | + | + | + | + | | | |
| Spectrin, alpha, non- erythrocytic 1 (alpha-fodrin) (SPTAN1) | 2 | J05243 | | + | + | | | + | |
| spermidine/spermine N1- acetyltransferase (SAT) | 11 | M55580 | | | | | | | |
| spermidine/spermine N1- acetyltransferase (SAT) (non-exact, 84%) | 1 | U40369 | | | | | | | |
| spermine synthase (SMS) | 1 | AD001528 | + | + | + | + | | + | |
| SPF31 (SPF31) | 1 | AF083190 | + | + | + | + | | + | |
| sphingomyelin phosphodiesterase 1, acid lysosomal (acid sphingomyelinase) (SMPD1) | 1 | X52679 | | + | + | | + | | |
| SPINDLIN HOMOLOG (PROTEIN DXF34) | 1 | Q99865 | | | | | | | |
| spinocerebellar ataxia 1 (olivopontocerebellar ataxia 1, autosomal dominant, ataxin 1) (SCA1) | 3 | X79204 | В | + | | | + | | |

| WO 00/40749 | | | | | | | | | .1/CA00/00003 | |
|--|-------------|----------|-------------|----------------|----------|------------|---|-----|---------------|----------|
| spinocerebellar ataxia 2 (olivopontocerebellar ataxia 2, autosomal dominant, ataxin 2) (SCA2) | 1 | U70323 | В | | | | + | | | |
| spinocerebellar ataxia 7 (olivopontocerebellar atrophy with retinal | 2 | AJ000517 | | + | • | | | | <u> </u> | |
| degeneration) (SCA7) spliceosome associated | 3 | U41371 | | + | + | + | + | + | | \dashv |
| protein (SAP 145) splicing factor (CC1.3) | 2 | L10910 | + | + | + | + | + | + | | \dashv |
| (CC1.3) splicing factor SRp40-1 | 7 | U30826 | + | + | + | + | + | + | | \dashv |
| (ŚRp4Ŏ) splicing factor, arginine/serine-rich 11 | 3 | M74002 | В | + | + | | + | + | | |
| (SFRS11) splicing factor, | 4 | L41887 | | + | + | + | | + | | |
| arginine/serine-rich 7 (35kD) (SFRS7) Src-like adapter protein | | U30473 | | | | | | | | |
| (non-exact, 76%aa) | | | | + | + | + | | + | | |
| Src-like-adapter (SLA) | 6 | D89077 | | ļ , | <u> </u> | ļ <u>,</u> | | ļ , | · | |
| Src-like-adapter (SLA) (low match) | 1 | D89077 | | | | | | | | |
| Src-like-adapter (SLA) (low score) | 1 | U44403 | | | | | | | | |
| stannin (SNN) | 2 | AF030196 | + | + | + | + | | + | | |
| STAT induced STAT inhibitor 3 (SSI-3) | 1 | AB004904 | | | | + | | | | |
| STE20-like kinase 3 (MST-3) | 2 | AF024636 | + | + | + | + | | + | | |
| step II splicing factor SLU7 (SLU7) | 1 | AF101074 | | + | | + | + | + | | |
| steroid sulfatase | 1 | M17591 | | | | | L | ' | | |
| steroid sulfatase (microsomal), arylsulfatase C, isozyme S (STS) | | J04964 | | + | + | + | | | | |
| sterol carrier protein 2 (SCP2) | 1 | M55421 | | + | + | + | + | + | | |
| sterol O-acyltransferase (acyl-Coenzyme A: cholesterol acyltransferase) 1 (SOAT1) | 1 | AF059202 | | | | | + | | | |
| stimulated trans-acting factor (50 kDa) (STAF50) | 6 | X82200 | + | + | | + | | | | |
| Striatin, calmodulin-binding protein (STRN) (low match, 71%aa) | 1 | U17989 | | | | | | | | |
| Stromal antigen 2 (STAG2) | 2 | Z75331 | | | + | + | + | + | | \neg |
| stromal interaction molecule 1 (STIM1) | 3 | U52426 | + | + | + | + | | + | - | |
| structure specific recognition protein 1 (SSRP1) | 1 | M86737 | | + | + | + | | + | | |
| succinate dehydrogenase complex, subunit A, flavoprotein (Fp) (SDHA) | 5 | L21936 | | | + | | | | | |
| succinate dehydrogenase complex, subunit B, iron sulfur (Ip) (SDHB) | 1 | U17248 | + | + | + | + | | + | | |
| succinate dehydrogenase complex, subunit C, integral membrane protein, | 1 | U57877 | + | + | + | + | | + | | |
| 15kD (SDHC) succinate dehydrogenase complex, subunit D, integral membrane protein | 3 | AB006202 | | + | + | | + | | | |
| (SDHD) succinate-CoA ligase, GDP-forming, beta subunit | 1 | AF058954 | | + | + | + | + | + | | \dashv |
| (SUCLG2) | | L | | | <u> </u> | | | | | |

PCT/CA00/00005

| WO 00/40/49 | | | | | | | | | C1/CA00/00003 |
|--|----------|----------|----------|--|----------|---------------------------------------|----------|----------|---------------|
| succinyl CoA synthetase | 1 | Z68204 | 1 | | | | | | |
| sudD (suppressor of bimD6, Aspergillus | 2 | AF013591 | | + | | | + | + | |
| nidulans) homolog (SUDD) | | 1.40000 | | | <u> </u> | | <u> </u> | <u> </u> | |
| sulfotransferase family 1A, phenol-preferring, member 1 (SULT1A1) | 1 | L19999 | | + | | | + | + | |
| sulfotransferase family 1A, phenol-preferring, member 3 (SULT1A3) (non-exact 67%) | 1 | U37686 | | | | | | | |
| superoxide dismutase 1, soluble (amyotrophic lateral sclerosis 1 (adult)) (SOD1) | 4 | X02317 | | + | + | | + | + | |
| superoxide dismutase 2, mitochondrial (SOD2) | 5 | Y00985 | | + | + | + | + | + | |
| supervillin (SVIL) | 2 | AF051851 | | | + | + | 1 | + | |
| suppression of | 2 | U15131 | <u> </u> | + | | + | | + | |
| tumorigenicity 5 (ST5) suppression of | 1 | U15779 | | | | ـــــــــــــــــــــــــــــــــــــ | | | |
| tumorigenicity 5 (ST5) | 1 | 013/79 | | | | | | | |
| suppressor of K+ transport defect 1 (SKD1) | 1 | AF038960 | | | + | + | | | |
| suppressor of Ty (S.cerevisiae) 3 homolog (SUPT3H) | 1 | AF064804 | + | + | + | + | | + | |
| suppressor of Ty (S.cerevisiae) 4 homolog 1 (SUPT4H1) | 2 | U38817 | + | + | + | + | | + | |
| suppressor of Ty (S.cerevisiae) 5 homolog (SUPT5H) | 2 | U56402 | | + | | | | + | |
| suppressor of Ty (S.cerevisiae) 6 homolog (SUPT6H) | 2 | U46691 | + | + | + | + | + | + | |
| suppressor of variegation 3-9 (Drosophila) homolog 1 (SUV39H1) | 1 | AF019968 | | + | + | + | | | |
| survival of motor neuron 1, telomeric (SMN1) | 1 | U18423 | | | | | | | |
| SWI/SNF related, matrix | 1 | M88163 | | | + | + | | + | |
| associated, actin dependent regulator of chromatin, subfamily a, member 1 (SMARCA1) (non-exact, 75%) | | | | | | | | | |
| SWI/SNF related, matrix | 2 | D26155 | | + | | | | | |
| associated, actin dependent regulator of | | ļ | | | | | | | |
| chromatin, subfamily a, member 2 (SMARCA2) | | | | | | | | | |
| SWI/SNF related, matrix associated, actin | 1 | D26156 | + | + | + | + | + | + | |
| dependent regulator of chromatin, subfamily a, member 4 (SMARCA4) | | | | | | | | | |
| SWI/SNF related, matrix | 4 | U66616 | + | + | + | + | + | + | |
| associated, actin dependent regulator of chromatin, subfamily c, | | | | | | | | | |
| member 2 (SMARCC2) SWI/SNF related, matrix | 2 | AF035262 | B, W | + | + | | + | + | |
| associated, actin dependent regulator of chromatin, subfamily e, | | | | | | · | | | |
| member 1 (SMARCE1) synaptobrevin-like 1 | 1 | X95803 | | + | + | + | | + | |
| (\$YBL1) synaptosomal-associated | 2 | AJ011915 | | + | + | + | | + | |
| protein, 23kD (SNAP23) | | | | | | | | | |
| syndecan binding protein (syntenin) (SDCBP) | 15 | AF006636 | + | + | + | + | | + | |

| WO 00/40749 | | | | | | | | rc | CT/CA00/00005 |
|---|--------------|------------|---|----|---|---|----------|----|---------------|
| synovial sarcoma, translocated to X chromosome (SSXT) | 2 | X79201 | | + | | | | | |
| syntaxin 16 | | AF038897 | | 11 | | | 1 | | |
| syntaxin 3A (STX3A) | 2 | U32315 | | + | | + | | + | |
| syntaxin 6 (STX6) | | AJ002078.1 | | 11 | | | | | |
| SYNTAXIN BINDING PROTEIN 3 (UNC-18 | 1 | O00186 | | | | | | | |
| HOMOLOG 3) (UNC-18C) syntaxin-16C | | AF008937 | | + | | | | | |
| SYT interacting protein | | AF080561 | | ++ | + | + | | + | |
| (SIP) | - 4 | M88282 | | ļ | | + | \dashv | | |
| late expression (TACTILE) | 2 | X58744 | | - | | | | _ | |
| T cell receptor V alpha gene segment V-alpha-7 (clone IGRa11) | | | | | | | | | |
| T cell receptor V alpha gene segment V-alpha-w27 | 1 | X58740 | | | | | | | |
| T3 receptor-associating cofactor-1 | 5 | 583390 | + | + | + | + | + | + | |
| tafazzin (cardiomyopathy, dilated 3A (X-linked); endocardial fibroelastosis 2; Barth syndrome) (TAZ) | 1 | X92763 | + | + | | + | | + | |
| TAFII100 protein (non- exact 53%) | 1 | U80191 | | | | | | | |
| tankyrase, TRF1- interacting ankyrin-related ADP-ribose polymerase (TNKS) | 1 | AF082556 | - | + | + | + | | + | |
| TAP1, TAP2, LMP2, LMP7 | 1 | X66401 | | | | | | | |
| TAR DNA-binding protein- | 6 | U23731 | + | + | + | + | | + | |
| Tat interactive protein (60kD) (TIP60) | 2 | U40989 | + | + | + | + | | + | |
| TATA box binding protein (TBP)-associated factor, RNA polymerase II, C1, 130kD (TAF2C1) (non- exact, 55%) | 1 | O00268 | | | | | | | |
| TATA box binding protein (TBP)-associated factor, RNA polymerase II, F, 55kD (TAF2F) | 4 | X97999 | | + | + | + | + | + | |
| TATA box binding protein (TBP)-associated factor, RNA polymerase II, G, 32kD (TAF2G) | 2 | U21858 | | + | + | + | + | + | |
| TATA box binding protein (TBP)-associated factor, RNA polymerase II, I, 28kD (TAF2I) | 1 | D63705 | + | + | + | + | | + | |
| Tax1 (human T-cell leukemia virus type I) binding protein 1 (TAX1BP1) | 1 | U33821 | | + | + | + | + | + | |
| T-box 2 (TBX2) (non-exact 77%) | 1 | U28049 | | | + | + | | + | |
| TBP-associated factor 172 (TAF-172) | 1 | AJ001017 | | + | | + | | + | |
| T-cell death-associated gene 8 (TDAG8) | 1 | U95218 | | | | + | | | |
| T-cell leukemia/lymphoma 1A (TCL1A) | 1 | X82240 | + | | | | | | |
| T-cell leukemia/lymphoma 1A (TCL1A) (low match) | 1 | X82240 | | | | | | | |
| T-cell receptor (delta D2- J1-region) (clone K3B) | 1 | M22197 | | | | | | | |

| WO 00/40/49 | | | | | | | | | |
|---|--------------|--------------------|---------------|--------------|----------|----------|-----|----------|---------------------|
| T-cell receptor (V beta 5.1, J beta 1.5, C beta 1) (low match) | 1 | M97705 | | | | | | | |
| T-cell receptor alpha delta (=M94081) | 2 | AE000662 | | | | | | | |
| T-cell receptor alpha enhancer-binding protein, short form (=X58636 Mouse LEF1 lymphoid enhancer binding factor 1 | 1 | B39625 | | | | | | | |
| (=D16503)) T-cell receptor delta gene | 1 | M22197 | | | | | | | |
| D2-J1-region, clone K3B T-cell receptor germline beta chain gene V-region | 1 | M11955 | | | | | | | |
| (V) V-beta-MT1-1 T-cell receptor germline | 1 | M14159 | + | | | | | | only in blood |
| beta-chain gene J2.1 exon T-cell receptor germline delta-chain D-J region | 2 | M22152 | | | | | | _ | |
| T-cell receptor interacting molecule (TRIM) protein | 2 | AJ224878 | | | | | | + | |
| T-cell receptor rearranged delta-chain, V-region (V-idelta 3-J) | 1 | M21784 | | | | | | | |
| T-cell receptor, alpha (V,D,J,C) (TCRA) | 3 | AE000660 | + | + | + | + | | + | |
| T-cell receptor, beta cluster (TCRB) | 3 | L34740 | + | + | + | + | + | + | high in pancreas |
| T-cell receptor, delta (V.D.J.C) (TCRD) | 2 | X73617 | | | + | + | | + | |
| T-cell, immune regulator 1 (TCIRG1) | 3 | U45285 | | | | | | | only found in tumor |
| TCF-1 mRNA for T cell factor 1 | 1 | X59870 | | | | | | | |
| TCF-1 mRNA for T cell factor 1 (splice form B) (low match) | 1 | X59870 | | | | | | | |
| T-COMPLEX PROTEIN 1, ETA SUBUNIT (TCP-1- ETA) (CCT-ETA) (HIV-1 NEF INTERACTING PROTEIN) | 1 | Q99832 | | | | | | | ÷ |
| T-COMPLEX PROTEIN 1, THETA SUBUNIT (TCP-1- THETA) (CCT-THETA) (KIAA0002) | 1 | P50990 | | | | | | | |
| TCR eta =1 cell receptor(eta-exon) | 1 | S94421 | | | | | | | |
| TCR V Beta 13.2 | 1 | X75419 | | | | | | | |
| TERA | 1 20 | AC004472 X75861 | - | + | + | + | + | + | |
| transcript (TEGT) | 33 | | | + | + | <u> </u> | ļ . | + | |
| tetracycline transporter-like protein (TETRAN) | 2 | L11669 | + | <u> </u> | + | + | | + | |
| tetratricopeptide repeat domain 1 (TTC1) | | U46570 U46571 | | + | | + | | <u> </u> | |
| tetratricopeptide repeat domain 2 (TTC2) | 1 | D84296 | + | + | + | + | _ | ļ . | |
| tetratricopeptide repeat domain 3 (TTC3) | 1 | D86970 | ·+ | <u>'</u> | + | + | | + | |
| TGFB1-induced anti- apoptotic factor 1 (TIAF1) thioredoxin reductase 1 | 3 | S79851 | · | + | + | + | _ | + | |
| (TXNRD1) THIOREDOXIN- | | P30048 | | | <u> </u> | <u> </u> | | - | |
| DEPENDENT PEROXIDE REDUCTASE PRECURSOR, mitochondrial (ANTI- OXIDANT PROTEIN 1) | , | . 300-10 | | | | | | | |
| (AOP-1) | | | | | | <u> </u> | L | | <u></u> |

PCT/CA00/00005

| WO 00/40/49 | | | | | | | | | |
|--|----|------------|---|---|---|---|---|---|------------------|
| threonyl-tRNA synthetase (TARS) | 1 | M63180 | | + | + | + | | + | |
| thrombin inhibitor | 1 | Z22658 | | | | | | | |
| thrombospondin 1 (THBS1) | 2 | X04665 | | + | + | + | + | + | |
| thromboxane A synthase 1 (platelet, cytochrome P450, subfamily V) (TBXAZ1) | 1 | M80647 | | + | | + | + | + | |
| thymidine kinase 2, mitochondrial (TK2) | 2 | X76104 | | + | + | | + | | |
| thymidylate kinase (CDC8) | 1 | L16991 | | + | + | + | | + | |
| thymine-DNA glycosylase (TDG) | 2 | U51166 | + | + | + | + | | + | |
| Thymosin, beta 10 (TMSB10) | 2 | M20259 | + | + | + | + | + | + | |
| thymosin, beta 4, X chromosome (TMSB4X) | 29 | M17733 | | + | + | + | | + | |
| thyroid autoantigen 70kD (Ku antigen) (G22P1) | 7 | J04611 | | | | | | | |
| thyroid hormone receptor coactivating protein (SMAP) | 1 | AF016270 | | + | | + | | + | |
| thyroid hormone receptor interactor 7 (TRIP7) | 2 | L40357 | | + | + | + | | + | |
| thyroid hormone receptor interactor 8r (TRIP8) | 4 | L40411 | | + | | | | | |
| thyroid hormone receptor- associated protein, 230 kDa subunit (TRAP230) | 1 | D83783 | | | | | _ | _ | |
| thyroid receptor interacting protein 15 (TRIP15) | 2 | L40388 | + | + | + | + | | | |
| TI-227H | 1 | D50525 | | | | | | | |
| TIA1 cytotoxic granule- associated RNA-binding protein (TIA1) | 1 | M77142 | | + | + | + | | + | |
| tissue inhibitor of metalloproteinase 1 (erythroid potentiating activity, collagenase inhibitor) (TIMP1) | 1 | X02598 | 4 | + | + | + | + | + | |
| tissue inhibitor of metalloproteinase 2 | 1 | M32304 | + | + | + | + | | + | high in placenta |
| tissue specific transplantation antigen P35B (TSTA3) | 1 | U58766 | + | + | + | + | | + | |
| titin (TTN) | 1 | X64697 | + | + | + | + | | + | high in muscle |
| TNF receptor-associated factor 2 (TRAF2) | 1 | U12597 | | + | + | + | | + | |
| TNF receptor-associated factor 3 (TRAF3) | 1 | AF110908.1 | | + | | | | | |
| TNF receptor-associated factor 6 (TRAF6) (low match) | 1 | 078798 | | | | | | | |
| toll-like receptor 1 (TLR1) | 1 | U88540 | | | | + | | | |
| toll-like receptor 2 (TLR2) | 1 | U88878 | + | + | | + | | + | |
| toll-like receptor 4 (TLR4) | 1 | U88880 | | + | | | + | | |
| toll-like receptor 5 (TILR5) | 1 | AF051151 | | + | | + | | | |
| topoisomerase (DNA) I (TOP1) | 1 | J03250 | | + | + | + | | | |
| topoisomerase (DNA) II beta (180kD) (TOP2B) | 2 | X68060 | + | + | + | + | | + | |
| topoisomerase (DNA) III beta (TOP3B) | 3 | D87012 | + | | | | | | |
| TR3beta | 1 | D85245 | | + | | | | | |
| TRAF family member- associated NF-kB activator (TANK) | 3 | U63830 | + | + | + | + | + | + | |
| TRANSALDOLASE | 1 | P37837 | | | | | | | |
| transaldolase 1 (TALDO1) | 4 | L19437 | | + | + | + | + | + | |

| transaldolase-related protein | 1 | AF010398 | | | | | | | ! |
|---|---|----------|---|---|---|----------|----------|---|------|
| transcobalamin II (TCII) | 1 | AF047576 | | 1 | | | | | |
| transcription elongation | 2 | Z47087 | + | + | + | + | | + | |
| factor B (SIII), polypeptide 1-like (TCEB1L) | | | | | | | | | |
| transcription elongation factor B (SIII), polypeptide 3 (110kD, elongin A) (TCEB3) | 1 | L47345 | + | + | + | + | + | + | |
| (HTF4, helix-loop-helix transcription factors 4) | 1 | M83233 | + | + | + | + | | + | , |
| transcription factor 17 (TCF17) | 2 | D89928 | | + | | + | | | |
| transcription factor 4 (TCR4) | 2 | X52079 | | + | + | + | | + | |
| transcription factor 6-like 1 (mitochondrial transcription factor 1-like) (TCF6L1) | 2 | M62810 | + | + | + | + | | | |
| transcription factor 7-like 2 (T-cell specific, HMG-box) (TCF7L2) | 1 | Y11306 | | + | + | + | | + | |
| transcription factor binding to IGHM enhancer 3 (TFE3) | 1 | X96717 | + | + | + | + | + | + | |
| transcription factor IL-4 Stat | 7 | AF067575 | + | + | + | | | _ | |
| transcription factor IL-4 Stat (low match) | 1 | U16031 | | | | ļ | | | |
| transcription factor ISGF-3 (=M97936) | 4 | M97935 | | | | | | | |
| transcription factor REST | 1 | A56138 | | | | <u> </u> | <u> </u> | | |
| transcription factor TFIID | | Z22828 | | | | <u> </u> | <u> </u> | | |
| transcriptional adaptor 2 (ADA2, yeast, homolog)- like (TADA2L) | 1 | AF064094 | | | | | | | |
| transcriptional intermediary factor 1 (TIF1) (non-exact 72%) | 1 | AF009353 | | | | | | | |
| transducin (beta)-like 1 (TBL1) | 1 | Y12781 | + | + | + | + | | + | |
| transducin-like enhancer of split 3, homolog of Drosophila E(sp1) (TLE3) | 1 | M99438 | + | + | | | | | |
| Transformation/transcription domain-associated protein (TRRAP) | 1 | AF076974 | + | + | + | + | | + | |
| transformation-sensitive, similar to Saccharomyces cerevisiae STI1 (STI1L) | 2 | M86752 | | + | + | + | | + | |
| transforming growth factor beta-activated kinase 1 (TAK1) (non-exact 78%) | 1 | AB009356 | | | | | | | |
| transforming growth factor beta-stimulated protein TSC-22 (TSC22) | 3 | AJ222700 | + | + | + | + | | + | |
| transforming growth factor, beta receptor III (betaglycan, 300kD) | 1 | L07594 | | + | + | + | | + | |
| (TGFBR3) transforming growth factor, beta-induced, 68kD (TGFBI) | 2 | 4507466 | + | + | + | + | + | + | |
| TRANSFORMING GROWTH FACTOR-BETA INDUCED PROTEIN IG-H3 PRECURSOR (BETA IG-H3) | 2 | Q15582 | | | | | | | |
| transforming, acidic coiled- coil containing protein 1 (TACC1) (non-exact 70%) | 1 | AF049910 | | | | | | | |

| WO 00/40/49 | | | | | | | | | |
|---|----|----------|---|---|---|---|---|---|------------------------|
| transgelin 2 (TAGLN2) | 14 | D21261 | + | + | + | + | + | + | |
| transgelin 2 (TAGLN2) (non-exact) | 1 | D21261 | | | | | | | |
| trans-Golgi network protein (46, 48, 51kD isoforms) (TGN51) | 2 | AF029316 | | + | | + | | | |
| transient receptor potential channel 1 (TRPC1) | 1 | X89066 | | + | + | + | | + | |
| transketolase (Wernicke- Korsakoff syndrome) (TKT) | 7 | L12711 | | + | + | + | | + | |
| translation factor sui1 homolog (GC20) | 1 | AF064607 | | + | + | + | + | + | |
| translin (TSN) | 3 | X78627 | + | + | + | + | | + | |
| translin-associated factor X (TSNAX) | 1 | X95073 | | + | + | + | | + | |
| transmembrane | 1 | U79725 | | | | | | | |
| glycoprotein (A33) transmembrane protein (63kD), endoplasmic reticulum/Golgi intermediate compartment (P63) | 1 | X69910 | + | + | + | + | | + | |
| transmembrane protein 1 (TMEM2) | 1 | AB001523 | | + | | + | | + | |
| TRANSMEMBRANE PROTEIN SEX PRECURSOR (non-exact 65%) | 1 | P51805 | | | | | | | |
| transmembrane trafficking protein (TMP21) | 2 | X97442 | + | + | + | + | + | + | |
| transporter 1, ABC (ATP binding cassette) (TAP1) | 3 | L21208 | + | + | + | + | | + | |
| Treacher Collins- Franceschetti syndrome 1 (TCOF1) | 2 | U40847 | + | + | + | + | | + | high in many libraries |
| triosephosphate isomerase 1 (TPI1) | 2 | X69723 | + | + | + | + | + | + | |
| tropomyosin | 2 | X04201 | | + | + | + | | + | |
| tropomyosin 4 (TPM4) | 2 | X05276 | + | + | + | + | | + | |
| TRPM-2 protein | 2 | M63376 | | | | | | | |
| tryptase I precursor (non- exact 64%)(=P20231) | 1 | A35863 | | | | | | | |
| tryptophan rich basic protein (WRB) | 1 | Y12478 | | | | | | | |
| tryptophanyl-tRNA synthetase (WARS) | 1 | X59892 | + | + | + | + | + | + | |
| Ts translation elongation factor, mitochondrial (TSFM) | 1 | L37936 | + | + | | + | | + | |
| ttopoisomerase (DNA) II beta (180kD) | 1 | Z15115 | | + | + | | | + | |
| Tu translation elongation factor, mitochondrial (TUFM) | 4 | L38995 | | | | | | | |
| tuberous sclerosis 1 (TSC1) | 1 | AF013168 | | + | + | + | | + | |
| tuberous sclerosis 2 (TSC2) | 1 | X75621 | | + | + | + | | + | |
| tubulin, alpha 1 (testis specific) (TUBA1) | 1 | X06956 | | + | | | + | | |
| tubulin, alpha, ubiquitous (K-ALPHA-1) | 11 | K00558 | + | + | + | + | + | + | high in many libraries |
| tubulin, alpha, ubiquitous (K-ALPHA-1) (low match) | 1 | K00558 | | | | | | | |
| tubulin-specific chaperone (TBCC) | 1 | U61234 | | + | + | + | | + | |
| tumor necrosis factor (ligand) superfamily, member 10 (TNFSF10) | 7 | U37518 | | + | + | + | | + | |

| WO 00/40/49 | | | | | | | | • • | |
|--|-------------|----------|--------|----------|---|---|---|----------|---|
| tumor necrosis factor (ligand) superfamily, member 13 (TNFSF13) | 1 | AF046888 | + | + | | + | | + | |
| tumor necrosis factor (ligand) superfamily, | 1 | AF036581 | | | | | | | |
| member 14 (TNFSF14) tumor necrosis factor (ligand) superfamily. | 1 | D38122 | + | | | | | | Found only in library 386: T-cell lymphoma |
| member 6 (TNFSF6) tumor necrosis factor (ligand) superfamily, | | L09753 | В опіу | <u> </u> | | | | | |
| member 8 (TNFSF8) tumor necrosis factor alpha-inducible cellular | | AF061034 | | + | + | + | | + | |
| protein containing leucine zipper domains (FIP2) | 2 | M63928 | | + | | | + | <u> </u> | |
| Tumor necrosis factor receptor superfamily member 7 (TNFRSF7) | | | | | | | | | |
| tumor necrosis factor receptor superfamily, member 10b (TNFRSF10B) | 1 | AF016266 | | + | + | + | + | + | |
| tumor necrosis factor receptor superfamily, member 10c, decoy without an intracellular domain (TNFRSF10C) | 3 | AF012629 | | | | | + | | |
| tumor necrosis factor receptor superfamily, member 10d, decoy with truncated death domain (TNFRSF10D) (non-exact 84%) | 1 | AF023849 | | | | | | | found only in prostate |
| tumor necrosis factor receptor superfamily, member 12 (translocating chain-association membrane protein) (TNFRSF12) | | U94508 | + | + | + | + | | + | |
| tumor necrosis factor receptor superfamily, member 14 (herpesvirus entry mediator) (TNFRSF14) | 1 | U70321 | + | + | + | + | | + | |
| tumor necrosis factor receptor superfamily, member 1B (TNFRSF1B) | 5 | U52165 | + | + | + | + | | + | |
| tumor necrosis factor receptor superfamily, member 6 (TNFRSF6) | 1 | X63717 | B, W | | | | | + | |
| tumor necrosis factor receptor superfamily, member 7 (TNFRSF7) | 1 | M63928 | + | + | | | | | |
| tumor necrosis factor, alpha-induced protein 2 (TNFAIP2) | 8 | M92357 | | + | + | | + | | |
| tumor necrosis factor, alpha-induced protein 3 (TNFAIP3) | 2 | M59465 | | | | | | | |
| tumor protein 53-binding protein, 1 (TP53BP1) | 1 | AF078776 | | + | + | + | | + | |
| tumor protein p53 (Li- Fraumeni syndrome) (TP53) | 1 | M14695 | + | + | | | | + | |
| Tumor protein p53-binding protein (TP53BPL) | 1 | U82939 | + | | | + | | + | |
| tumor protein, translationally-controlled 1 (TPT1) | 35 | X16064 | | | | | | | |
| tumor protein, translationally-controlled 1 (TPT1) (low score) | 1 | X16064 | | | | | | | |
| tumor rejection antigen | | X15187 | | + | + | + | + | + | |

| WO 00/40/42 | | | | | | | | | |
|--|-------------|-----------|--------|----------|---|----------|----------|---|-------------------------------------|
| tumorous imaginal discs (Drosophila) homolog (TID1) | 2 | AF061749 | | + | | | | | |
| TXK tyrosine kinase (TXK) | 2 | L27071 | | T | | | | | |
| type II integral membrane protein (NKG2-E) | 1 | AJ001685 | | | | | + | | found only in fetal liver/spleen |
| TYRO protein tyrosine kinase binding protein | 3 | AF019562 | | | + | | | | |
| (TYROBP) tyrosine 3- | | X57346 | + | + | + | + | | + | high in ecnorm |
| monooxygenase/tryptopha n 5-monooxygenase activation protein, beta polypeptide (YWHAB) | , | 737340 | , _ | | · | | | | |
| tyrosine 3- monooxygenase/tryptopha n 5-monooxygenase activation protein, zeta polypeptide (YWHAZ) | 1 | M86400 | | | | | | | |
| tyrosine 3- | 1 | M86400 | | +1 | | | | | |
| monooxygenase/tryptopha n 5-monooxygenase activation protein, zeta polypeptide (YWHAZ) | | | | | | | | | |
| Tyrosine kinase 2 (TYK2) | 3 | X54637 | | + | + | + | | + | |
| TYROSINE-PROTEIN KINASE ZAP-70 (70 KD ZETA-ASSOCIATED PROTEIN) (SYK-RELATED TYROSINE KINASE) | 2 | P43403 | | | | | | | |
| tyrosyl-tRNA synthetase (YARS) | 1 | U89436 | + | + | + | + | | + | |
| U1 small nuclear RNA | 1 | M14387 | | | | | | | |
| U19H snoRNA (=M63485 R.norvegicus matrin 3) | 1 | AJ224166 | | | | | | | |
| U2(RNU2) small nuclear RNA auxillary factor 1 (non-standard symbol) (U2AF1) | 1 | M96982 | | + | + | + | | + | |
| U22 snoRNA host gene (UHG) | 2 | U40580 | | | | | | | |
| U4/U6-associated RNA splicing factor (HPRP3P) | 4 | AF016370 | | + | + | + | | + | |
| U49 small nuclear RNA | 1 | X96649 | | | | T | | | |
| U5 snRNP-specific protein (220 kD), ortholog of S. cerevisiae Prp8p (PRP8) | 1 | AB007510 | + , | + | + | + | | + | |
| U5 snRNP-specific protein, 116 kD (U5-116KD) | 4 | D21163 | + | + | + | + | | + | |
| U5 snRNP-specific protein, 200 kDa (DEXH RNA helicase family) (U5-200- KD) | 3 | 270200 | | | | | | | |
| Uba80 mRNA for ubiquitin | 4 | S79522 | + | + | + | + | + | + | high in ovary |
| ubiquinol-cytochrome c reductase (6.4kD) subunit (UQCR) | 1 | D55636 | + | + | + | + | + | + | high in fetal lung |
| UBIQUINOL- CYTOCHROME C REDUCTASE IRON- SULFUR SUBUNIT PRECURSOR (RIESKE IRON-SULFUR PROTEIN) (RISP) (low match) | 1 | P47985 | | | | | | | |
| ubiquitin A-52 residue | 2 | X56999 | - | | | | | | |
| ribosomal protein fusion product 1 (UBA52) | | AF00/1510 | | <u> </u> | + | _ | | + | |
| ubiquitin activating enzyme E1-like protein (GSA7) | 1 | AF094516 | | + | | <u> </u> | <u> </u> | | high in over |
| ubiquitin C (UBC) | 5 | AB009010 | | + | + | + | + | + | high in overy |

| WO 00/40/49 | | | | | | | | | |
|--|---|----------|---|---|---|---|----------|---|--|
| ubiquitin carboxyl-terminal esterase L3 (ubiquitin thiolesterase) (UCHL3) | 1 | M30496 | + | + | + | + | i | + | |
| ubiquitin fusion degradation 1-like (UFD1L) | 1 | U64444 | + | + | + | + | | + | |
| ubiquitin protein ligase E3A (human papilloma virus E6- associated protein, Angelman syndrome) (UBE3A) | 1 | U84404 | В | + | + | | | + | |
| ubiquitin specific protease 10 (USP10) | 4 | D80012 | + | + | + | + | | + | |
| ubiquitin specific protease 11 (USP11) | 1 | U44839 | + | + | + | + | + | + | |
| ubiquitin specific protease 15 (USP15) | 3 | AB011101 | + | + | + | + | | + | |
| ubiquitin specific protease 19 (USP19) | 1 | AB020698 | | + | | | | | |
| ubiquitin specific protease 4 (proto-oncogene) (USP4) | 1 | AF017305 | В | + | + | | + | + | |
| ubiquitin specific protease 4 (proto-oncogene) (USP4) (non-exact, 66%) | 1 | AF017306 | | | | | | | |
| ubiquitin specific protease 7 (herpes virus-associated) (USP7) | 1 | Z72499 | | + | + | + | | + | |
| ubiquitin specific protease 8 (USP8) | 5 | D29956 | | + | + | + | | + | |
| UBIQUITIN-ACTIVATING ENZYME E1 (A1S9 PROTEIN) (56%) | 1 | P22314 | | | | | | | |
| ubiquitin-activating enzyme E1 (A1S9T and BN75 temperature sensitivity complementing) (UBE1) | 1 | M58028 | + | + | + | + | | + | |
| ubiquitin-activating enzyme E1, like (UBE1L) | 1 | L34170 | + | + | | + | | + | |
| UBIQUITIN-BINDING PROTEIN P62; phosphotyrosine independent ligand for the Lck SH2 domain p62 (P62) | 7 | U41806 | | | + | | + | | |
| ubiquitin-conjugating enzyme E2 variant 1 (UBE2V1) | 2 | U49278 | + | + | + | + | + | + | |
| ubiquitin-conjugating enzyme E2 variant 2 (UBE2V2) | 1 | X98091 | | | | | | | |
| UBIQUITIN- CONJUGATING ENZYME E2-17 KD (UBIQUITIN- PROTEIN LIGASE) | 1 | Q16781 | | | | | | | |
| ubiquitin-conjugating enzyme E2B (RAD6 homolog) (UBE2B) | 1 | M74525 | + | + | + | + | | + | |
| ubiquitin-conjugating enzyme E2G 2 (homologous to yeast UBC7) (UBE2G2) | 1 | AF032456 | + | + | + | + | | + | |
| ubiquitin-conjugating enzyme E2H (homologous to yeast UBC8) (UBE2H) | 1 | Z29328 | + | + | + | + | | + | |
| ubiquitin-conjugating enzyme E2L 1 (UBE2L1) | 1 | X92962 | | + | + | | <u> </u> | + | |
| ubiquitin-conjugating enzyme E2L 3 (UBE2L3) | 3 | AJ000519 | | + | + | + | _ | + | |
| ubiquitin-conjugating | 4 | AF031141 | | + | + | + | + | + | |
| enzyme E2L 6 (UBE2L6) ubiquitin-like 1 (sentrin) | 2 | U61397 | + | + | + | + | | + | |

| 2 | X85019 | | | | | | | |
|----------|--|---|--|--|--|--|--------------------|--------------------|
| | | | | | | | | |
| 1 | X92689 | | | | | | | |
| | | | | | | | | |
| | 124804 | | | + | + | | + | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| <u> </u> | | | | | | | | |
| 1 | | | | | | | | |
| 1 | | | | | | | | |
| 2 | | | | | | | | |
| 1 | AF070626 | + | + | + | + | + | + | |
| 1 | AF040966 | | | | | | | |
| 1 | D79984 | + | + | + | + | + | + | |
| 74 | S73591 | + | + | + | + | | + | high in heart |
| 1 | S73591 | | | | | | | |
| 1 | S73591 | | | | | | | |
| 1 | 573591 | | | | | | | |
| 1 | X53461 | + | + | | + | | + | |
| 2 | X99050 | | + | + | + | | + | |
| 4 | X71490 | | + | + | + | + | + | |
| 1 | M63167 | + | + | + | + | | + | |
| 3 | AJ132100 | | 1 | | | | | |
| 3 | Z46389 | + | | + | + | | + | |
| 1 | M59834 | | 1 | | | | + | |
| | 576992 | + | + | | | | | |
| 1 | D10656 | W | + | + | | + | | |
| 1 | M29366 | | | | | | + | |
| 1 | P13611 | | | | | | | |
| 1 | M36196 | | + | + | + | | + | |
| | 2 3 1 1 1 2 1 1 1 1 2 4 1 3 3 1 1 1 | 1 X92689 2 L24804 3 U57053 1 U94592 1 U94592 1 AC002310 1 AF070542 2 Z70223 1 AF070626 1 AF040966 1 D79984 74 S73591 1 S73591 1 S73591 1 S73591 1 S73591 1 X53461 2 X99050 4 X71490 1 M63167 3 AJ132100 3 Z46389 1 M59834 1 S76992 1 D10656 1 M29366 | 1 X92689 2 L24804 3 U57053 1 U94592 1 U94592 1 AC002310 1 AF070542 2 Z70223 1 AF040966 1 D79984 + 74 S73591 + 1 S73591 1 S73591 1 S73591 1 X53461 + 2 X99050 4 X71490 1 M63167 + 3 AJ132100 3 Z46389 + 1 M59834 1 S76992 + 1 D10656 W | 1 X92689 2 L24804 + 3 U57053 1 U94592 1 U94592 1 AC002310 1 AF070542 2 Z70223 1 AF070626 + + 1 AF040966 1 D79984 + + 74 S73591 + + 1 S73591 1 S73591 1 S73591 1 X53461 + + 2 X99050 + 4 X71490 + 1 M63167 + + 3 AJ132100 3 Z46389 + 1 M59834 1 S76992 + + 1 D10656 W + 1 M29366 | 1 X92689 2 L24804 + + + 3 U57053 1 U94592 1 AC002310 1 AF070542 2 Z70223 1 AF040966 1 D79984 + + + 1 S73591 + + 1 S73591 1 S73591 1 S73591 1 X53461 + + 2 X99050 + + 4 X71490 + + 1 M63167 + + + 3 AJ132100 3 Z46389 + + + 1 M59834 1 S76992 + + 1 M29366 1 P13611 | 1 X92689 2 L24804 + + + + + 3 U57053 | 1 X92689 2 L24804 | 1 X92689 2 L24804 |

| WO 00/40/49 | | | | | | | | | |
|--|----|----------|---|---|---|----------|---|---|------------------------|
| vesicle-associated membrane protein 3 (cellubrevin) (VAMP3) | 1 | U64520 | | | | | | | |
| v-fos FBJ munne osteosarcoma viral oncogene homolog (FOS) | 26 | K00650 | | + | + | + | + | + | high in aorta |
| v-fos FBJ murine osteosarcoma viral oncogene homolog (FOS) (low match) | 1 | K00650 | | | | | | | |
| villin 2 (ezrin) (VIL2) | 1 | X51521 | + | + | + | + | | + | |
| villin-like protein | 1 | D88154 | | | | | | | |
| vimentin (VIM) | 12 | X56134 | | + | + | + | + | + | high in many libraries |
| vinculin (VCL) | 4 | M33308 | | + | + | + | | + | |
| vitamin A responsive; cytoskeleton related (JWA) | 6 | AF070523 | + | + | + | + | | + | |
| v-jun avian sarcoma virus 17 oncogene homolog (JUN) | 2 | U65928 | , | | | | | | |
| v-myb avian myeloblastosis viral oncogene homolog (MYB) | 1 | M15024 | | | + | | + | | |
| voltage-dependent anion channel 1 (VDAC1) | 1 | L06132 | + | + | + | + | | + | |
| voltage-dependent anion channel 3 (VDAC3) | 4 | U90943 | · | + | + | + | | + | |
| von Hippel-Lindau syndrome (VHL) | 1 | L15409 | | + | + | + | | + | |
| von Willebrand factor (vWF) (low matched) | 1 | X06828 | | | | <u> </u> | | | |
| v-raf murine sarcoma 3611 viral oncogene homolog 1 (ARAF1) | 2 | L24038 | + | + | + | + | | | |
| v-raf-1 murine leukemia viral oncogene homolog 1 (RAF1) | 1 | X03484 | + | + | + | + | | + | |
| v-ral simian leukemia viral oncogene homolog B (ras related; GTP binding protein) (RALB) | 3 | M35416 | | | | | | | |
| V-rel avian reticuloendotheliosis viral oncogene homolog A (nuclear factor of kappa light polypeptide gene enhancer in B-cells 3 (p65)) (RELA) | 1 | L19067 | | + | + | + | | + | |
| v-yes-1 Yamaguchi sarcoma viral related oncogene homolog (LYN) | 2 | M16038 | + | + | | + | | + | |
| WD repeat domain 1 (WDR1) | 1 | AB010427 | + | + | + | + | + | + | |
| WDR1 (=AF020260) | 1 | AF020056 | | | | | | | |
| WD-repeat protein (HAN11) | 2 | U94747 | | + | + | | | + | |
| Williams-Beuren syndrome chromosome region 1 (WBSCR1) | 12 | AF045555 | + | + | + | + | + | + | |
| Wiskott-Aldrich syndrome protein interacting protein (WASPIP) | 4 | X86019 | + | + | + | | | + | |
| X (inactive)-specific transcript (XIST) | 2 | M97168 | | | | | | | |
| xeroderma pigmentosum, complementation group C (XPC) | 3 | D21089 | + | + | + | + | | | |
| XIAP associated factor-1 | 2 | X99699 | - · · · · · · · · · · · · · · · · · · · | | | + | | | |
| XIB | 1 | X90392 | | + | + | | + | + | |
| X-linked anhidroitic ectodermal dysplasia | 1 | AF003528 | | | | | | | |

| WO 00/40/42 | | T T T T T T T T T T T T T T T T T T T | | | | | | | Ibiah ia aslasia |
|--|----|---------------------------------------|------|--------------|--|--|--|--|------------------|
| X-ray repair complementing defective | 1 | M30938 | + | + | + | + | | • | high in spleen |
| repair in Chinese hamster | | | | 1 | | | | | |
| cells 5 (double-strand- | | | | | | | | | |
| break | | | | | | | | | |
| rejoining; Ku autoantigen, | | | | 1 | | | | | |
| 80kD) (XRCC5) | | X TRASERA | | | ļ | | | | |
| XRP2 protein | 1 | AJ007590 | | <u> </u> | | | | | |
| yeloid differentiation | 1 | U84408 | | + | + | + | | + | |
| primary response gene | | | |] | l | | | | |
| (88) (MYD88) | 1 | L05148 | + | - | | + | | | |
| zeta-chain (TCR) associated protein kinase | , | 1 (03)40 | | 1 | | • | | | |
| (70kD) (ZAP70) | | : | | 1 | | | | | |
| zeta-chain (TCR) | 1 | L05148 | | 1 | | | | | |
| associated protein kinase | | | | 1 | | | | | l |
| (70kD) (ZAP70) (low | | | | 1 | | | | | - |
| match) | 2 | U69274 | + | + | + | + | - | + | |
| zinc finger protein (Hs.47371) | 2 | 003274 | • | 1 | , , | | İ | ' | |
| zinc finger protein | · | U69645 | + | + | + | + | | + | |
| (Hs.78765) | • | | | | | 1 | | | |
| zinc finger protein 10 (KOX | 1 | X78933 | | | | Ī | | | + only |
| 1) (ZNF10) | | ļ | | | | Щ | L | ļ | |
| ZÍNC FINGER PROTEIN | 1 | Q15973 | | 1 | | | | 1 | |
| 124 (HZF-16) (non-exact | | | | 1 | | | | 1 | ` |
| zinc finger protein 124 | 1 | S54641 | | +- | | | - | | |
| (HZF-16) (ZNF124) (non- | • | | | 1 | | | 1 | | |
| exact, 78%) | | | | 1 | | | | | |
| ZINC FINGER PROTEIN | 1 | P52736 | | | | | | | |
| 133 | | LIGOREZ | | - | + | | _ | | |
| zinc finger protein 136 (clone pHZ-20) (ZNF136) | 1 | U09367 | | | * | + | <u> </u> | ļ | 1 |
| zinc finger protein 140 | 1 | U09368 | | ++ | _ | + | \vdash | + | |
| (clone pHZ-39) (ZNF140) | • | | | ' | | | | | |
| zinc finger protein 140 | 1 | AF060865 | | | | | | _ | |
| (clone pHZ-39) (ZNF140) | |] | | 1 | | | | | |
| (non-exact 59%) | | | | ļ | | | ١ | L | ļ |
| zinc finger protein 140 | 1 | U09368 | | 1 | | : | | ŀ | |
| (clone pHZ-39) (ZNF140) (non-exact 73%) | | | | 1 | | | | | |
| zinc finger protein 140 | | S66508 | | + | | | | | |
| (clone pHZ-39) (ZNF140) | • | | | | | | | | |
| (non-exact 73%aa) | | | | <u> </u> | | | L | | |
| zinc finger protein 140 | 1 | U09368 | | | | | | | |
| (clone pHZ-39) (ZNF140) | | | | | | 1 | | | |
| (non-exact, 80%) zinc finger protein 143 | 2 | U09850 | + | + | + | + | + | + | |
| (clone pHZ-1) (ZNF143) | | 003030 | | ' | ' | ` | Ι . | 1 | |
| zinc finger protein 143 | 1 | U09850 | · | 1 | | | | | |
| (clone pHZ-1) (ZNF143) | | 1 | | . | | | | | |
| (low match) | | | | ļ | | ļ | <u> </u> | <u> </u> | |
| zinc finger protein 148 | 11 | AF039019 | + | } | | | | | |
| (pHZ-52) (ZNF148) | 1 | Q13105 | | | | - - | \vdash | | |
| ZINC FINGER PROTEIN 151 (MIZ-1 PROTEIN) (low | ' | 413105 | | | | | | | |
| match) | | 1 | | | 1 | İ | | ľ | |
| zinc finger protein 173 | 1 | U09825 | В, Т | + | + | | + | | |
| (ZNF173) | | L | | ļ | | | | <u> </u> | |
| zinc finger protein 192 | 1 | U57796 | | | | | | | |
| (ZNF192) (non-exact, 66%) | | A 1994004 | | + | + | + | | | <u> </u> |
| zinc finger protein 198 (ZNF198) | 1 | AJ224901 | | - | • |] ~ | i | |] |
| zinc finger protein 2 (ZNF2) | 1 | X60152 | | +- | \vdash | | | | |
| (low match) | ' | 7.00102 | | | | | ۱. | | |
| zinc finger protein 200 | 1 | AF060866 | | + | <u> </u> | + | | | |
| (ZNF200) | | | | <u></u> | | | <u></u> | _ | <u> </u> |
| zinc finger protein 207 | 6 | AF046001 | + | + | + | + | + | + | high in prostate |
| (ZNF207) | | AEDEODTO | L | ↓ | <u> </u> | | ├ | 1 | |
| zinc finger protein 216 | 2 | AF062072 | + | + | + | + | | + | |
| (ZNF216) | | <u> </u> | L | | L | Ь. | | Щ. | |

| zinc finger protein 217 (ZNF217) | 1 | AF041259 | Tactiv | ated - | | | | + | |
|--|---------------------------------------|----------|--------------|--------|---|---|----------|---|----------------------|
| ZINC FINGER PROTEIN 22 (ZINC FINGER PROTEIN KOX15) (non- | 1 | P17026 | | | | | | | |
| exact 58%) zinc finger protein 230 | | U95044 | | + | | | | - | |
| (ZNF230) Zinc finger protein 239 | 1 | L26914 | | + | | + | | | |
| (ANF239) zinc finger protein 261 | 1 | AB002383 | | + | + | + | | + | |
| (ZNF261) zinc finger protein 262 | · · · · · · · · · · · · · · · · · · · | AB007885 | | + | + | + | | + | |
| (ANF262) zinc finger protein 263 | · · · · · · | D88827 | | | | ļ | | | |
| (ZNF263) zinc finger protein 264 | · · | AB007872 | | + | + | + | | | |
| (ZNF264) ZINC FINGER PROTEIN | ' 1 | Q06730 | | | | | <u> </u> | | |
| 33A (ZINC FINGER PROTEIN KOX31) (KIAA0065) (HA0946) | ' | | | | _ | | | | |
| zinc finger protein 42 (myeloid-specific retinoic cid- responsive) (ZNF42) | 1 | M58297 | + | + | + | + | | + | |
| zinc finger protein 43 (HTF6) (ZNF43) (low match) | 1 | X59244 | | | | ! | | | |
| zinc finger protein 43 (HTF6) (ZNF43) (non- exact, 54%) | 1 | X59244 | | | | | | | |
| zinc finger protein 43 (HTF6) (ZNF43) (non- exact, 71%) | 1 | X59244 | | | | | | | |
| ZINC FINGER PROTEIN 43 (ZINC PROTEIN HTF6) (non-exact 67%) | 1 | P28160 | | | | | | | |
| zinc finger protein 45 (a Kruppel-associated box (KRAB) domain polypeptide) (ZNF45) | 1 | L75847 | | | | | | | only found in testis |
| ZINC FINGER PROTEIN 46 (ZINC FINGER PROTEIN KUP) (non-exact 62%) | 1 | P24278 | | | | | | | |
| zinc finger protein 6 (CMPX1) (ZNF6) | 1 | X56465 | | + | + | + | | + | |
| zinc finger protein 74 (Cos52) (ZNF74) (non- exact, 67%) | 1 | X71623 | | | | | | | |
| zinc finger protein 76 (expressed in testis) (ZNF76) | 1 | M91592 | | + | + | + | | + | |
| ZINC FINGER PROTEIN 83 (ZINC FINGER PROTEIN HPF1) (non- exact 65%) | 1 | P51522 | | | | | | | |
| zinc finger protein 84 (HPF2) (ZNF84) | 1 | M27878 | Tactivated | + | + | | | + | |
| zinc finger protein 85 (ZNF85)) | 2 | U35376 | | + | + | + | | | |
| zinc finger protein 9 (ZNF9) | 5 | M28372 | | + | + | + | + | + | |
| ZINC FINGER PROTEIN 93 (=ZINC FINGER PROTEIN HTF34) (non- exact 70%) | 1 | P35789 | | | | | | | |
| zinc finger protein C2H2-25 (ZNF25) | 3 | U38904 | | + | + | + | | | |
| zinc finger protein clone L3-4 | 1 | AF024706 | | | | | | | |
| zinc finger protein homologous to Zfp-36 in (mouse (ZFP36) | 4 | M92843 | + | | | | | | blood only |

| ZINC FINGER PROTEIN HRX (ALL-1) (71%a.a.) | 1 | Q03164 | | | | | | |
|--|---|--------|---|---|----------|--|---|------|
| zinc finger protein HZF4 | 1 | X78927 | | | <u> </u> | | | |
| zinc finger protein RIZ | 1 | D45132 | + | + | + | + | + | |
| zinc finger protein, subfamily 1A, 1 (Ikaros) (LYF1) | 1 | U40462 | + | | | | | |
| zinc finger protein, subfamily 1A, 1 (Ikaros) (LYF1) (low match) | 1 | U40462 | | | | | | |
| zinc finger transcriptional regulator (GOS24) | 1 | M92844 | | | | | | |
| zinc-finger helicase (hZFH) | 2 | U91543 | + | + | + | + | + | ï |
| Zn-15 related zinc finger protein (rff) | 1 | U22377 | | + | + | + | | |
| Zn-15 related zinc finger protein (rlf) (non-exact 56%) | 1 | U22377 | | | | | | |
| ZNF80-linked ERV9 long terminal repeat | 1 | X83497 | | | | | | |
| ZW10 (Drosophila) homolog, centromere/kinetochore protein (ZW10) | 2 | U54996 | | + | | | | |
| zyxin (ZYX) | 4 | X95735 | | | | | | |

Column 1: List of unique genes derived from 6,283 known ESTs from blood cells.

Column 2: Number of genes found in randomly sequenced ESTs from blood cells.

Column 3: Accession number. Column 4: "+" indicates the presence of the unique gene in publicly available cDNA libraries of blood (Bl), brain (Br), heart (H), kidney (K), liver (Li) and lung (Lu). **Comparison to previously identified tissue-specific genes was determined using the GenBank of the National Centre of Biotechnology Information (NCBI) Database.

10

15

5

Discussion

Every cell and tissue comprising the human body share the necessary genetic information required to maintain cellular homeostasis. These "housekeeping" genes function in basic cellular maintenance, including energy metabolism and cellular structure in all cell types. However, in certain situations, even the housekeeping genes show altered expression. Thus, it is necessary to define the use of these genes as internal controls from one investigation to another. Current results from the human blood cell EST database indicate that over 50% of the transcripts are

widely expressed throughout the human body. Most of the cell or tissue specific genes are also detectable in blood cells by RT-PCR analysis.

5

10

15

20

25

For example, isoformic myosin heavy chain genes are known to be generally expressed in cardiac muscle tissue. In the rodent, the βMyHC gene is only highly expressed in the fetus and in diseased states such as overt cardiac hypertrophy, heart failure and diabetes; the αMyHC gene is highly expressed shortly after birth and continues to be expressed in the adult heart. In the human, however, βMyHC is highly expressed in the ventricles from the fetal stage through adulthood. This highly expressed βMyHC, which harbours several mutations, has been demonstrated to be involved in familial hypertrophic cardiomyopathy (Geisterfer-Lowrance *et al.* 1990). It was reported that mutations of βMyHC can be detected by PCR using blood lymphocyte DNA (Ferrie et al., 1992). Most recently, it was also demonstrated that mutations of the myosin-binding protein C in familial hypertrophic cardiomyopathy can be detected in the DNA extracted from lymphocytes (Niimura *et al.*, 1998).

Similarly, APP and APC, which are known to be tissue specific and predominantly expressed in the brain and intestinal tract, are also detectable in the transcripts of blood. These cell- or tissue-specific transcripts are not detectable by Northern blot analysis. However, the low number of transcript copies can be detected by RT-PCR analysis. These findings strongly demonstrate that genes preferentially expressed in specific tissues can be detected by a highly sensitive RT-PCR assay. In recent years, evidence has been obtained to indicate that expression of cell or tissue-restricted genes can be detected in the peripheral blood of patients with metastatic transitional cell carcinoma (Yuasa et al. 1998) and patients with prostate cancer (Gala et al. 1998).

Atrial natriuretic factor (ANF) and zinc finger protein (ZFP), which are known to be highly expressed in heart tissue biopsies and in the plasma of heart failure patients, are also detectable in the transcripts of blood. Differential expression of zinc finger protein among the normal, diabetic and asymptomatic preclinical

subjects may have additional value as a prophylactic "early warning system". On a related note, there is now more attention/discussion in the cardiovascular disease field being focused on Syndrome X, loosely defined as a continuum of hypertension, increasing sugar levels, diabetes, kidney failure, culminating in heart failure, with the possibility of stroke and heart attack at any time in the continuum. The early identification of patients at risk of organ failure has been a challenge to the medical community for some time and the present method has the potential of resolving or, at least, ameliorating this challenge.

5

10

15

20

25

The present invention demonstrates that a simple drop of blood may be used to determine the quantitative expression of various mRNAs that reflect the health/disease state of the subject through the use of RT-PCR analysis. This entire process takes about three hours or less. The single drop of blood may also be used for multiple RT-PCR analyses. There is no need for large samples and/or costly and time-consuming separation of cell types within the blood for this method as compared to the methods described by Kimoto (1998) and Chelly et al. (1989; 1988). It is believed that the present finding can potentially revolutionize the way that diseases are detected, diagnosed and monitored because it provides a non-invasive, simple, highly sensitive and quick screening for tissue-specific transcripts. The transcripts detected in whole blood have potential as prognostic or diagnostic markers of disease, as they reflect disturbances in homeostasis in the human body. Delineation of the sequences and/or quantitation of the expression levels of these marker genes by RT-PCR will allow for an immediate and accurate diagnostic/prognostic test for disease or to assess the efficacy and monitor a particular therapeutic.

In addition to RT-PCR, other methods of amplifying may also be used for the purpose of measuring/quantitating tissue-specific transcripts in human blood. For example, mass spectrometry may be used to quantify the transcripts (Koster et al., 1996; Fu et al., 1998). The application of presently disclosed method for detecting tissue-specific transcripts in blood does not restrict to subjects undergoing course of

therapy or treatment, it may also be used for monitoring a patient for the onset of overt symptoms of a disease. Furthermore, the present method may be used for detecting any gene transcripts in blood. A kit for diagnosing, prognosing or even predicting a disease may be designed using gene-specific primers or probes derived from a whole blood sample for a specific disease and applied directly to a drop of blood. A cDNA library specific for a disease may be generated from whole blood samples and used for diagnosis, prognosis or even predicting a disease.

The following references were cited herein:

Claudio JO et al. (1998). Genomics 50:44-52.

5

10 Chelly J et al. (1989). Proc. Nat. Acad. Sci. USA. 86:2617-2621.

Chelly J et al. (1988). Nature 333:858-860.

Drews J & Ryser S (1997). Nature Biotech. 15:1318-9.

Ferrie RM et al. (1992). Am. J. Hum. Genet. 51:251-62.

Fu D-J et al. (1998). Nat. Biotech 16: 381-4.

15 Gala JL et al. (1998). Clin. Chem. 44(3):472-81.

Geisterfer-Lowrance AAT et al. (1990). Cell 62:999-1006.

Groden J et al. (1991). Cell 66:589-600.

Hwang DM et al. (1997). Circulation 96:4146-4203.

Jandreski MA & Liew CC (1987). Hum. Genet. 76:47-53.

20 Jin O et al. (1990). Circulation 82:8-16

Kimoto Y (1998). Mol. Gen. Genet 258:233-239.

Koster M et al. (1996). Nat. Biotech 14: 1123-8.

Liew & Jandreski (1986). Proc. Nat. Acad. Sci. USA. 83:3175-3179

Liew CC et al. (1990). Nucleic Acids Res. 18:3647-3651.

25 Liew CC (1993). J Mol. Cell. Cardiol. 25:891-894

Liew CC et al. (1994). Proc. Natl. Acad. Sci. USA. 91:10645-10649.

Liew et al. (1997). Mol. and Cell. Biochem. 172:81-87.

Niimura H et al. (1998). New Eng. J. Med. 338:1248-1257.

Ogawa M (1993). Blood 81:2844-2853.

5

10

15

Santoro IM & Groden J (1997). Cancer Res. 57:488-494.

Yuasa T et al. (1998). Japanese J. Cancer Res. 89:879-882.

Any patents or publications mentioned in this specification are indicative of the levels of those skilled in the art to which the invention pertains. Further, these patents and publications are incorporated by reference herein in their entirety to the same extent as if each individual publication was specifically and individually indicated to be incorporated by reference.

One skilled in the art will appreciate readily that the present invention is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those objects, ends and advantages inherent herein. The present examples, along with the methods, procedures, treatments, molecules, and specific compounds described herein are presently representative of preferred embodiments, are exemplary, and are not intended as limitations on the scope of the invention. Changes therein and other uses will occur to those skilled in the art which are encompassed within the spirit of the invention as defined by the scope of the claims.

WO 00/40749 WHAT IS CLAIMED IS:

5

- 1. A method for detecting expression of a gene in blood from a subject, comprising the steps of:
 - a) quantifying RNA from a subject blood sample; and
- b) detecting expression of said gene in the quantified RNA, wherein the expression of said gene in said quantified RNA indicates expression of said gene in the subject blood.
- 10 2. The method of claim 1, wherein the quantification is performed by mass spectrometry.
 - 3. A method for detecting expression of one or more genes in blood from a subject, comprising the steps of:
 - a) obtaining a subject blood sample;
 - b) extracting RNA from said blood sample;
 - c) amplifying said RNA;
 - d) generating expressed sequence tags from the amplified RNA product; and
- e) detecting expression of said genes in the expressed sequence tags, wherein the expression of said genes in said expressed sequence tags indicates expression of said genes in the subject blood.
- 4. The method of claim 3, wherein said genes are non-cancer-25 associated genes.
 - 5. The method of claim 3, wherein said genes are tissue-specific genes.

6. The method of claim 3, wherein said subject is a fetus, an embryo, a child, an adult or a non-human animal.

- 5 7. The method of claim 3, wherein the amplification is performed by RT-PCR.
- 8. The method of claim 7, wherein said RT-PCR utilizes primers selected from the group consisting of random sequence primers and gene-specific primers.
 - 9. A method for detecting expression of one or more genes in blood from a subject, comprising the steps of:
 - a) obtaining a subject blood sample;
 - b) extracting DNA fragment(s) from said blood sample;
 - c) amplifying said DNA fragment(s); and
 - d) detecting expression of said genes in the amplified DNA product, wherein the expression of said genes in said amplified DNA product indicates expression of said genes in the subject blood.

20

25

- 10. A method for monitoring a course of therapeutic treatment in an individual, comprising the steps of:
 - a) obtaining a blood sample from said individual;
 - b) extracting RNA from said blood sample;
 - c) amplifying said RNA;
- d) generating expressed sequence tags from the amplified RNA product; and

e) detecting expression of genes in said expressed sequence tags, wherein the expression of said genes is associated with the effect of said therapeutic treatment; and

- f) repeating steps a)-e), wherein the course of said therapeutic treatment is monitored by detecting the change of expression of said genes in the expressed sequence tags.
 - 11. The method of claim 10, wherein the amplification is performed by RT-PCR.

10

- 12. The method of claim 11, wherein the change of expression of said genes in the expressed sequence tags is monitored by sequencing the expressed sequence tags and comparing the resulting sequences at various time points.
- 13. The method of claim 11, wherein the change of expression of said genes in the expressed sequence tags is monitored by performing single nucleotide polymorphism analysis and detecting the variation of a single nucleotide in the expressed sequence tags at various time points.
- 20 14. The method of claim 10, wherein said individual is monitored for the onset of overt symptoms of a disease, and wherein the expression of said genes is associated with the onset of said symptoms.
- 15. A method for diagnosing a disease in a test subject, comprising the steps of:
 - a) generating a cDNA library for said disease from a whole blood sample from a normal subject;

b) generating expressed sequence tag (EST) profile from the normal subject cDNA library;

c) generating a cDNA library for said disease from a whole blood sample from a test subject;

5

15

20

- d) generating EST profile from the test subject cDNA library; and
- e) comparing the test subject EST profile to the normal subject EST profile, wherein if said test subject EST profile differs from said normal subject EST profile, said test subject might be diagnosed with said disease.
- 16. A kit for diagnosing, prognosing or predicting a disease, comprising:
 - a) gene-specific primers; wherein said primers are designed in such a way that the sequences of said primers contain the opposing ends of two adjacent exons for the specific gene with the intron sequence excluded; and
 - b) a carrier, wherein said carrier immobilizes said primer(s).
 - 17. The kit of claim 16, wherein said gene-specific primer(s) are selected from the group consisting of insulin-specific primers, atrial natriuretic factor-specific primers, zinc finger protein gene-specific primers, beta-myosin heavy chain gene-specific primers, amyloid precurser protein gene-specific primers, and adenomatous polyposis-coli protein gene-specific primers.
 - 18. The kit of claim 17, wherein the sequences of said genespecific primers are selected from the group consisting of SEQ ID Nos. 1 and 2, and SEQ ID Nos. 5 and 6.
 - 19. A method for diagnosing, prognosing or predicting a disease in a test subject, comprising the step of:

PCT/CA00/00005 WO 00/40749

applying the kit of claim 16 to a test subject whole blood sample, wherein quantitative expression levels of specific genes associated with said disease are detected and compared to the levels of said specific genes expressed in a normal subject, therefore, said disease may be diagnosed, prognosed or predicted.

5

The method of claim 19. wherein said method is used for 20. monitoring a course of therapeutic treatment or monitoring the onset of overt symptoms of said disease.

10 comprising: 21.

- probes derived from a whole blood sample for a specific a) disease; and
 - a carrier, wherein said carrier immobilizes said probes. b)

A kit for diagnosing, prognosing or predicting a disease,

15


20

25

A method for diagnosing, prognosing or predicting a disease in 22. a test subject, comprising the step of:

applying the kit of claim 21 to a test subject whole blood sample, wherein quantitative expression levels of specific genes associated with said disease are detected and compared to the levels of said specific genes expressed in a normal subject, therefore, said disease may be diagnosed, prognosed or predicted.

- The method of claim 22, wherein said method is used for 23. monitoring a course of therapeutic treatment or monitoring the onset of overt symptoms of said disease.
- A cDNA library specific for a disease, wherein said cDNA 24. library is generated from whole blood samples.







FIGURE 1 1/7



2/7
SUBSTITUTE SHEET (RULE 26)




FIGURE 3 3/7






FIGURE 4 4/7








FIGURE 5

5/7




FIGURE 6 6/7


Total: 13,283 ESTs Known: 6,283

Mitochondrial: 405

Ribosome: 498 Repeat: 868 Mis.: 156 Novel: 2,718

Human Blood

Human Fetal Heart



■5% **■**8%

■22%

%9□

Metabolism

%9**■**

■ Unclassified

□26%

■29%




FIGURE 7

7/7

SEQUENCE LISTING

| <: | 110> | Liew, Choong-Chin |
|----------|-------------------|--|
| | 120> | Method for the Detection of Gene Transcripts |
| in Bloo | od and Us | ses Thereof |
| <u>,</u> | 130> | 2173/0003 |
| • | 130> | 2173, 0003 |
| <: | 150> | US Number not yet assigned |
| < | 151> | 2000-01-04 |
| | | |
| <: | 150> | US 60/115,125 |
| < | 151> | 1999-01-06 |
| <. | 160> | 10 |
| | | |
| < | 210> | 1 |
| <. | 211> | 18 |
| < | 212> | DNA |
| < | 213> | artificial sequence |
| < | 220> | |
| < | 221> | primer_bind |
| | 223> antitati | forward primer of exon 1 of insulin gene used ve RT-PCR analysis |
| < | 400> | 1 |
| gccctc | tggg gac | ctgac 18 |
| | | |
| < | 210> | 2 |
| <. | 211> | 18 |
| < | 212> | DNA |
| < | 213> | artificial sequence |
| < | 220> | |
| < | 221> | primer_bind |
| gene u | 223> sed for (| reverse primer of exons 1 and 2 of insulin quantitative RT-PCR analysis |
| < | 400> | 2 |
| cccacc | tgca ggt | ectct 18 |

| WO 00/40749 | | PCT/CA00/00005 |
|----------------------|--|---------------------|
| <210> | 3 | |
| <211> | 24 | |
| <212> | DNA | |
| <213> | artificial sequence | |
| <220> | | |
| <221> | <pre>primer_bind</pre> | |
| | forward primer of β MyH α | C gene used for |
| <400> | 3 | |
| gctggaacgt | agagactccc tgct | 24 |
| | | |
| <210> | 4 | |
| <211> | 24 | |
| <212> | DNA | |
| <213> | artificial sequence | |
| <220> | | |
| <221> | <pre>primer_bind</pre> | |
| <223> quantitativ | reverse primer of eta MyH $_{\!$ | C gene used for |
| <400> | 4 | |
| ggatccttcc | agatcatcca cttg | 24 |
| | | |
| <210> | 5 | |
| <211> | 20 | |
| <212> | DNA | |
| <213> | artificial sequence | |
| <220> | | |
| <221> | <pre>primer_bind</pre> | |
| <223> RT-PCR | forward primer of ANF use | ed for quantitative |
| | analysis | |
| <400> | 5 | |
| | aatttgctgg | 20 |

| WO 00/40749 |) | PCT/CA00/00005 |
|----------------------|--------------------------------|----------------|
| <210> | 6 | |
| <211> | 20 | |
| <212> | DNA | |
| <213> | artificial sequence | |
| <220> | | |
| <221> | primer_bind | |
| <223> RT-PCR anal | | r quantitative |
| <400> | 6 | |
| gcagatcgat | cagaggagtc | 20 |
| <210> | 7 | |
| <211> | 20 | |
| <212> | DNA | |
| <213> | artificial sequence | |
| <220> | | |
| <221> | <pre>primer_bind</pre> | |
| <223> RT-PCR | forward primer of APP used for | r quantitative |
| | analysis | |
| <400> | 7 | |
| ggatgcttca | tgtgaacgtg | 20 |
| | | |
| <210> | 8 | |
| <211> | 19 | |
| <212> | DNA | |
| <213> | artificial sequence | |
| <220> | | |
| <221> | - | |
| <223> RT-PCR | reverse primer of APP used fo | r quantitative |
| | analysis | |
| <400> | 8 | |
| tcattcacac | cagcacatg | 19 |

| WO 00/40749 | PCT/CA00/0000 | 5 |
|------------------------|---|---|
| <210> | 9 | |
| <211> | 21 | |
| <212> | DNA | |
| <213> | artificial sequence | |
| <220> | | |
| <221> | primer_bind | |
| <223> RT-PCR analys | <u>-</u> | |
| <400> | 9 | |
| cacargagre ar | ggtcaacg a 21 | |
| | | |
| <210> | 10 | |
| <211> | 22 | |
| <212> | DNA | |
| <213> | artificial sequence | |
| <220> | | |
| <221> | primer_bind | |
| <223> RT-PCR analys | reverse primer of ZFP used for quantitative | |
| <400> | 10 | |
| ggattaaaat ga | agcaccca ga 22 | |

Inter Inal Application No PCT/CA 00/00005

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, MEDLINE, CHEM ABS Data, BIOSIS, EMBASE, EMBL

| Category ° | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|--|-----------------------|
| X | WO 98 49342 A (COCKBAIN JULIAN R M ;FORSKNINGSPARKEN I AAS AS (NO); LOENNEBORG AN) 5 November 1998 (1998-11-05) | 1-15, 21-24 |
| Υ | the whole document | 19,20 |
| X | WO 98 24935 A (AN GANG ;HARA MARK O (US); RALPH DAVID (US); VELTRI ROBERT (US); U) 11 June 1998 (1998-06-11) | 1-15, 21-24 |
| Y | the whole document | 19,20 |
| X | EP 0 534 640 A (PFIZER) 31 March 1993 (1993-03-31) | 16 |
| Υ | the whole document | 17-20 |
| | -/ | |
| | | |
| | | |

| X Further documents are listed in the continuation of box C. | X Patent family members are listed in annex. |
|--|---|
| Special categories of cited documents: A* document defining the general state of the art which is not considered to be of particular relevance E* earlier document but published on or after the international filling date L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) O* document referring to an oral disclosure, use, exhibition or other means P* document published prior to the international filling date but later than the priority date claimed | "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention. "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone. "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family |
| Date of the actual completion of the international search | ☐ate of mailing of the international search report |
| 27 June 2000 | 12/07/2000 |
| Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentiaan 2 NL – 2280 HV Rijswijk Tel. (+31-70) 340–2040, Tx. 31 651 epo nl, Fax: (+31-70) 340–3016 | Authorized officer Hagenmaier, S |

2

Inter anal Application No PCT/CA 00/00005

| Continu | ation) DOCUMENTS CONSIDERED TO BE RELEVANT | 1/CA 00/00005 |
|------------|--|----------------------|
| Category * | | Relevant to daim No. |
| Y | PATENT ABSTRACTS OF JAPAN vol. 1997, no. 11, 28 November 1997 (1997-11-28) & JP 09 187299 A (NIPPON BIO SERAPII KK), 22 July 1997 (1997-07-22) abstract & DATABASE GCG_GENESEQ 'Online! AC:V00198, March 1998 (1998-03) NIPPON BIOTHERAPY: "HUMAN INSULIN SENSE PRIMER INS-7" abstract | 17,18 |
| Y | DATABASE GENBANK 'Online! AC:V00565, March 1995 (1995-03) BELL ET AL.: "HUMAN GENE FOR PREPROINSULIN" XP002141055 abstract | 17,18 |
| Y | DATABASE GENBANK 'Online! AC:M54947, April 1993 (1993-04) SEIDMAN ET AL.: "HUMAN ATRIAL NATRIURETIC FACTOR GENE" XP002141054 abstract | 17,18 |
| Y | DATABASE GENBANK 'Online! AC:X52889, September 1993 (1993-09) LIEW: "HUMAN GENE FOR CARDIAC BETA MYOSIN HEAVY CHAIN" XP002141056 abstract | 17 |
| Υ | DATABASE GENBANK 'Online! AC:2808656, December 1998 (1998-12) BERNOT ET AL.: "A TRANSCRIPTIONAL MAP OF THE FMF REGION/ZINC FINGER PROTEIN" XP002141057 abstract | |
| Υ | YOSHIKAI ET AL.: "GENOMIC ORGANIZATION OF THE HUMAN AMYLOID BETA-PROTEIN PRECURSOR GENE" GENE, vol. 87, 1990, pages 257-263, XP002141053 the whole document | 17 |
| Y | DATABASE GENBANK 'Online! AC:M73548, January 1995 (1995-01) JOSLYN ET AL.: "HUMAN POLYPOSIS LOCUS mRNA" XP002141058 abstract | 17 |
| | -/ | |

2

Inter anal Application No
PCT/CA 00/00005

| | | PCI/CA 00 | |
|------------|---|-----------|-----------------------|
| | ation) DOCUMENTS CONSIDERED TO BE RELEVANT | | Relevant to claim No. |
| Category ° | Citation of document, with indication, where appropriate, of the relevant passages | | neievani to daim No. |
| A | WO 98 18906 A (CLAYCOMB WILLIAM CREIGHTON; PROCTER & GAMBLE (US)) 7 May 1998 (1998-05-07) the whole document | | |
| A | US 5 352 775 A (MARKHAM ALEXANDER F ET AL) 4 October 1994 (1994-10-04) the whole document | | |
| A | DE 44 35 919 C (DEUTSCHES KREBSFORSCH) 7 December 1995 (1995-12-07) the whole document | | |
| A | US 5 837 449 A (ECKER DAVID J ET AL) 17 November 1998 (1998-11-17) the whole document | | |
| A | WO 98 33942 A (NAKAO KOICHI ;BRISTOW MICHAEL R (US); LEINWAND LESLIE A (US); MINO) 6 August 1998 (1998-08-06) the whole document | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | , | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | <u> </u> |

2

information on patent family members

Inter onal Application No PCT/CA 00/00005

| | tent document in search report | | Publication date | | Patent family member(s) | Publication date |
|----|---|-------|---------------------|------|-------------------------|-------------------------|
| WO | 9849342 | A | 05-11-1998 | AU | 7222698 A | 24-11-1998 |
| | | | | EP | 0979308 A | 16-02-2000 |
| | | | | NO | 995296 A | 14-12-1999 |
| WO | 9824935 | Α | 1·1-06-1998 | AU | 5515198 A | 29-06-1998 |
| | | | | EP | 0960214 A | 01-12-1999 |
| EP | 0534640 | Α | 31-03-1993 | AT | 143700 T | 15-10-1996 |
| | • | | | CA | 2078703 A | 24-03-1993 |
| | | | | DE | 69214243 D | 07-11-1996 |
| | | | | DE | 69214243 T | 06 - 02-1997 |
| | | | | DK | 534640 T | 17-03-1997 |
| | | | | ES | 2092056 T | 16-11-1996 |
| | | | | FΙ | 924242 A | 24-03-1993 |
| | | | | GR · | 3021721 T | 28-02-1997 |
| | | | | JP | 2703156 B | 26-01-1998 |
| | | | | JP | 5192199 A | 03-08-1993 |
| | | | | US | 5643730 A | 01-07-1997 |
| JP | 09187299 | Α | 22-07-1997 | NONE | | |
| WO | 9818906 | А | 07-05-1998 | AU | 5149998 A | 22-05-1998 |
| | • | | | EP | 0956341 A | 17-11-1999 |
| | | | | NO | 991894 A | 24-06-1999 |
| US | 5352775 | A | 04-10-1994 | US | 5648212 A | 15-07-1997 |
| | | | | US | 5691454 A | 25-11-1997 |
| | | | | US | 5783666 A | 21-07-1998 |
| | | | | US | RE36713 E | 23-05-2000 |
| | | | | AU | 1366992 A | 27-08-1992 |
| | | | | EP | 0569527 A | 18-11-1993 |
| | | | | WO | 9213103 A | 06-08-1992 |
| DE | 4435919 | С | 07-12-1995 | WO | 9611267 A | 18-04-1996 |
| | | | | EP | 0784680 A | 23-07-1997 |
| | | | | JP | 10506789 T | 07-07-1998 |
| US | 5837449 | Α | 17-11-1998 | AU | 3249793 A | 28-07-1993 |
| | | | | .CA | 2126451 A | 08-07-1993 |
| | | | | EP | 0644889 A | 29-03-1995 |
| | | | | JP | 6511387 T | 22-12-1994 |
| | | | | WO | 9313114 A | 08-07-1993 |
| MU | 9833942 | A | 06-08-1998 | AU | 6141098 A | 25-08-1998 |